

**OCR**

Oxford Cambridge and RSA

**...day June 20XX – Morning/Afternoon**

**A Level Chemistry B (Salters)**

**H433/01 Fundamentals of chemistry**

**SAMPLE MARK SCHEME**

**Duration:** 2 hours 15 minutes

**MAXIMUM MARK 110**

**This document consists of 24 pages**

## MARKING INSTRUCTIONS

### PREPARATION FOR MARKING

#### SCORIS

1. Make sure that you have accessed and completed the relevant training packages for on-screen marking: *scoris assessor Online Training*; *OCR Essential Guide to Marking*.
2. Make sure that you have read and understood the mark scheme and the question paper for this unit. These are posted on the RM Cambridge Assessment Support Portal <http://www.rm.com/support/ca>
3. Log-in to scoris and mark the **required number** of practice responses (“scripts”) and the **required number** of standardisation responses.

YOU MUST MARK 10 PRACTICE AND 10 STANDARDISATION RESPONSES BEFORE YOU CAN BE APPROVED TO MARK LIVE SCRIPTS.

#### MARKING

1. Mark strictly to the mark scheme.
2. Marks awarded must relate directly to the marking criteria.
3. The schedule of dates is very important. It is essential that you meet the scoris 50% and 100% (traditional 50% Batch 1 and 100% Batch 2) deadlines. If you experience problems, you must contact your Team Leader (Supervisor) without delay.
4. If you are in any doubt about applying the mark scheme, consult your Team Leader by telephone, email or via the scoris messaging system.

5. Work crossed out:
- where a candidate crosses out an answer and provides an alternative response, the crossed out response is not marked and gains no marks
  - if a candidate crosses out an answer to a whole question and makes no second attempt, and if the inclusion of the answer does not cause a rubric infringement, the assessor should attempt to mark the crossed out answer and award marks appropriately.
6. Always check the pages (and additional objects if present) at the end of the response in case any answers have been continued there. If the candidate has continued an answer there then add a tick to confirm that the work has been seen.
7. There is a NR (No Response) option. Award NR (No Response)
- if there is nothing written at all in the answer space
  - OR if there is a comment which does not in any way relate to the question (e.g. 'can't do', 'don't know')
  - OR if there is a mark (e.g. a dash, a question mark) which isn't an attempt at the question.
- Note: Award 0 marks – for an attempt that earns no credit (including copying out the question).
8. The scoris **comments box** is used by your Team Leader to explain the marking of the practice responses. Please refer to these comments when checking your practice responses. **Do not use the comments box for any other reason.**
- If you have any questions or comments for your Team Leader, use the phone, the scoris messaging system, or email.
9. Assistant Examiners will send a brief report on the performance of candidates to their Team Leader (Supervisor) via email by the end of the marking period. The report should contain notes on particular strengths displayed as well as common errors or weaknesses. Constructive criticism of the question paper/mark scheme is also appreciated.

10. For answers marked by levels of response:

Read through the whole answer from start to finish, concentrating on features that make it a stronger or weaker answer using the indicative scientific content as guidance. The indicative scientific content indicates the expected parameters for candidates' answers, but be prepared to recognise and credit unexpected approaches where they show relevance.

Using a 'best-fit' approach based on the science content of the answer, first decide which set of level descriptors, Level 1, Level 2 or Level 3, **best** describes the overall quality of the answer using the guidelines described in the level descriptors in the mark scheme.

Once the level is located, award the higher or lower mark.

**The higher mark** should be awarded where the level descriptor has been evidenced and all aspects of the communication statement (in italics) have been met.

**The lower mark** should be awarded where the level descriptor has been evidenced but aspects of the communication statement (in italics) are missing.

**In summary:**

- **The science content determines the level.**
- **The communication statement determines the mark within a level.**

Level of response questions on this paper are **32(b)** and **34(b)**.

## 11. Annotations

<b>Annotation</b>	<b>Meaning</b>
<b>DO NOT ALLOW</b>	Answers which are not worthy of credit
<b>IGNORE</b>	Statements which are irrelevant
<b>ALLOW</b>	Answers that can be accepted
( )	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument
✓	Marking point

12. **Subject-specific Marking Instructions**

**INTRODUCTION**

Your first task as an Examiner is to become thoroughly familiar with the material on which the examination depends. This material includes:

- the specification, especially the assessment objectives
- the question paper
- the mark scheme.

You should ensure that you have copies of these materials.

You should ensure also that you are familiar with the administrative procedures related to the marking process. These are set out in the OCR booklet **Instructions for Examiners**. If you are examining for the first time, please read carefully **Appendix 5 Introduction to Script Marking: Notes for New Examiners**.

Please ask for help or guidance whenever you need it. Your first point of contact is your Team Leader.

## SECTION A

Question	Answer	Marks	Guidance
1	D	1	
2	D	1	
3	D	1	
4	B	1	
5	A	1	
6	A	1	
7	D	1	
8	B	1	
9	B	1	
10	C	1	
11	C	1	
12	B	1	
13	D	1	
14	A	1	
15	B	1	
16	C	1	
17	A	1	
18	B	1	
19	B	1	
20	A	1	

Question	Answer	Marks	Guidance
21	B	1	
22	B	1	
23	D	1	
24	C	1	
25	A	1	
26	A	1	
27	B	1	
28	B	1	
29	A	1	
30	D	1	
	Total	30	

## SECTION B

Question			Answer	Marks	Guidance
31	(a)	(i)	Cerium(IV) oxide ✓	1	<b>ALLOW</b> without parentheses
		(ii)	$4\text{CeO}_2 \rightarrow 2\text{Ce}_2\text{O}_3 + \text{O}_2$ <b>OR</b> $2\text{CeO}_2 \rightarrow \text{Ce}_2\text{O}_3 + 0.5\text{O}_2$ ✓ $\text{Ce}_2\text{O}_3 + \text{H}_2\text{O} \rightarrow 2\text{CeO}_2 + \text{H}_2$ ✓	2	
		(iii)	Enthalpy changes of formation ✓	1	<b>ALLOW</b> bond enthalpies/energies
	(b)	(i)	$E^\ominus$ of $\text{Ce}^{4+}/\text{Ce}^{3+}$ in $\text{H}_2\text{SO}_4(\text{aq})$ is more positive than $E^\ominus$ of $\text{Cl}_2/\text{Cl}^-$ ✓  So will oxidise $\text{Cl}^-$ to $\text{Cl}_2$ ✓  In $\text{HCl}(\text{aq})$ $E^\ominus$ of $\text{Ce}^{4+}/\text{Ce}^{3+}$ is less positive so will not oxidise $\text{Cl}^-$ to $\text{Cl}_2$ <b>AND</b> so $\text{H}_2\text{SO}_4(\text{aq})$ is used ✓	3	<b>ALLOW</b> less negative   <b>ALLOW</b> more negative
		(ii)	<b>FIRST CHECK ANSWER ON ANSWER LINE</b> answer = 83% to 2 sig figs award 5 marks  $n(\text{AsO}_3^{3-})$ used in titration = $3.00/1000 \times 0.500$ <b>OR</b> 0.00150 (mol) (conc. of $\text{AsO}_3^{3-}$ is double that of $\text{As}_2\text{O}_3$ ) ✓  $n(\text{Ce}^{4+})$ used in titration = $2 \times (3.00/1000 \times 0.500)$ <b>OR</b> $2 \times 0.00150$ <b>OR</b> 0.00300 (mol) ✓  $M_r$ of $\text{CeO}_2$ = 172.1 ✓  mass $\text{CeO}_2$ present in 100 $\text{cm}^3$ of solution = $4 \times 172.1 \times 0.00300$ = 2.0652 g ✓  % purity of $\text{CeO}_2$ sample = $2.0652 \times 100/2.5$ = 83% to 2 sig figs ✓	5	<b>ALLOW ECF</b> from first marking point          <b>ALLOW ECF</b> from second marking point

Question	Answer	Marks	Guidance
	<p>(iii) Use a more dilute <math>\text{AsO}_3^{3-}</math> solution to increase volume of titre / reduce % error in titre</p> <p><b>OR</b></p> <p>Use more ceria sample to increase volume of titre / reduce % error in titre ✓</p>	1	
	<p>(iv) <b>FIRST CHECK ANSWER ON ANSWER LINE</b>            answer = <math>1.7 \times 10^{22}</math> atoms of oxygen award 2 marks</p> <p><math>n(\text{CeO}_2) = 2.5/172.1 = 0.01453</math> (mol) ✓</p> <p><math>n(\text{O}) = 0.01453 \times 2 = 0.02905</math> (mol)</p> <p><math>0.02905 \times 6.02 \times 10^{23} = 1.7 \times 10^{22}</math> atoms of oxygen ✓</p>	2	<p><b>ALLOW ECF</b> from first marking point  <b>ALLOW 2 or more sig figs</b></p>
	<p>(v) Reactants adsorbed onto surface of catalyst and form bonds to surface (<b>AW</b>) ✓</p> <p>Bonds within reactants weaken and break ✓</p> <p>New bonds form (<b>AW</b>) ✓</p> <p>Products formed desorb/leave from catalyst (<b>AW</b>) ✓</p>	4	
	<b>Total</b>	<b>19</b>	

Question		Answer	Marks	Guidance
32	(a)	<p><i>Two marking points from the following:</i></p> <ul style="list-style-type: none"> <li>• Large amounts of arable land are required to produce the crops required to obtain large amounts ethanol</li> <li>• (Environmental problem caused by) disposal of fermentation waste</li> <li>• Current car engines need to be modified to use high concentrations of ethanol</li> </ul> <p>✓ ✓</p>	2	<p><b>ALLOW</b> ethanol has a lower enthalpy change of combustion than petrol</p> <p><b>IGNORE</b> better for the environment</p>


Question	Answer	Marks	Guidance
(b)*	<p><i>Please refer to the marking instructions on page 4 of this mark scheme for guidance on how to mark this question.</i></p> <p><b>Level 3 (5–6 marks)</b> Analyses information <b>AND</b> spectral data to provide evidence to support the correct and full identification of all compounds <b>A, B, C and D</b>. Evidence from reactions of <b>A AND</b> no reaction of <b>B</b> with Tollens' reagent or NaOH <b>AND</b> MS spectrum <b>AND</b> <math>^{13}\text{C}</math> NMR spectrum.</p> <p><i>The information and evidence used is relevant and fully supports the identification. The answer is clear and logically structured.</i></p> <p><b>Level 2 (3–4 marks)</b> Analyses information <b>AND</b> data to provide evidence to support the partial identification of compound <b>A</b> as a secondary alcohol, <b>B</b> as a ketone and <b>C</b> and <b>D</b> as alkenes. Evidence from reactions of <b>A AND</b> no reaction of <b>B</b> with Tollens' reagent or NaOH <b>AND EITHER</b> MS spectrum <b>OR</b> <math>^{13}\text{C}</math> NMR spectrum</p> <p><i>The information and evidence used is in the most-part relevant and supports the identification. The answer is presented with some structure.</i></p> <p><b>Level 1 (1–2 marks)</b> Analyses information <b>OR</b> data to provide evidence allowing partial identification of the compounds <b>A AND B OR C AND D</b> using reactions of <b>A OR</b> no reaction of <b>B</b> with Tollens' reagent or NaOH <b>OR</b> using information from MS Spectrum <b>OR</b> <math>^{13}\text{C}</math> NMR spectrum</p>	6	<p><b>Indicative scientific points may include:</b></p> <p>Full identification Compound <b>A</b> is <math>\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3</math> Compound <b>B</b> is <math>\text{CH}_3\text{CH}_2\text{COCH}_3</math> Compounds <b>C &amp; D</b> are <math>\text{CH}_3\text{CH}_2\text{CH}=\text{CH}_2</math> and <math>\text{CH}_3\text{CH}=\text{CHCH}_3</math></p> <p><b>Evidence from spectral data</b></p> <p><b>MS Spectrum:</b></p> <ul style="list-style-type: none"> <li><math>M_r (\text{C}_x\text{H}_y\text{O}) = 74</math> <math>M_r (\text{C}_x\text{H}_y) 74 - 16 = 58</math> so <math>x = 4</math> and <math>y = 10</math>.</li> </ul> <p><math>^{13}\text{C}</math> NMR: 2 from</p> <ul style="list-style-type: none"> <li>4 carbon environments</li> <li>no C=O or C=C</li> <li>C–O (and C–C) present.</li> </ul> <p><b>Evidence from the Reactions:</b></p> <ul style="list-style-type: none"> <li><b>A</b> is alcohol from formula plus <math>\text{H}^+/\text{Cr}_2\text{O}_7^{2-}</math> reaction</li> <li>Heating <b>A</b> with <math>\text{Al}_2\text{O}_3</math> results in elimination of water from <b>A</b> forms 2 different alkenes, <b>C</b> and <b>D</b></li> <li>Thus <b>A</b> secondary and <b>B</b> a ketone.</li> <li><b>A</b> reacts with <math>\text{H}^+/\text{Cr}_2\text{O}_7^{2-}</math> when heated <math>\rightarrow</math> <b>B</b> is aldehyde, ketone or carboxylic acid</li> <li>No reaction with Tollens' <math>\rightarrow</math> <b>B</b> is <b>NOT</b> an aldehyde</li> <li>No reaction with NaOH <math>\rightarrow</math> <b>B</b> is <b>NOT</b> a carboxylic acid</li> <li>Conclusion: <b>B</b> is a ketone <b>AND</b> <b>A</b> a secondary alcohol.</li> </ul>

Question	Answer	Marks	Guidance
	<p><i>The information and evidence is used to make a partial identification of <b>A AND B OR C</b> and <b>D</b>. The evidence chosen does not fully support the identification and is not presented in a logical order.</i></p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b> No response or no response worthy of credit.</p>		For Level 1: partial identification of <b>A</b> required. May be supplemented by partial identification of <b>B OR</b> partial identification of <b>C</b> and <b>D</b> .
(c)	Substrate/reactant has specific <u>shape</u> ✓ Fits <u>active site</u> in enzyme ✓	<b>2</b>	
(d)	$n(\text{O}_2) = 25 \times 0.21/24.0 = 0.219 \text{ mol}$ ✓ $M_r$ of $\text{C}_2\text{H}_5\text{OH} = 46.0$ $n(\text{C}_2\text{H}_5\text{OH}) = 4.0/46.0 = 0.087 \text{ mol}$ Recognition of ratio $\text{O}_2 / \text{C}_2\text{H}_5\text{OH}$ : $0.219 / 0.087 = 2.5$ ✓ This is smaller than the required ratio of 3 (from $\text{C}_2\text{H}_5\text{OH} + 3\text{O}_2 \rightarrow 2\text{CO}_2 + 3\text{H}_2\text{O}$ ), so the ethanol is not completely burned. ✓	<b>3</b>	<b>ALLOW ECF</b> from first marking point.
	<b>Total</b>	<b>13</b>	

Question		Answer	Marks	Guidance	
33	(a)	$\begin{array}{c} \text{CH}_2\text{OOC}(\text{CH}_2)_{10}\text{CH}_3 \\   \\ \text{CHOOC}(\text{CH}_2)_{10}\text{CH}_3 \\   \\ \text{CH}_2\text{OOC}(\text{CH}_2)_{10}\text{CH}_3 \end{array}$ <p>Any of the ester groups circled correctly ✓</p>	1	<b>IGNORE</b> circled adjacent carbons	
	(b)	(i)	$K_{\text{sp}} = [\text{Na}^+(\text{aq})] \times [\text{L}^-(\text{aq})]$ ✓	1	State symbols required <b>ALLOW</b> $K_{\text{sp}} = [\text{Na}^+(\text{aq})] [\text{L}^-(\text{aq})]$
		(ii)	$[\text{Na}^+]$ increases but $K_{\text{sp}}$ remains constant ✓ so NaL precipitates to make $[\text{L}^-]$ smaller / to move equilibrium left ✓	2	
		(iii)	<p><b>FIRST CHECK ANSWER ON ANSWER LINE</b> minimum mass of NaCl = 34.2 g award 4 marks</p> <p>Solubility of F = <math>24.0/222 = 0.108 \text{ mol dm}^{-3}</math></p> <p><math>K_{\text{sp}} = (\text{solubility})^2 = 0.0117</math> ✓</p> <p><math>0.0117 = [\text{Na}^+(\text{aq})] \times 1.0 \times 10^{-2}</math> ✓</p> <p><math>[\text{NaCl}]</math> to exceed <math>K_{\text{sp}} = 0.0117 / 1.0 \times 10^{-2} = 1.17 \text{ (mol dm}^{-3}\text{)}</math> ✓</p> <p>Min mass of NaCl to add to <math>500 \text{ cm}^3</math> to form ppt of F = <math>(1.17/2) \times 58.5 = 34.2 \text{ g}</math> ✓</p> <p><b>Assumption:</b> volume of solution does not change when NaCl added ✓</p>	5	<b>IGNORE</b> units

Question	Answer	Marks	Guidance
(c)	<p><b>FIRST CHECK ANSWER ON ANSWER LINE</b>            Answer = 0.043 award 4 marks</p> <p>concentration of H<sub>2</sub>O = 5.00 – 1.20 = 3.80 (mol dm<sup>-3</sup>) ✓</p> <p>concentration of acid H = 0.60 (mol dm<sup>-3</sup>)  <b>AND</b>            concentration of CH<sub>3</sub>OH = 1.20 (mol dm<sup>-3</sup>) ✓</p> $K_c = \frac{[(\text{CH}_2\text{COOH})_2][\text{CH}_3\text{OH}]^2}{[(\text{CH}_2\text{COOCH}_3)_2][\text{H}_2\text{O}]^2} = \frac{0.6 \times 1.2^2}{1.4 \times 3.8^2} \checkmark$ <p>= 0.043 ✓ (no units)</p>	4	<p><b>ALLOW</b> 2 or more sig figs</p>
(d)	<p><b>FIRST CHECK ANSWER ON ANSWER LINE</b>            Answer = 22.5 g award 4 marks</p> <p><math>[\text{H}^+] = 10^{-4.8} = 1.585 \times 10^{-5} \text{ mol dm}^{-3} \checkmark</math></p> $K_a = \frac{[\text{H}^+][\text{salt}]}{[\text{acid}]} \text{ AND } \frac{[\text{salt}]}{[\text{acid}]} = \frac{K_a}{[\text{H}^+]} \checkmark$ $= \frac{1.74 \times 10^{-5}}{1.585 \times 10^{-5}} = 1.0979 \checkmark$ <p>Therefore amount of ethanoate required            = 1.0979 × n(ethanoic acid)            = 1.0979 × 0.250 = 0.274 (mol)</p> <p>0.274 × 82.0 = 22.5 (g) ✓</p>	4	<p><b>ALLOW ECF</b> from first marking point</p> <p><b>ALLOW ECF</b> from third marking point</p>
	<b>Total</b>	<b>17</b>	

Question		Answer	Marks	Guidance
34	(a)	Only small amount of raspberry ketone present in raspberries / difficult / high cost to extract ✓	1	
	(b)*	<p><i>Please refer to the marking instructions on page 4 of this mark scheme for guidance on how to mark this question.</i></p> <p><b>Level 3 (5–6 marks)</b> Fully describes the differences between the two compounds in detail with both ring descriptions correct. Describes at least three reactions with examples and equations.</p> <p><i>The full description is detailed and correct. There is a clear and logical structure. The reactions are relevant and fully supported with examples and equations. Demonstrates a clear and confident knowledge of relevant technical language (names of compounds, 'substitution', 'elimination', 'delocalisation').</i></p> <p><b>Level 2 (3–4 marks)</b> Describes the differences between the two compounds in detail including electron delocalisation in phenol. Describes at least two reactions with equations but not necessarily showing reactions for both structures.</p> <p><i>The description is detailed and is presented with some structure. The reactions are in the most-part relevant and supported by equations. Demonstrates ability to answer question with some indications of a sound grasp of technical language.</i></p>	6	<p><b>Indicative scientific points may include:</b></p> <p><b>Ring structures:</b></p> <ul style="list-style-type: none"> <li>• saturated ring of 6 carbons in alcohol</li> <li>• unsaturated ring of 6 carbons in phenol with 6 delocalised electrons</li> </ul> <p><b>Reactions of –OH group:</b></p> <ul style="list-style-type: none"> <li>• acidic in phenol neutral in alcohol e.g. with alkalis* (NOT with carbonates)</li> <li>• nucleophilic substitution in alcohol e.g. with halide*</li> <li>• elimination in alcohol not in phenol e.g. form alkenes* with <math>Al_2O_3 / H_2SO_4</math></li> <li>• phenols give purple colour with <math>FeCl_3</math></li> <li>• phenols will not react with carboxylic acids but alcohols will*.</li> </ul> <p>* an equation can be written here</p> <p>If other correct reactions are given which are <b>NOT</b> different between the two compounds mark lower at each level</p>

Question	Answer	Marks	Guidance
	<p><b>Level 1 (1–2 marks)</b> Identifies the differences between the two structures mentioning phenol and alcohol. Describes at least two reactions.</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b> No response or no response worthy of credit.</p>		
(c)	<p>Gas–liquid chromatography <b>OR</b> Thin layer chromatography <b>OR</b> paper chromatography ✓</p>	1	<b>ALLOW</b> glc or tlc
(d)	 <p>Arrow from negative charge on cyanide ✓</p> <p>Partial charges on C=O correct <b>AND</b> arrow on carbonyl ✓</p> <p>Intermediate correct ✓</p> <p>Arrow from O to H <b>AND</b> correct cyanohydrin structure ✓</p>	4	<p>Arrows <b>MUST BE</b> double headed <b>AND</b> pointing towards correct atom</p> <p><b>ALLOW</b> lone pair inserted on C and arrow starting from there</p>
(e)	<p>(compound contains a) chiral C / chiral centre / asymmetric C ✓</p> <p>(the enantiomers are) mirror images ✓</p>	2	<b>ALLOW</b> 3-D sketch showing stereoisomers

H433/01


Mark Scheme

June 20XX

Question		Answer	Marks	Guidance
	(f)	Stronger intermolecular bonds present in raspberry ketone (than in 4-phenylbutan-2-one) ✓  due to hydrogen bonding ✓  so more energy needed to separate molecules ✓	3	<b>ALLOW</b> intermolecular forces
			<b>Total</b>	<b>17</b>

SPECIMEN

Question		Answer	Marks	Guidance
35	(a)	$Al^{3+}$ since group 3, $O^{2-}$ since Group 6; charges balance (in $Al_2O_3$ ) ✓	1	
	(b)	(i) $Al^{3+} + 3e^- \rightarrow Al$ ✓	1	<b>ALLOW</b> 'e' without minus <b>IGNORE</b> state symbols
		(ii) <b>FIRST CHECK ANSWER ON ANSWER LINE</b> Answer = 35000 <b>OR</b> 35280 award 3 marks  $n(Al_2O_3) = 100000/102$ <b>OR</b> 980 (mol) ✓  half a mol $CO_2$ for every mol O ✓ (stated or shown in calc)  volume $CO_2 = 24 \times 980 \times 3/2$ = 35000 <b>OR</b> 35280 ✓	3	<b>ALLOW ECF</b> from first marking point
	(c)	(i) cathode: $2H_2O + 2e^- \rightarrow H_2 + 2OH^-$ ✓  anode: $2H_2O \rightarrow 4H^+ + O_2 + 4e^-$ ✓	2	<b>ALLOW</b> 'e' without minus <b>IGNORE</b> state symbols <b>ALLOW</b> multiples and halves <b>ALLOW</b> $2H^+ + 2e^- \rightarrow H_2$ and $4OH^- \rightarrow O_2 + 2H_2O + 4e^-$
		(ii) white ppt ✓  after adding $Ba^{2+}$ / solution of (named) barium salt <b>OR</b> (formation of) barium sulfate/ $BaSO_4$ ✓	2	Any named barium salt must be soluble

Question	Answer	Marks	Guidance
(d) (i)	 <p>aluminium</p> <p>silver</p> <p>1.0 mol dm<sup>-3</sup> Al<sup>3+</sup></p> <p>1.0 mol dm<sup>-3</sup> Ag<sup>+</sup></p> <p>salt bridge</p> <p>labelled metal rods in labelled solutions of corresponding ions ✓  both ions 1.0 mol dm<sup>-3</sup> ✓  wires, voltmeter and salt bridge ✓</p>	3	<p><b>ALLOW:</b></p> <ul style="list-style-type: none"> <li>cell either way round</li> <li>any unambiguous representation of the voltmeter</li> <li>descriptions of solutions as, e.g., 'AgNO<sub>3</sub>' and 'Al(NO<sub>3</sub>)<sub>3</sub>' (but molarity must be correct, in next mark, for ,e.g., Al<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>).</li> </ul> <p><b>IGNORE</b></p> <p>temperature  any other label on salt bridge</p>
(ii)	<p><b>FIRST CHECK ANSWER ON ANSWER LINE</b>  Answer = - 1.68 V award 2 marks</p> $\frac{RT}{nF} \ln 0.1 = 8.314 \times 298 \times (-2.3) / 3 \times 9.65 \times 10^4$ $= -0.02 \text{ (-0.0197) (V) } \checkmark$ $E = -1.66 - 0.02 = -1.68 \text{ V } \checkmark$	2	
	<b>Total</b>	<b>14</b>	

## Summary of updates

---

Date	Version	Change
January 2019	2.0	Minor accessibility changes to the paper: i) Additional answer lines linked to Level of Response questions ii) One addition to the rubric clarifying the general rule that working should be shown for any calculation questions

SPECIMEN

**H433/01**

**Mark Scheme**

**June 20XX**

**BLANK PAGE**

SPECIMEN

**H433/01**

**Mark Scheme**

**June 20XX**

**BLANK PAGE**

SPECIMEN

**H433/01**

**Mark Scheme**

**June 20XX**

**BLANK PAGE**

SPECIMEN