

Please write clearly in	lock capitals.	
Centre number	Candidate number	
Surnama		
Surname		
Forename(s)		
Candidate signature		

# AS CHEMISTRY

Paper 1: Inorganic and Physical Chemistry

Friday 27 May 2016

Morning

Time allowed: 1 hour 30 minutes

#### **Materials**

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a calculator, which you are expected to use where appropriate.

## Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

## Information

- The maximum mark for this paper is 80.
- The Periodic Table/Data Sheet is provided as in insert.

#### Advice

You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.



7404/1

Section A			
	Answer <b>all</b> questions in this section.		
1	This question is about electron configuration.		
0 1 . 1	Give the full electron configuration of an Al atom and of a Cr <sup>3+</sup> ion.		
		[2 marks]	
	Al atom  Cr <sup>3+</sup> ion		
	CI 1011		
0 1 . 2	Deduce the formula of the ion that has a charge of 2+ with the same electron configuration as krypton.	tron	
		[1 mark]	
0 1 . 3	Deduce the formula of the compound that contains 2+ ions and 3- ions the have the same electron configuration as argon.	at both	
		[1 mark]	








2	This question is about Period 3 of the Periodic Table.	
0 2 . 1	Deduce which of Na <sup>+</sup> and Mg <sup>2+</sup> is the smaller ion. Explain your answer.	[2 marks]
	Smaller ion	-
	Explanation	
0 2 . 2	Write an equation to represent the process that occurs when the first io	nisation
	energy for sodium is measured.	[1 mark]
		[ · ··································



0 2 . 3 The first ionisation energies of some Period 3 elements are shown in Figure 1.





Complete **Figure 1** by plotting the approximate first ionisation energy values for magnesium and sulfur.

Explain why the first ionisation energy of sulfur is different from that of phosphorus.

[4 marks]

	_	



This question is about a white solid, MHCO<sub>3</sub>, that dissolves in water and reacts with hydrochloric acid to give a salt.

$$MHCO_3 + HCl \rightarrow MCl + H_2O + CO_2$$

A student was asked to design an experiment to determine a value for the  $M_r$  of MHCO<sub>3</sub>. The student dissolved 1464 mg of MHCO<sub>3</sub> in water and made the solution up to 250 cm<sup>3</sup>.

25.0 cm<sup>3</sup> samples of the solution were titrated with 0.102 mol dm<sup>-3</sup> hydrochloric acid. The results are shown in **Table 1**.

Table 1

	Rough	1	2	3
Initial burette reading / cm <sup>3</sup>	0.00	10.00	19.50	29.25
Final burette reading / cm <sup>3</sup>	10.00	19.50	29.25	38.90
Titre / cm <sup>3</sup>	10.00	9.50	9.75	9.65

0	3		1	Calculate the mean titre and use this to determine the amount, in moles,	of HCl
		_		that reacted with 25.0 cm <sup>3</sup> of the MHCO <sub>3</sub> solution.	
				·	[3 marks]
		_		_	
0	3	] .	2	, , , ,	
				Then calculate the experimental value for the $M_r$ of MHCO <sub>3</sub> .	
				Give your answer to the appropriate number of significant figures.	
					[3 marks]



0 3 . 3	The student identified use of the burette as the largest source of uncertainty in the experiment.
	Using the same apparatus, suggest how the procedure could be improved to reduce the percentage uncertainty in using the burette.
	Justify your suggested improvement.  [2 marks]
	Suggestion
	Justification
0 3 . 4	Another student is required to make up 250 cm <sup>3</sup> of an aqueous solution that contains a known mass of MHCO <sub>3</sub> . The student is provided with a sample bottle containing the MHCO <sub>3</sub> .
	Describe the method, including apparatus and practical details, that the student should use to prepare the solution.
	[6 marks]
	More answer space is available on page 8



'n	•	

-	
-	
•	
-	
-	
· ·	
-	
<del>-</del>	



**Table 2** shows some data about the elements bromine and magnesium.

Table 2

Element	Melting point / K	Boiling point / K	
Bromine	266	332	
Magnesium	923	1383	


0   4   .   1	different from that of magnesium. Suggest why magnesium is a liquid over a much greater temperature range compared to bromine.
	[5 marks]



5 Figure 2 represents two glass flasks, P and Q, connected via a tap.

Flask **Q** (volume =  $1.00 \times 10^3 \text{ cm}^3$ ) is filled with ammonia (NH<sub>3</sub>) at 102 kPa and 300 K. The tap is closed and there is a vacuum in flask **P**. (Gas constant R =  $8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

Figure 2



0   5	] . <u>[1</u>	Calculate the mass of ammonia in flask <b>Q</b> .  Give your answer to the appropriate number of significant figures.	[3 marks]



0 5 . 2	When the tap is opened, ammonia passes into flask <b>P</b> . The temperature decreases by 5 °C. The final pressure in both flasks is 75.0 kPa. Calculate the volume, in cm³, of flask <b>P</b> .	
		[3 marks]
	Turn over for the next question	



4	-	ı	
ı	٩	ì	

0	6		1	Explain how ions are accelerated, detected and have their abundance determined
		-		in a time of flight (TOF) mass spectrometer.

[3 marks]

0	6		Calculate the mass, in kg, of a single <sup>52</sup> Cr <sup>+</sup> ion.
			Assume that the mass of a 52Cr+ ion is the same as that of a 52Cr atom

(The Avogadro constant L =  $6.022 \times 10^{23} \text{ mol}^{-1}$ )

[1 mark]

 $\fbox{ \ \ \, 0\ \ \, 6\ \ \, }$  .  $\fbox{ \ \ \, 3\ \ \, }$  In a TOF mass spectrometer the kinetic energy (KE) of a  $^{52}Cr^+$  ion was 1.269 x  $10^{-13}\,J$ 

Calculate the velocity of the ion using the equation.

$$KE = \frac{1}{2}mv^2$$

 $(m = \text{mass/kg and } v = \text{velocity/ms}^{-1})$ 


[2 marks]



**0 6 . 4** Bromine has two isotopes, <sup>79</sup>Br and <sup>81</sup>Br, in approximately equal abundance. In a TOF mass spectrometer bromine forms ions with formula  $[Br_2]^+$ 

Sketch the pattern of peaks you would expect to see in the mass spectrum of a sample of bromine.

[2 marks]



 $oxed{0}$  .  $oxed{5}$  A sample of xenon has  $A_r$  = 131.31. The sample consists of four isotopes. The abundances of three of the isotopes are shown in **Table 3**. The data for one of the isotopes,  ${}^m\!Xe$ , is missing.

Table 3

Isotope	<sup>129</sup> Xe	<sup>131</sup> Xe	<sup>132</sup> Xe	<sup>m</sup> Xe
% abundance	28.0	25.0	27.0	To be calculated

Use the data to calculate the abundance of isotope <sup>m</sup>Xe and calculate m, the mass number of <sup>m</sup>Xe. Show your working.

[4 marks]



7 Ammonia reacts with aluminium chloride as shown by the	equation:
--	-----------

$$\mathsf{NH_3} \; + \; \mathsf{AlCl_3} \; \rightarrow \; \mathsf{H_3NAlCl_3}$$

[3 marks]



0 7 . 2	Name the type of bond formed between N and Al in $H_3NAlCl_3$ and exploit this bond is formed.	ain how [2 marks]
	Type of bond	
	Explanation	
0 7 . 3	Explain how the value of the Cl-Al-Cl bond angle in AlCl $_3$ changes, if a formation of the compound $H_3NAlCl_3$	t all, on [2 marks]
	Turn over for the next question	



8	A student oxidised a solution of hydrochloric acid with a few drops of sodi chlorate(I) solution. The reaction mixture effervesced and turned pale gree gas formed bleached universal indicator paper.	
0 8 . 1	Write a half-equation for the oxidation of chloride ions.	
		[1 mark]
0 8 . 2	Write a half-equation for the reduction of chlorate(I) ions to chlorine in aci conditions.	dic
		[1 mark]
0 8 . 3	Write an overall equation for the redox reaction of chlorate(I) ions with hydrochloric acid.	[1 mark]
0 8 . 4	A solution of sodium chlorate(I) was added to a colourless solution of pote iodide. Suggest what is observed.	assium
	Explain the reaction that leads to this observation.	[3 marks]



9	
ch the A	student was given a powder made from a mixture of anhydrous barium aloride and anhydrous magnesium chloride. The student dissolved 1.056 g of the powder in water in a conical flask and added an excess of sulfuric acid. White precipitate formed and was filtered off, washed and dried. The mass of this solid was 0.764 g.
	entify the white precipitate and calculate the percentage, by mass, of agnesium chloride in the powder.
	[4 marks]
<u> </u>	
_	
_	
_	
_	
_ _	

Turn over for the next question



## **Section B**


		3332	
	Answer a	II questions in the spaces prov	rided
	swer per question is all	lowed. he circle alongside the appropr	riate answer.
CORRECT MET	DD WRONG METHOD	os 🗴 💿 📾 🔯	
If you want	o change your answer	you must cross out your origin	al answer as shown.
If you wish t shown.	return to an answer pre	eviously crossed out, ring the an	swer you now wish to select as
	your working out in the badditional sheets for this	blank spaces around the questic working.	ons but this will not be marked.
			0
1   0	vvnich eiement is in th	ne d-block of the Periodic Table	
	A Selenium		[1 mark]
	<b>B</b> Antimony		<u> </u>
	<b>C</b> Tantalum		0
	<b>D</b> Lead		0
1 1	Which species contain	ns an element with an oxidation	n state of +4?
			[1 mark]
	A NO <sub>2</sub> <sup>+</sup>		0
	B ClO <sub>3</sub>		0
	C H <sub>2</sub> SO <sub>3</sub>		
	<b>D</b> PCl <sub>5</sub>		0



1 2	There are 392 mol of pure gold in a bar measuring 10 cm by 10 cm by 40 cm. What is the density of gold in kg dm <sup>-3</sup> ?			
	VVI	lat is the density of gold in kg diff	I	[1 mark]
	A	193	$\bigcirc$	
	В	19.3	$\bigcirc$	
	С	1.93	$\bigcirc$	
	D	0.193	$\circ$	
1   3	Ion	s of two isotopes of iron are		
		<sup>53</sup> Fe <sup>2+</sup>		
	Wh	ich statement is correct?		[1 mark]
		The ions of both the isotopes have the electronic configuration $1s^22s^22p^63s^23p^64s^23d^6$	0	[ i iliai kj
	В	The ions of both the isotopes contains 26 neutrons	$\bigcirc$	
	С	<sup>53</sup> Fe <sup>2+</sup> has fewer protons than <sup>56</sup> Fe <sup>2+</sup>	0	
	D	After acceleration to the same kinetic energy <sup>56</sup> Fe <sup>2+</sup> will move more slowly than <sup>53</sup> Fe <sup>2+</sup>	0	



1 4 The successive ionisation energies for element X are shown in Figure 3.



Which element is X?

[1 mark]

A Nitrogen

**B** Phosphorus

 $\circ$ 

**C** Aluminium

0

**D** Boron

0

1 5 Which of these decreases down Group 2?

[1 mark]

A First ionisation energy

**B** Atomic radius

0

**C** Number of protons

 $\circ$ 

**D** Reactivity with water

 $\circ$ 



Refer to the unbalanced equation below when answering questions 16 and 17.

$$\mathsf{K_2Cr_2O_7} \ + \ 3\mathsf{H_2C_2O_4} \ + \ \_\mathsf{H_2SO_4} \ \to \ \mathsf{Cr_2(SO_4)_3} \ + \ \_\mathsf{H_2O} \ + \ 6\mathsf{CO_2} \ + \ \mathsf{K_2SO_4}$$

1 6 In the balanced equation the mole ratio for sulfuric acid to water is

[1 mark]

**A** 1:4

 $\circ$ 

**B** 1:2

**C** 4:7

0

**D** 4:9

0

1 7 What is the reducing agent in this reaction?

[1 mark]

A H<sup>+</sup>

0

**B**  $C_2O_4^{2-}$ 

0

C K<sup>+</sup>

 $\circ$ 

**D**  $Cr_2O_7^{2-}$ 

0

റ	2
_	4

1 8	Which substance exists as a macromolecule?	
	<b>A</b> Cu	[1 mark]
	<b>B</b> SiO <sub>2</sub>	
	<b>C</b> P <sub>4</sub> O <sub>10</sub>	
	D MgO	
	g-	
1 9	A pale brown mixture of $NO_2$ and $N_2O_4$ is allowed to reach equilibri gas syringe according to the following equation.	um in a sealed
	$2NO_2(g) \rightleftharpoons N_2O_4(g)$	
	When the plunger is pushed further into the syringe the pressure in mixture becomes paler in colour.	ncreases and the
	When the syringe is placed in a hot oven the mixture becomes dar	ker in colour.
	Which of the following statements is correct?	[4 manula]
	<b>A</b> NO <sub>2</sub> is brown and the forward reaction is exothermic.	[1 mark]
	<b>B</b> NO <sub>2</sub> is brown and the forward reaction is endothermic.	0
	<b>C</b> NO <sub>2</sub> is colourless and the forward reaction is exothermic.	0
	<b>D</b> NO <sub>2</sub> is colourless and the forward reaction is endothermic.	



	23		
2 0	Which molecule has the largest dipole?		
	A ClF <sub>3</sub>	0	[1 mark]
	<b>B</b> BF <sub>3</sub>	$\bigcirc$	
	C SF <sub>6</sub>	$\bigcirc$	
	<b>D</b> CF <sub>4</sub>	$\bigcirc$	
2   1	In a molecule of a hydrocarbon, the fraction by mass of carbon is	<u>9</u> 11	
	What is the empirical formula of the hydrocarbon?		[4
	A CH	$\bigcirc$	[1 mark]
	B CH <sub>3</sub>	$\bigcirc$	
	<b>C</b> C <sub>3</sub> H <sub>8</sub>	$\bigcirc$	
	<b>D</b> C <sub>5</sub> H <sub>12</sub>	$\bigcirc$	



2 2
-----

30 cm³ of xenon are mixed with 20 cm³ of fluorine. The gases react according to the following equation. Assume that the temperature and pressure remain constant.

$$Xe(g) + F_2(g) \rightarrow XeF_2(g)$$

What is the final volume of gas after the reaction is complete?

[1 mark]

Δ	50	cm

0

0

 $\subset$ 

2	3

Which of the following solutions would react exactly with a solution containing 0.0500 mol sulfuric acid?

[1 mark]

Α	50.0	cm³	of	1.00	mol	dm <sup>-3</sup>	KOF
---	------	-----	----	------	-----	------------------	-----

**B** 100.0 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> KOH

0

C 100.0 cm<sup>3</sup> of 2.00 mol dm<sup>-3</sup> Ba(OH)<sub>2</sub>

0

**D** 50.0 cm<sup>3</sup> of 1.00 mol dm<sup>-3</sup> Ba(OH)<sub>2</sub>

0



2 4

In a car airbag, sodium azide (NaN $_{\!\scriptscriptstyle 3}$ ) decomposes to form sodium metal and nitrogen gas.

$$2NaN_3(s) \rightarrow 2Na(s) + 3N_2(g)$$

The sodium metal then reacts with potassium nitrate to produce more nitrogen gas.

$$10Na(s) + 2KNO_3(s) \rightarrow N_2(g) + 5Na_2O(s) + K_2O(s)$$

If 2.00 mol of sodium azide react in this way, how many molecules of  $N_2$  will be formed?

(The Avogadro constant L =  $6.022 \times 10^{23} \text{ mol}^{-1}$ )

[1 mark]

**A**  $2.41 \times 10^{24}$ 

 $\circ$ 

**B**  $1.93 \times 10^{24}$ 

0
---


**C**  $1.81 \times 10^{24}$ 

0
---


**D**  $9.63 \times 10^{23}$ 

0
---


**END OF QUESTIONS** 











#### **Copyright Information**

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright @ 2016 AQA and its licensors. All rights reserved.

