

Please write clearly in block	capitals.	
Centre number	Candidate number	
Surname		
Forename(s)		
Candidate signature		

AS CHEMISTRY

Paper 1: Inorganic and Physical Chemistry

Friday 26 May 2017

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
Section B		
TOTAL		

Section A			
Answer all questions in this section.			
0 1	This question is about atomic structure.		
0 1 . 1	Write the full electron configuration for each of the following species.	[2 marks]	
	Cl ⁻		
F	=e ²⁺		
0 1 . 2	Write an equation, including state symbols, to represent the process the when the third ionisation energy of manganese is measured.	at occurs [1 mark]	
0 1 . 3	State which of the elements magnesium and aluminium has the lower		
	first ionisation energy. Explain your answer.		
		[3 marks]	



0 1 . 4

A sample of nickel was analysed in a time of flight (TOF) mass spectrometer. The sample was ionised by electron impact ionisation. The spectrum produced showed three peaks with abundances as set out in **Table 1**.

Table 1

m/z	Abundance/%
58	61.0
60	29.1
61	9.9

Give the symbol, including mass number, of the ion that would reach the detector first in the sample.

Calculate the relative atomic mass of the nickel in the sample. Give your answer to one decimal place.

[3 marks]

|| |**| ||**

0	2	This questi

ion is about energetics.

0 2 1

Write an equation, including state symbols, for the reaction with an enthalpy change equal to the enthalpy of formation for iron(III) oxide.

[1 mark]

2

Table 2 contains some standard enthalpy of formation data.

Table 2

	CO(g)	Fe ₂ O ₃ (s)
Δ _f H ^e / kJ mol ⁻¹	-111	-822

$$Fe_2O_3(s) + 3CO(g) \rightarrow 2Fe(s) + 3CO_2(g)$$
 $\Delta H = -19 \text{ kJ mol}^{-1}$

$$\Delta H = -19 \text{ kJ mol}^{-1}$$

Use these data and the equation for the reaction of iron(III) oxide with carbon monoxide to calculate a value for the standard enthalpy of formation for carbon dioxide.

Show your working.

[3 marks]

$$\Delta_{\mathrm{f}}H^{\mathrm{e}}$$
 kJ mol^{-1}



0 2 . 3 Some enthalpy data are given in Table 3.

Table 3

Process	Δ <i>H</i> / kJ mol ⁻¹
$N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$	-92
$N_2(g) \rightarrow 2N(g)$	+944
$H_2(g) \rightarrow 2H(g)$	+436

Use the data from **Table 3** to calculate the bond enthalpy for N–H in ammonia. **[3 marks]**

N–H bond enthalpy	kJ mol ⁻¹

Question 2.3 Give one reason why the bond enthalpy that you calculated in Question **2.3** is different from the mean bond enthalpy quoted in a data book (388 kJ mol⁻¹).

[1 mark]



0	3

A student planned and carried out an experiment to determine the enthalpy of reaction when magnesium metal displaces zinc from aqueous zinc sulfate.

$$Mg(s) + Zn^{2+}(aq) \rightarrow Mg^{2+}(aq) + Zn(s)$$

The student used this method:

- A measuring cylinder was used to transfer 50 cm³ of a 1.00 mol dm⁻³ aqueous solution of zinc sulfate into a glass beaker.
- · A thermometer was placed in the beaker.
- 2.08 g of magnesium metal powder were added to the beaker.
- The mixture was stirred and the maximum temperature recorded.

The student recorded a starting temperature of 23.9 $^{\circ}\text{C}$ and a maximum temperature of 61.2 $^{\circ}\text{C}$.

0 3 . 1

Show by calculation which reactant was in excess.

Use the data to calculate the experimental value for enthalpy of reaction in kJ mol⁻¹(Assume that the specific heat capacity of the solution is 4.18 J K⁻¹g⁻¹ and the density of the solution is 1.00 g cm⁻³).

[6 marks]

Reactant in excess	

Enthalpy of reaction kJ mol⁻¹



0 3 . 2	Another student used the same method and obtained a value for the enthalpy of reaction of -142 kJ mol^{-1}	
	A data book value for the enthalpy of reaction is -310 kJ mol ⁻¹	
	Suggest the most likely reason for the large difference between the student's experimental value and the data book value.	
	[1 mark]	

Question 3 continues on the next page



0 3 . 3	Suggest how the students' method, and the analysis of the results, could be improved in order to determine a more accurate value for the enthalpy of reaction.		
	Justify your suggestions.		
	Do not refer to the precision of the measuring equipment. Do not change the amounts or the concentration of the chemicals.		
	[6 marks]		



0 4

When substances ${\bf P}$ and ${\bf Q}$ react together to form substance ${\bf R}$ an equilibrium is established according to the equation

$$P(g) + Q(g) \rightleftharpoons 2R(g)$$

The equilibrium constant expression is $K_c = \frac{[R]^2}{[P][Q]}$

1.0 mol of **P** and 1.0 mol of **Q** were mixed in a container with volume 1.0 dm³

At equilibrium, x mol of **P** had reacted.

0 4 . 1

The amount, in moles, of each of P and Q at equilibrium is (1-x).

Deduce in terms of x the amount, in moles, of \mathbf{R} in the equilibrium mixture.

[1 mark]

0 4 . 2

At 298 K the value of the equilibrium constant $K_c = 3.6$

Calculate a value for the equilibrium concentration, in mol dm^{-3} , of **R.**

[3 marks]

Equilibrium concentration of R

 $\mathrm{mol}\ \mathrm{dm}^{-3}$



0 5	This question is al	bout intermolecular forces.	
0 5 . 1	Give the meaning	of the term electronegativity.	[1 mark]
0 5 . 2		anent dipole-dipole forces arise	between hydrogen chloride
	molecules.		[2 marks]
0 5 . 3	Complete Table 4	by naming the shape of each n	nolecule.
	Place a tick (✓) in	the final column if the molecule	has a permanent dipole. [4 marks]
		Table 4	[+ mana]
	Molecule	Name of shape	Tick (✓) if molecule has a permanent dipole
	SiH ₄		
	PH ₃		
	BeCl ₂		



CH₃Cl

IB/M/Jun17/7404/1

0 6	Copper can be produced from rock that contains CuFeS ₂
0 6 . 1	Balance the equations for the two stages in this process. [2 marks]
Cı	$uFeS_2 +O_2 +SiO_2 \rightarrowCu_2S +Cu_2O +SO_2 +FeSiO_3$
0	$Cu_2S + \dots Cu_2O \rightarrow \dots Cu + \dots SO_2$
0 6 . 2	Suggest two reasons why the sulfur dioxide by-product of this process is removed from the exhaust gases. [2 marks]
	Reason 1
	Reason 2

Question 6 continues on the next page



0 6 . 3	A passenger jet contains 4050 kg of copper wiring.				
	A rock sample contains 1.25% CuFeS ₂ by mass.				
	Calculate the mass, in tonnes, of rock needed to produce enough copper wire				
	a passenger jet. (1 tonne = 1000 kg) [4	marks]			
	Mass of rock tonnes				



0	6	4

Copper can also be produced by the reaction of carbon with copper(II) oxide according to the equation

$$2\text{CuO} \ + \ \text{C} \rightarrow 2\text{Cu} \ + \ \text{CO}_2$$

Calculate the percentage atom economy for the production of copper by this process.

Give your answer to the appropriate number of significant figures.

[2 marks]

Percentage atom economy





An aqueous solution ${\bf Y}$ is known to contain one type of group 2 metal ion and one type of negative ion.

Aqueous solutions of sulfuric acid and magnesium nitrate are added to separate samples of solution $\bf Y$. The observations are shown in **Table 5**.

Table 5

Solution added	Observation with solution Y
Sulfuric acid	A white precipitate forms
Magnesium nitrate	A white precipitate forms

0 7.1	Suggest the identity of the group 2 metal ion present in solution Y.		
	Write an ionic equation, including state symbols, for the reaction that takes place when sulfuric acid is added to solution Y . [2 marks]		
	Group 2 metal ion		
	Ionic equation		
0 7.2	Suggest the identity of the negative ion present in solution Y.		
	Write an ionic equation, including state symbols, for the reaction that takes place when magnesium nitrate is added to solution Y .		
	[2 marks]		
	Negative ion		
	Ionic equation		

0 8	When an acidified solution of sodium nitrite (NaNO ₂) is added to aqueopotassium iodide, iodine and nitrogen monoxide (NO) are formed.	ous
0 8 . 1	Give the oxidation state of nitrogen in the following species.	[2 marks]
	NO ₂	
0 8 . 2	Write a half-equation for the conversion of NO_2^- in an acidic solution into NO	[1 mark]
0 8 . 3	Write a half-equation for the conversion of I^- into I_2	[1 mark]
0 8 . 4	Write an overall ionic equation for the reaction of NO_2^- in an acidic sol with I^-	ution [1 mark]
0 8 . 5	State the role of NO_2^- in the reaction with I^-	[1 mark]
	Question 8 continues on the next page	



0	8		6	
•	•	•	_	

In aqueous solution, nitrite ions react with acidified chlorate(V) ions according to the equation

$$2{\rm ClO_3}^- \ + \ 5{\rm NO_2}^- \ + \ 2{\rm H}^+ \rightarrow {\rm Cl_2} \ + \ 5{\rm NO_3}^- \ + \ {\rm H_2O}$$

A 25.0 cm 3 sample of an aqueous solution of sodium nitrite required 27.40 cm 3 of a 0.0200 mol dm $^{-3}$ solution of potassium chlorate(V) for complete reaction.

Calculate the concentration, in g dm⁻³, of sodium nitrite in the sample.

[4 marks]

Concentration of sodium nitrite ___ g dm⁻³



Se	ction	В
-	CUOII	_

Answer all questions in this section.

Only on	e answer	per	question	is	allowed.
----------------	----------	-----	----------	----	----------

For each answer completely fill in the circle alongside the appropriate answer.

CORRECT METHOD

WRONG METHODS

&	•		Φ

If you want to change your answer you must cross out your original answer as shown.



If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown.

You may do your working in the blank space around each question but this will not be marked. Do **not** use additional sheets for this working.

0 9

Which is the correct crystal structure for the substance named?

[1 mark]

	Substance	Structure
Α	lodine	Simple molecular
В	Diamond	Ionic
С	Sodium chloride	Giant covalent
D	Graphite	Metallic

\bigcirc

\circ

\bigcirc

1 0

Which is the best technique to remove the silver chloride that forms when aqueous solutions of silver nitrate and sodium chloride react?

[1 mark]

A Refluxing	ĺ
-------------	---



B Evaporation

 \bigcirc

C Filtration

0

D Distillation



1 1	Which statement about astatine is correct?	[1 mark]	
	A Astatine has a greater electronegativity than bromine		
	B Astatine is a better oxidising agent than bromine		
	C Astatine has a greater boiling point than bromine		
	D Astatine has a greater first ionisation energy than bromine	0	
1 2	Which statement about time of flight mass spectrometry is correct?	[1 mark]	
	A The current in the detector is proportional to the ion abundance	0	
	B Sample particles gain electrons to form positive ions		
	C Particles are detected in the order of their kinetic energies	0	
	D lons are accelerated by a magnetic field	0	
1 3	Chlorine exists as two isotopes ³⁵ Cl and ³⁷ Cl in the ratio 3:1		
	Which statement about peaks in the mass spectrum of Cl_2 is correct	ct? [1 mark]	
	A Peaks at $m/z = 70$ and 74 in the ratio 3:1	0	
	B Peaks at $m/z = 70$, 72 and 74 in the ratio 9:6:1	0	
	C Peaks at $m/z = 70$, 72 and 74 in the ratio 9:3:1	0	
	D Peaks at $m/z = 70$ and 72 in the ratio 3:1	0	



1 4	A 4.85 g sample of anhydrous sodium sulfate is dissolved in water and the solution made up to 250 cm ³ in a volumetric flask.		
	What is the concentration in mol dm ⁻³ of sodium sulfate in the solution?	[1 mark]	
	A 0.0341		
	B 0.137		
	C 0.163		
	D 0.273		
1 5	Which of these contains the greatest number of atoms?	[1 mark]	
	A 127 mg of iodine		
	B 1.54×10^{-4} kg of phosphorus		
	C 81.0 mg of carbon dioxide		
	D 1.70×10^{-4} kg of ammonia		
1 6	25.0 cm ³ samples of NaOH solution were taken by pipette from a beaker. The were then titrated with an aqueous solution of ethanoic acid. The concentrate ethanoic acid calculated from the experimental results was found to be lower than the actual value.		
	Which of these could explain the difference?	1 mark]	
	A Rinsing the pipette with distilled water before filling with NaOH		
	B Rinsing the burette with distilled water before filling with ethanoic acid		
	C Rinsing the walls of the conical flask with distilled water during the titration		
	D Rinsing the beaker with distilled water before filling with NaOH		



1 7	A 20.0 cm 3 sample of a 0.400 mol dm $^{-3}$ aqueous solution of a metal bromide (MBr $_n$) reacts exactly with 160 cm 3 of 0.100 mol dm $^{-3}$ aqueous silver nitrate.			
	What is the formula of the metal bromide?			
		[1 mark]		
	A MBr			
	B MBr ₂			
	C MBr ₃			
	D MBr ₄			
1 8	Which species has one or more bond angle(s) of 90°?	[1 mark]		
	A CH₄			
	B NH ₄ ⁺	0		
	C CIF ₄	0		
	D AlCl ₄			
1 9	The forward reaction in this equilibrium is endothermic			
	$COCl_2(g) \Rightarrow CO(g) + Cl_2(g)$			
	Which statement is correct?			
		[1 mark]		
	A If the total pressure is increased at constant temperature, the proportion of COCl ₂ in the equilibrium mixture will decrease			
	B Use of a catalyst will increase the proportion of COCl ₂ in the			
	equilibrium mixture at constant temperature and pressure			
	C Reducing the equilibrium concentration of CO will increase the value of the equilibrium constant			

Raising the temperature from 373 K to 473 K will increase the value of the equilibrium constant



D

 \circ

2	0	Which of these is not a redox reaction?
---	---	--

[1 mark]

$$\textbf{A} \quad \text{Cu}_2\text{O} \ + \ \text{H}_2\text{SO}_4 \rightarrow \text{CuSO}_4 \ + \ \text{Cu} \ + \ \text{H}_2\text{O}$$

B MgO + 2HCl
$$\rightarrow$$
 MgCl₂ + H₂O

$$\textbf{C} \quad \mathsf{SnCl}_2 \ + \ \mathsf{HgCl}_2 \rightarrow \mathsf{Hg} \ + \ \mathsf{SnCl}_4$$

$$\textbf{D} \quad \text{MnO}_2 \ + \ \text{4HCl} \rightarrow \text{MnCl}_2 \ + \ \text{2H}_2\text{O} \ + \ \text{Cl}_2$$

 \bigcirc

2 1 Which of these has the highest first ionisation energy?

[1 mark]

A Na

 \circ

B Al

C Si

D Cl

0

2 2 What is the empirical formula of an oxide of nitrogen that contains 26% nitrogen by mass?

[1 mark]

A NO_2

0

 $B N_2O_3$

 \bigcirc

 $\mathbf{C} \quad N_2O_5$

 \circ

 $D N_4O_5$

 \bigcirc

Which species is **not** produced by a redox reaction between solid sodium iodide and concentrated sulfuric acid?

[1 mark]

A Na₂SO₄

 \bigcirc

B H₂S


C S

0


D SO₂

0


END OF QUESTIONS











For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2017 AQA and its licensors. All rights reserved.

