



GCE

Mathematics B (MEI)

Unit **H630/01**: Pure Mathematics and Mechanics

Advanced Subsidiary GCE

Mark Scheme for June 2018

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2018

Annotations and abbreviations

Annotation in scoris	Meaning
✓ and ✕	
BOD	Benefit of doubt
FT	Follow through
ISW	Ignore subsequent working
M0, M1	Method mark awarded 0, 1
A0, A1	Accuracy mark awarded 0, 1
B0, B1	Independent mark awarded 0, 1
SC	Special case
^	Omission sign
MR	Misread
Highlighting	
Other abbreviations in mark scheme	Meaning
E1	Mark for explaining a result or establishing a given result
dep*	Mark dependent on a previous mark, indicated by *
cao	Correct answer only
oe	Or equivalent
rot	Rounded or truncated
soi	Seen or implied
www	Without wrong working
AG	Answer given
awrt	Anything which rounds to
BC	By Calculator
DR	This indicates that the instruction In this question you must show detailed reasoning appears in the question.

Subject-specific Marking Instructions for AS Level Mathematics B (MEI)

- a Annotations should be used whenever appropriate during your marking. The A, M and B annotations must be used on your standardisation scripts for responses that are not awarded either 0 or full marks. It is vital that you annotate standardisation scripts fully to show how the marks have been awarded. For subsequent marking you must make it clear how you have arrived at the mark you have awarded.
- b An element of professional judgement is required in the marking of any written paper. Remember that the mark scheme is designed to assist in marking incorrect solutions. Correct solutions leading to correct answers are awarded full marks but work must not be judged on the answer alone, and answers that are given in the question, especially, must be validly obtained; key steps in the working must always be looked at and anything unfamiliar must be investigated thoroughly. Correct but unfamiliar or unexpected methods are often signalled by a correct result following an apparently incorrect method. Such work must be carefully assessed. When a candidate adopts a method which does not correspond to the mark scheme, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner.
If you are in any doubt whatsoever you should contact your Team Leader.
- c The following types of marks are available.

M

A suitable method has been selected and *applied* in a manner which shows that the method is essentially understood. Method marks are not usually lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. In some cases the nature of the errors allowed for the award of an M mark may be specified.

A

Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated Method mark is earned (or implied). Therefore M0 A1 cannot ever be awarded.

B

Mark for a correct result or statement independent of Method marks.

E

A given result is to be established or a result has to be explained. This usually requires more working or explanation than the establishment of an unknown result.

Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored. Sometimes this is reinforced in the mark scheme by the abbreviation isw. However, this would not apply to a case where a candidate passes through the correct answer as part of a wrong argument.

- d When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. (The notation 'dep*' is used to indicate that a particular mark is dependent on an earlier, asterisked, mark in the scheme.) Of course, in practice it may happen that when a candidate has once gone wrong in a part of a question, the work from there on is worthless so that no more marks can sensibly be given. On the other hand, when two or more steps are successfully run together by the candidate, the earlier marks are implied and full credit must be given.
- e The abbreviation FT implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A and B marks are given for correct work only – differences in notation are of course permitted. A (accuracy) marks are not given for answers obtained from incorrect working. When A or B marks are awarded for work at an intermediate stage of a solution, there may be various alternatives that are equally acceptable. In such cases, what is acceptable will be detailed in the mark scheme. If this is not the case please, escalate the question to your Team Leader who will decide on a course of action with the Principal Examiner. Sometimes the answer to one part of a question is used in a later part of the same question. In this case, A marks will often be 'follow through'. In such cases you must ensure that you refer back to the answer of the previous part question even if this is not shown within the image zone. You may find it easier to mark follow through questions candidate-by-candidate rather than question-by-question.
- f Unless units are specifically requested, there is no penalty for wrong or missing units as long as the answer is numerically correct and expressed either in SI or in the units of the question. (e.g. lengths will be assumed to be in metres unless in a particular question all the lengths are in km, when this would be assumed to be the unspecified unit.) We are usually quite flexible about the accuracy to which the final answer is expressed; over-specification is usually only penalised where the scheme explicitly says so. When a value is given in the paper only accept an answer correct to at least as many significant figures as the given value. This rule should be applied to each case. When a value is not given in the paper accept any answer that agrees with the correct value to 2 s.f. Follow through should be used so that only one mark is lost for each distinct accuracy error, except for errors due to premature approximation which should be penalised only once in the examination. There is no penalty for using a wrong value for g. E marks will be lost except when results agree to the accuracy required in the question.
- g Rules for replaced work: if a candidate attempts a question more than once, and indicates which attempt he/she wishes to be marked, then examiners should do as the candidate requests; if there are two or more attempts at a question which have not been crossed out, examiners should mark what appears to be the last (complete) attempt and ignore the others. NB Follow these maths-specific instructions rather than those in the assessor handbook.
- h For a genuine misreading (of numbers or symbols) which is such that the object and the difficulty of the question remain unaltered, mark according to the scheme but following through from the candidate's data. A penalty is then applied; 1 mark is generally appropriate, though this may differ for some units. This is achieved by withholding one A mark in the question. Marks designated as cao may be awarded as long as there are no other errors. E marks are lost unless, by chance, the given results are established by equivalent working. 'Fresh starts' will not affect an earlier decision about a misread. Note that a miscopy of the candidate's own working is not a misread but an accuracy error.

- i If a graphical calculator is used, some answers may be obtained with little or no working visible. Allow full marks for correct answers (provided, of course, that there is nothing in the wording of the question specifying that analytical methods are required). Where an answer is wrong but there is some evidence of method, allow appropriate method marks. Wrong answers with no supporting method score zero. If in doubt, consult your Team Leader.



- j If in any case the scheme operates with considerable unfairness consult your Team Leader.

Question			Answer	Marks	AOs	Guidance
1			$\frac{8}{(3-\sqrt{5})} \times \frac{(3+\sqrt{5})}{(3+\sqrt{5})} = 6 + 2\sqrt{5}$	M1 A1 [2]	1.1a 1.1b	Attempt to rationalize the denominator Must be in correct notation
2			EITHER $(3)^3 + 3(3)^2(-2x) + 3(3)(-2x)^2 + (-2x)^3$ $= 27 - 54x + 36x^2 - 8x^3$	M1 M1 A1 A1 [4]	1.1a 1.1b 1.1b 1.1b	Use of Binomial coefficients Powers of 3 and (-2x) Condone no brackets or (2x) used. At least 3 simplified terms correct All correct and simplified
			OR $(3-2x)^2 = (9-12x+4x^2)$ $(3-2x)(9-12x+4x^2)$ $= 27 - 54x + 36x^2 - 8x^3$	M1 M1 A1 A1 [4]		Attempting to square Multiplying their answer by third bracket At least 3 simplified terms correct All correct and simplified


H630/01

Mark Scheme

June 2018

Question			Answer	Marks	AOs	Guidance
3			$\begin{pmatrix} 8 \\ 0 \end{pmatrix} + \begin{pmatrix} 2a \\ -3a \end{pmatrix} + \begin{pmatrix} 0 \\ b \end{pmatrix} = 0$ $a = -4,$ $b = -12$	M1 A1 A1 [3]	2.5 1.1b 1.1b	Setting up a correct vector equilibrium equation or two separate equations.
4			$T - \frac{1}{2}mg = 4ma$ $mg - T = ma$ $a = \frac{g}{10} \text{ m s}^{-2}, T = \frac{9mg}{10} \text{ N}$	B1 B1 M1dep A1 [4]	3.1b 3.1b 1.1a 1.1b	Award for first of these two equations Signs consistent with first equation Attempting to solve the simultaneous equations leading to either T or a so Both T and a needed simplified
5	i		$y = 4 \cos x$ $y = 2 \sin x$ 	B1 B1 [2]	1.2 1.2	Correct shapes and relative sizes Graphs labelled with correct amplitudes clear Ignore any graph beyond the given interval
	ii		$4 \cos x = 2 \sin x$ $\tan x = 2$ $x = \arctan 2$ $4 \cos x = 4 \times \frac{1}{\sqrt{5}} = \frac{4}{5} \sqrt{5}$  $\text{Point of intersection is } \left(\arctan 2, \frac{4}{5} \sqrt{5} \right)$	M1 A1 M1 A1 [4]	3.1a 1.1b 1.1a 1.1b	Forming equation leading to $\tan x = k$ oe Award for 63.4° for this first A mark Attempting to find exact value for $\cos x$ or $\sin x$ Must be a coordinate in this form
	iii		The student's argument is invalid as the period of $\tan x$ is 180° .	E1 [1]	2.3	Must state or imply that the argument is invalid and give a correct reason Allow for "roots occur every 180° " oe

Question			Answer	Marks	AOs		Guidance
6	i		$f(-1) = 4 \times (-1)^3 - 3(-1) + 1 = -4 + 3 + 1 = 0$ Therefore $(x + 1)$ is a factor	M1 A1 [2]	2.1 2.2a	DR Use of $f(-1)$ must be seen. Do not allow for algebraic division. Clear conclusion must be made	Allow without conclusion if preceded by “If $f(-1) = 0$ then $(x + 1)$ will be a factor” or similar
		ii	$f(x) = (x + 1)(4x^2 - 4x + 1) = 0$ $= (x + 1)(2x - 1)^2 = 0$ $x = -1, \frac{1}{2}$ [repeated]	M1 A1 A1 [3]	1.1a 1.1b 1.1b	DR Attempt to divide or to factorise by inspection with $4x^2$ correct quadratic factor seen or implied by correct linear factors Both roots seen derived from 3 correct linear factors or use of quadratic formula	Allow full credit for $(x + 1)(4x - 2)(x - 0.5)$ No marks for solving the cubic on the calculator
7			EITHER acceleration phase $v = 0 + 2.5 \times 2 = 5 \text{ m s}^{-1}$ slowing phase $v^2 = u^2 + 2as$ $0 = 5^2 + 2a \times 10$ $a = -1.25 \text{ m s}^{-2}$ $[-R] = 1.5 \times (-1.25) = -1.875$ Magnitude of $R = 1.875 \text{ N}$ (1.88 to 3sf)	M1 A1 M1 A1 M1 A1 [6]	3.1b 1.1b 3.1b 1.1b 1.1a 1.1b	Use of <i>suvat</i> equation(s) to find velocity. Do not allow if $s = 10$ used Use of <i>suvat</i> equation(s) with $s = 10$ to find acceleration FT their velocity. Must be correct sign. Use of Newton’s second law. FT their a $a \neq 2$ Must be positive	Must recognise two phases of motion for first 4 marks Consistent sign convention needed for full credit.


Question			Answer	Marks	AOs	Guidance
			<p>OR</p> <p>acceleration phase</p> $v = 0 + 2.5 \times 2 = 5 \text{ m s}^{-1}$ <p>v</p>  <p>5</p> <p>2.5</p> <p>T t</p> <p>using the distance to find the time it takes to stop using <i>areas</i> (second triangle):</p> $\frac{1}{2}(T - 2.5) \times 5 = 10$ <p>$T = 6.5$ so time to stop is 4 s.</p> <p>So $0 = 5 + 4a$</p> <p>Giving $a = -1.25 \text{ m s}^{-2}$</p> $[-R] = 1.5 \times (-1.25) = -1.875$ <p>Magnitude of $R = 1.875 \text{ N}$ (1.88 to 3sf)</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>[6]</p>	<p>Use of <i>suvat</i> equation(s) to find velocity. Do not allow if $s = 10$ used</p> <p>Use of area and <i>suvat</i> equation(s) to find acceleration</p> <p>Must be correct sign</p> <p>Use of Newton's second law.</p> <p>FT their $a \neq 2$ Must be positive</p>	<p>Must recognise two phases of motion for first 4 marks</p> <p>Consistent sign convention needed for full credit.</p>





Question			Answer	Marks	AOs	Guidance
8			EITHER			DR
			Equation of the form $y = k(x+1)(x-2)$	M1	1.1a	Allow with $k = 1$ and without $y =$
			$(0, -4)$ on curve so $k = 2$	M1	3.1a	Attempt to find $k \neq 1$
				A1	1.1b	All correct
			OR			
			Equation of the form $y = ax^2 + bx + c$			
			$(0, -4)$ on curve $c = -4$	(M1)		Uses one point to form an equation
			$(-1, 0)$ on the curve $0 = a - b - 4$	(M1)		Uses both other points and attempts to solve simultaneous equations
			$(2, 0)$ on the curve $0 = 4a - 2b - 4$			
			Solving simultaneous equations $a = 2, b = -2$	(A1)		All correct
			BOTH			
			Area = $\int_{-1}^2 (2x^2 - 2x - 4) \, dx$			
			$\left[\frac{2x^3}{3} - x^2 - 4x \right]_{-1}^2$	M1	1.1a	Integration – allow without limits – condone one error
				A1	1.1b	FT their quadratic
			$\left(\frac{2 \times 2^3}{3} - 2^2 - 4 \times 2 \right) - \left(\frac{2 \times (-1)^3}{3} - (-1)^2 - 4 \times (-1) \right)$	M1	1.1a	Substitution of limits clearly seen Complete argument leading to exact answer.
			$= -\frac{20}{3} - \frac{7}{3} = -9$	A1	2.1	Allow for 9 if there is an argument to explain the change of sign even if -9 not seen.
				E1	2.4	Must give modulus and explain the change of sign. FT if their definite integral is negative.
			Area is 9 below the x-axis.	[8]		
						“Area must be positive” is not sufficient explanation.

Question			Answer	Marks	AOs	Guidance
9	(i)		Using $y = f\left(\frac{x}{a}\right) \quad y = \left(\frac{x}{\frac{1}{2}} - 1\right)^2 = (2x - 1)^2$ $= 4x^2 - 4x + 1$	M1 A1 [2]	1.1a 2.1	Allow for 2 instead of $\frac{1}{2}$ used for method mark or attempt to write equation of quadratic that touches axis at (0.5, 0) AG Must be a convincing argument that references either stretch or $f(2x)$ or similar $(2x - 1)^2$ seen is sufficient for M1
	(ii)		EITHER C_2 is $y = 4.25x - x^2 - 3$ Normal to $y = 4x^2 - 4x + 1$ $\frac{dy}{dx} = 8x - 4$ At (0.1) $\frac{dy}{dx} = -4$ Gradient of normal is $\frac{1}{4}$ (0, 1) on line so equation of normal is $y = \frac{1}{4}x + 1$ Intersection of normal and C_2 $\frac{1}{4}x + 1 = 4.25x - x^2 - 3$ $4x^2 - 16x + 16 = 0$ EITHER $(x - 2)^2 = 0$ OR discriminant $16^2 - 4 \times 4 \times 16 = 0$ Repeated root so the normal is a tangent to C_2	B1 M1 M1 A1 M1 A1 E1 [7]	3.1a 1.1a 1.1b 1.1a 3.1a 1.1b 3.2a	Finding the equation of C_2 . Any form Finding the derivative Finding negative reciprocal of their gradient FT their value for derivative Attempt to solve simultaneous equations Repeated factor or root, or zero discriminant seen. Must interpret their solution in the context.

Question			Answer	Marks	AOs	Guidance
			OR C_2 is $y = 4.25x - x^2 - 3$	B1		Finding the equation of C_2 . Any form
			Normal to $y = 4x^2 - 4x + 1$ $\frac{dy}{dx} = 8x - 4$ At (0.1) $\frac{dy}{dx} = -4$	M1		Finding the derivative
			Gradient of normal is $\frac{1}{4}$	M1		Finding negative reciprocal of their gradient
			Equation of normal is $y = \frac{1}{4}x + 1$	A1		FT their value for derivative
			Point on C_1 where gradient is $\frac{1}{4}$ $\frac{dy}{dx} = 4.25 - 2x = \frac{1}{4}$ giving $x = 2$ $y = 1.5$	M1 A1		Attempting to find the point on C_1 where tangent parallel to the normal found. Both coordinates required
			EITHER So the equation of the tangent is $y - \frac{3}{2} = \frac{1}{4}(x - 2)$ Which is the same equation as the normal to C_1	E1		Correct equation for the tangent in form that makes it clear it is the same line as the normal.
			OR show that point (2, 1.5) lies on normal So the normal to C_1 is a tangent to C_2	(E1) [7]		

14

Question			Answer	Marks	AOs	Guidance
10	i			B1	1.1a	Two line segments with one horizontal
				B1	1.1a	($T, 4$) and ($12.5, 4$) labelled or indicated on scales. Allow their 2.5 marked instead of T . On axes labelled v and t oe
				[2]		
	ii		$\frac{1}{2} \times 4 \times (12.5 + (12.5 - T)) = 45$ $T = 2.5$	M1 A1 [2]	3.1a 1.1b	Attempt to find area of trapezium or both the triangle $\left(\frac{1}{2}T \times 4\right)$ and the rectangle $(12.5 - T) \times 4$. cao
	iii		EITHER $a = \frac{4}{2.5} = 1.6 \text{ m s}^{-2}$ $s = \frac{1}{2} \times 1.6t^2 = 0.8t^2$	M1 A1 [2]	1.1a 3.3	Soi FT their T
			OR $a = \frac{4}{2.5} = 1.6 \text{ m s}^{-2}$ $v = \int a \, dt = 1.6t + c$ When $t = 0, v = 0$ so $c = 0$ $s = \int v \, dt = 0.8t^2 + c$ When $t = 0, s = 0$ so $c = 0$ Giving $s = 0.8t^2$	M1 A1 [2]		Soi FT their T Must be complete solution – do not award without consideration of $+c$ at least once
	iv		$0.8t^2 = 4$ $t = \sqrt{5} = 2.24 \text{ s}$	B1FT [1]	3.4	FT their quadratic model in (iii)

Question			Answer	Marks	AOs		Guidance
	v			B1 [1]	1.1a	Must have curved section of the graph decreasing gradient. <i>S</i> must be labelled.	
	vi		<p>Total distance (area under the graph) can only be equal if $S > T$ If $S > T$</p>  <p>If $S = T$</p>  <p>If $S < T$</p> 	E1 [1]	3.5c	Needs to give reason relating to the refinement of the model. Graphs not required	“It takes longer to reach 4 ms^{-1} ” is not sufficient reason

Question			Answer	Marks	AOs	Guidance									
11	i		$y = kx^n$	B1 [1]	3.3	Allow any letters used for constant of proportionality and power.									
	ii		$\ln y = \ln(kx^n)$ $\ln y = \ln k + \ln(x^n) = \ln k + n \ln x$	M1 E1 [2]	2.1 2.1	Taking natural logs of both sides and one correct use of laws of logs used. Convincing argument. AG									
	iii		<table><tr><td></td><td>$\ln x$</td><td>$\ln y$</td></tr><tr><td>Mercury</td><td>-1.179</td><td>9.575</td></tr><tr><td>Jupiter</td><td>1.599</td><td>4.022</td></tr></table>		$\ln x$	$\ln y$	Mercury	-1.179	9.575	Jupiter	1.599	4.022	B1 B1 [2]	1.1b 1.1b	At least 2 correct values All correct and to 4sf
	$\ln x$	$\ln y$													
Mercury	-1.179	9.575													
Jupiter	1.599	4.022													
	iv		EITHER $b = \frac{9.575 - 4.022}{-1.179 - 1.599} = -1.999$ (-2.00 to 3sf) $a = 7.218$	M1 A1 A1 [3]	1.1a 3.1a 1.1b	using gradient formula Allow -2 a correct to at least 2 sf									
			OR $9.575 = a - 1.179b$ $4.022 = a + 1.599b$ Giving $a = 7.218$ $b = -1.999$ (-2.00 to 3sf)	M1 A1 A1 [3]		Setting up pair of equations by substitution of their values. Allow one slip. a correct to at least 2 sf b correct to at least 2 sf									
	v		$y = 1363x^{-2.00}$	B1 B1 [2]	2.2a 3.3	FT their equation in (i) awrt 1300 or 1400, or $e^{7.2}$ or better. FT their a Allow for x^{-2} or better. FT their b									
	vi			B1 B1 [2]	1.2 1.1b	Appropriate curve with at least one horizontal asymptote or vertical asymptote shown Both asymptotes correct Ignore $x < 0$ if shown									
	vii		Earth $x = 1$, $y = 1363 \times 1^{-2} = 1360 \text{ W m}^{-2}$ (3sf)	B1 [1]	3.4	FT their (v)									

Question			Answer	Marks	AOs		Guidance	
			SCHEME FOR CANDIDATES USING LOG BASE 10					
	ii			SC1		Correct use of log instead of ln and no other error		
	iii			B0 B1		All correct. Must be 4 sf		
		Mercury	-0.5122					4.158
		Jupiter	0.6946					1.747
	iv		As in main scheme $a = 3.1336$ $b = -1.999$	M1 A1 A1				
	v		$y = 1360x^{-2}$	B1 B1		FT their equation in (i) awrt 1300 or 1400, or $e^{3.1}$ (= 22.2), $10^{3.1}$ or better. FT their a allow for x^{-2} or better. FT their b		
	vii		Earth $x = 1$, $y = 1363 \times 1^{-2} = 1360 \text{ W m}^{-2}$ (3sf)	B1		FT their (v)		

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA
Registered Company Number: 3484466
OCR is an exempt Charity

OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2018

