

Please write clearly in	block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature			

GCSE CHEMISTRY

F

Foundation Tier F

Paper 1

Thursday 17 May 2018

Morning

Time allowed: 1 hour 45 minutes

Materials

For this paper you must have:

- a ruler
- a scientific calculator
- the periodic table (enclosed).

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions in the spaces provided.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- In all calculations, show clearly how you work out your answer.

Information

- There are 100 marks available on this paper.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- You are reminded of the need for good English and clear presentation in your answers.

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
TOTAL		
IOIAL		



0 1 This question is about mixtures. Substances are separated from a mixture using different methods. 0 1 Draw **one** line from each substance and mixture to the best method of separation. [3 marks] **Substance and mixture** Method of separation Chromatography Ethanol from ethanol and water Crystallisation Electrolysis Salt from sea water Filtration The different colours in black ink Fractional distillation



Do not write outside the

box

2 A student filters a mixture. 0 1 . Figure 1 shows the apparatus. Figure 1 Beaker. Filter paper cone Flask Suggest one improvement to the apparatus. [1 mark] 0 1 . 3 Complete the sentences. Choose answers from the box. [2 marks] condense evaporate freeze melt solidify In simple distillation, the mixture is heated to make the liquid The vapour is then cooled to make it



Do not write Figure 2 shows the arrangement of atoms in a pure metal and in a mixture of metals. Figure 2 Pure metal Mixture of metals Metal A Metal A Calculate the percentage of metal **B** atoms in the mixture of metals shown in Figure 2. [2 marks] Percentage of metal **B** atoms = What is a mixture of metals called? 1 5 [1 mark] Tick one box. An alloy A compound A molecule A polymer



outside the

box

0 1.6	Why is the mixture of metals in Figure 2 harder than the pure metal? [1 mar Tick one box.	Do not write outside the box
	The atoms in the mixture are different shapes.	
	The layers in the mixture are distorted.	
	The layers in the mixture slide more easily.	
	The mixture has a giant structure.	
0 1.7	A nanoparticle of pure metal A is a cube.	
	Each side of the cube has a length of 20 nm.	
	Figure 3 shows the cube.	
	Figure 3	
	20 nm	
	What is the volume of the nanoparticle?	
	Tick one box.	KJ
	20 nm ³	
	60 nm ³	
	400 nm ³	
	8000 nm ³	11



0 2	The halogens are elements in Group 7.	Do not write outside the box
0 2 . 1	Bromine is in Group 7.	
	Give the number of electrons in the outer shell of a bromine atom. [1 mark]	
0 2.2	Bromine reacts with hydrogen. The gas hydrogen bromide is produced.	
	What is the structure of hydrogen bromide? [1 mark]	
	Tick one box.	
	Giant covalent	
	Ionic lattice	
	Metallic structure	
	Small molecule	
0 2.3	What is the formula for fluorine gas? [1 mark] Tick one box.	
	F	
	F_2	
	F^2	
	2F	



A student mixes solutions of halogens with solutions of their salts.

Table 1 shows the student's observations.

Table 1

	Potassium chloride (colourless)	Potassium bromide (colourless)	Potassium iodide (colourless)
Chlorine (colourless)		Solution turns orange	Solution turns brown
Bromine (orange)	No change		Solution turns brown
lodine (brown)	No change	No change	

0 2 . 4	Explain how the reactivity of the halogens changes going down Group 7.		
	Use the results in Table 1 .	[3 marks]	

Question 2 continues on the next page



	A company uses chlorine to produce titanium chloride from titanium dioxide.	Do not write outside the box
0 2.5	What is the relative formula mass (M_r) of titanium dioxide, TiO_2 ?	
	Relative atomic masses (A_r): $O = 16$ $Ti = 48$ [1 mark]	
	Tick one box.	
	64	
	80	
	128	
	768	
0 2.6	The company calculates that 500 g of titanium dioxide should produce 1.2 kg of titanium chloride.	
	However, the company finds that 500 g of titanium dioxide only produces 900 g of titanium chloride.	
	Calculate the percentage yield.	
	[2 marks]	
	Percentage yield =%	
		9



Turn over for the next question	Do not write outside the
rum over for the next question	box
DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED	
ANSWER IN THE SPACES PROVIDED	





0 3

0 3 . 1

Do not wi	ite
outside ti	hε
hox	

This question is about the structure of the atom.					
Comple	ete the sentences.				
Choose	e answers from the box	x .			
Each w	ord may be used once	, more than onc	e, or not at all.	[5 marks]	
	electron	ion	neutron		
	nucleus	s prot	on		
The ce	ntre of the atom is the			<u> </u> .	
The two	o types of particle in th	e centre of the a	tom are the proton	ı	
and the					
James	Chadwick proved the	existence of the		<u>.</u>	
Niels Bohr suggested particles orbit the centre of the atom. This type of particle					
is the		·			
The two	o types of particle with	the same mass	are the neutron		
and the)				

Table 2 shows information about two isotopes of element **X**.

Table 2

	Mass number	Percentage (%) abundance
Isotope 1	63	70
Isotope 2	65	30



0 3.2	Calculate the relative atomic mass (A_r) of element X using the equation:	Do not write outside the box
$A_{r}=$	(mass number × percentage) of isotope 1 + (mass number × percentage) of isotope 2	
	Use Table 2 .	
	Give your answer to 1 decimal place. [2 marks]	
	$A_{r} = \underline{\hspace{2cm}}$	
0 3.3	Suggest the identity of element X.	
	Use the periodic table. [1 mark]	
	Element X is	
0 3.4	The radius of an atom of element \mathbf{X} is $1.2 \times 10^{-10} \text{m}$	
	The radius of the centre of the atom is $\frac{1}{10000}$ the radius of the atom.	
	Calculate the radius of the centre of an atom of element X .	
	Give your answer in standard form. [2 marks]	
	Radius = m	

10




0 4

A student investigated the electrolysis of sodium chloride solution.

Figure 4 shows the apparatus.

Figure 4




The student measured the volume of gas collected in each measuring cylinder every minute for 20 minutes.



0 4 . 1

Figure 5 shows the volume of hydrogen gas collected in the measuring cylinder after 8 minutes.

Figure 5



What is the volume of hydrogen gas collected?


[1 mark]


Volume = ____ cm³

Question 4 continues on the next page



Do not write outside the box





0 4.2	Which of the lines on Figure 6 show that the volume of gas collected is directly proportional to the time?		′
	Tick one box.		1 mark]
	Both lines		
	Chlorine line only		
	Hydrogen line only		
	Neither line		




0 4.3	Which of the lines on Figure 6 show a positive correlation between the volume of gas collected and time? [1 mark] Tick one box. Both lines Chlorine line only Hydrogen line only Neither line	Do not write outside the box
	Question 4 continues on the next page	



A teacher demonstrates the electrolysis of different substances using graphite electrodes.

Figure 7 shows the apparatus used.

Figure 7



0 4.4	Why can graphite conduct electricity? Tick one box.	[1 mark]
	Graphite exists in layers of atoms.	
	Graphite has a giant structure.	
	Graphite has a high melting point.	
	Graphite has delocalised electrons.	



0 4 . 5

The teacher demonstrates the electrolysis of:

- molten zinc chloride
- potassium bromide solution.

Complete **Table 3** to predict the products.

Choose answers from the box.

[4 marks]

Do not write outside the box

chlorine	bromine	hydrogen	oxygen	potassium	zinc	
İ						

Table 3

Substance electrolysed	Product at cathode (negative electrode)	Product at anode (positive electrode)
Molten zinc chloride		
Potassium bromide solution		

Turn over for the next question

8



0 5

A student investigated the mass of copper oxide produced by heating copper carbonate.

This is the method used.

- 1. Weigh an empty test tube.
- 2. Weigh 2.00 g of copper carbonate into the test tube.
- 3. Heat the copper carbonate until there appears to be no further change.
- 4. Re-weigh the test tube and copper oxide produced.
- 5. Subtract the mass of the empty tube to find the mass of copper oxide.
- 6. Repeat steps 1–5 twice.
- 7. Repeat steps 1–6 with different masses of copper carbonate.

Table 4 shows the student's results.

Table 4

Mass of copper	Mass of copper oxide in g			
carbonate in g	Trial 1	Trial 2	Trial 3	Mean
2.00	1.29	1.27	1.31	1.29
4.00	2.89	2.57	2.59	2.58
6.00	3.85	3.90	3.87	3.87
8.00	5.12	5.15	5.09	Х
10.00	6.42	6.45	6.45	6.44

The equation for the reaction is:

$$CuCO_3(s) \rightarrow CuO(s) + CO_2(g)$$

0 5 . 1 Complete the sentence.

[1 mark]

The state symbol shows carbon dioxide is a



0 5.2	Why do the contents of the test tube lose mass in the investigation? [1 mark]	Do not writ outside the box
0 5.3	Calculate the mean mass X in Table 4. [1 mark]	
	X = g	
0 5.4	One of the results in Table 4 is anomalous.	
	Which result is anomalous? [1 mark]	
	Mass of copper carbonateg Trial	
0 5.5	Suggest how the investigation could be improved to make sure the reaction is complete. [2 marks]	




Another student repeated the investigation using magnesium carbonate instead of copper carbonate.

The word equation for the reaction is:

magnesium carbonate → magnesium oxide + carbon dioxide

Figure 8 shows the results of the investigation.

Figure 8





Draw a line of best fit on Figure 8 . [1 mark]	Do not write outside the box
Determine the mass of magnesium oxide produced by 8.4 g of magnesium carbonate. Use Figure 8 .	
[1 mark]	
Mass = g	
Calculate the mass of magnesium oxide produced when 168 g of magnesium carbonate is heated. Use your answer to Question 05.7 [2 marks]	
Mass of magnesium oxide produced = g	
Turn over for the next question	10
	Determine the mass of magnesium oxide produced by 8.4 g of magnesium carbonate. Use Figure 8. [1 mark] Mass = g Calculate the mass of magnesium oxide produced when 168 g of magnesium carbonate is heated. Use your answer to Question 05.7 [2 marks] Mass of magnesium oxide produced = g



0 6

A student investigated the temperature change in displacement reactions between metals and copper sulfate solution.

This is the method used.

- 1. Measure 50 cm³ of the copper sulfate solution into a polystyrene cup.
- 2. Record the starting temperature of the copper sulfate solution.
- 3. Add the metal and stir the solution.
- 4. Record the highest temperature the mixture reaches.
- 5. Calculate the temperature increase for the reaction.
- 6. Repeat steps 1-5 with different metals.

0 6 . 1

Draw **one** line from each type of variable to the name of the variable in the investigation.

[2 marks]

Name of variable in the investigation

Concentration of solution

Dependent variable

Particle size of solid

Temperature change

Independent variable

Type of metal

Volume of solution



		I
0 6.2	The student used a polystyrene cup and not a glass beaker.	Do not write outside the box
	Why did this make the investigation more accurate? [1 mark]	
	Tick one box.	
	Glass is breakable	
	Glass is transparent	
	Polystyrene is a better insulator	
	Polystyrene is less dense	
	Question 6 continues on the next page	

Do not write outside the box

Table 5 shows the student's results.

Table 5


Metal	Temperature increase in °C
Magnesium	38
Nickel	8
Zinc	16

0 6.3 Complete Figure 9.

Use data from Table 5.

[2 marks]

Figure 9






0 6.4	The student concluded that the reactions between the metals and copper sulfate solution are endothermic.
	Give one reason why this conclusion is not correct. [1 mark]
0 6.5	The temperature increase depends on the reactivity of the metal.
	Write the metals magnesium, nickel and zinc in order of reactivity.
	Use Table 5. [1 mark]
	Most reactive
	Least reactive
0 6.6	Y is an unknown metal.
	Describe a method to find the position of Y in the reactivity series in Question 06.5 [3 marks]





Figure 10 shows the reaction profile for the reaction between zinc and copper sulfate solution.

Figure 10



0 6 . 7 Which letter represents the products of the reaction?

[1 mark]

Tick **one** box.

A B C D E

0 6 . 8 Which letter represents the activation energy?

[1 mark]

Tick **one** box.

A B C D E

12



Do not write outside the box Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 7	This question is about elements in Group 1.	Do not write outside the box
	A teacher burns sodium in oxygen.	
0 7.1	Complete the word equation for the reaction. [1 mark]	
	sodium + oxygen →	
0 7.2	What is the name of this type of reaction? [1 mark] Tick one box.	
	Decomposition	
	Electrolysis	
	Oxidation	
	Precipitation	
0 7.3	The teacher dissolves the product of the reaction in water and adds universal indicator.	
	The universal indicator turns purple.	
	What is the pH value of the solution?	
	Tick one box.	
	1 7 13	



0 7.4	The solution contains a substance with the formula NaOH		
	Give the name of the substance. [1 mark]		
	[1 mark]		
0 7.5	All alkalis contain the same ion.		
	What is the formula of this ion? [1 mark]		
	Tick one box.		
	H ⁺		
	Na⁺		
	OH ⁻		
	O ²⁻		
0 7.6	A solution of NaOH had a concentration of 40 g/dm ³		
	What mass of NaOH would there be in 250 cm³ of the solution? [2 marks]		
	Mass = g		



0 7.7

The melting points of the elements in Group 1 show a trend.

Table 6 shows the atomic numbers and melting points of the Group 1 elements.


Table 6

Element	Atomic number	Melting point in °C
Lithium	3	181
Sodium	11	98
Potassium	19	63
Rubidium	37	Х
Caesium	55	29

Plot the data from Table 6 on Figure 11.

[2 marks]








0 8	Soluble salts are formed by reacting metal oxides with acids.
0 8.1	Give one other type of substance that can react with an acid to form a soluble salt. [1 mark]
0 8.2	Calcium nitrate contains the ions Ca ²⁺ and NO ₃ ⁻
	Give the formula of calcium nitrate. [1 mark]
0 8.3	Describe a method to make pure, dry crystals of magnesium sulfate from a metal oxide and a dilute acid.
	[6 marks]



	Do not write outside the box
Turn over for the next question	8





0 9	This question is about metals and metal compounds.	
0 9 . 1	Iron pyrites is an ionic compound.	
	Figure 12 shows a structure for iron pyrites.	
	Figure 12	
	Key Fe S	
	Determine the formula of iron pyrites.	
	Use Figure 12.	[1 mark]
0 9.2	An atom of iron is represented as $^{56}_{26}$ Fe Give the number of protons, neutrons and electrons in this atom of iron. Number of protons Number of neutrons Number of electrons	[3 marks]
0 9 . 3	Iron is a transition metal.	
	Sodium is a Group 1 metal.	
	Give two differences between the properties of iron and sodium.	[2 marks]
	1	
	2	



	Nickel is extracted from nickel oxide by reduction with carbon.	
0 9.4	Explain why carbon can be used to extract nickel from nickel oxide.	[2 marks]
0 9.5	An equation for the reaction is:	
	$NiO + C \rightarrow Ni + CO$	
	Calculate the percentage atom economy for the reaction to produce nickel.	
	Relative atomic masses (A_r): $C = 12$ $Ni = 59$	
	Relative formula mass (M_r) : NiO = 75	
	Give your answer to 3 significant figures.	[3 marks]
		[o marko]
	Percentage atom economy =	%

11



1 0	Chemical reactions can produce electricity.			
1 0.1	Figure 13 shows a simple cell.			
		Figu	ure 13	
		Electrode A	Electrode B	
	Which of these combinations would not give a zero reading on the voltmeter in Figure 13 ?			oltmeter in [1 mark]
	Tick one box.	Tick one box.		
	Electrode A	Electrode B	Electrolyte	
	Copper	Copper	Sodium chloride solution	
	Zinc	Zinc	Water	
	Copper	Zinc	Sodium chloride solution	
	Copper	Zinc	Water	
	Figure 13? Tick one box. Electrode A Copper Zinc Copper	Electrode B Copper Zinc Zinc	Electrolyte Sodium chloride solution Water Sodium chloride solution	



	Alkaline batteries are non-rechargeable.	Do not write outside the box
1 0 . 2	Why do alkaline batteries eventually stop working? [1 mark]	
1 0.3	Why can alkaline batteries not be recharged? [1 mark]	
	Question 10 continues on the next page	



1 0 . 4

1 0.5

	38			
Hydrogen fuel cells and rechargeable lithium-ion batteries can be used to power electric cars.				
Complete the balanced equation f	or the overall reaction ir	n a hydrogen fuel cell. [2 marks]		
H ₂ +	<i>→</i>	H ₂ O		
Table 7 shows data about differen	nt ways to power electric	c cars.		
	Table 7			
	Hydrogen fuel cell	Rechargeable lithium-ion battery		
Time taken to refuel or recharge in minutes	5	30		
Distance travelled before refuelling or recharging in miles	Up to 415	Up to 240		
Distance travelled per unit of energy in km	22	66		
Cost of refuelling or recharging in £	50	3		
Minimum cost of car in £	60 000	18 000		
Evaluate the use of hydrogen fuel cells compared with rechargeable lithium-ion batteries to power electric cars. Use Table 7 and your own knowledge. [6 marks]				



30	
	Do not write outside the
	box
	11
END OF QUESTIONS	



There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED Copyright information For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

