

Please write clearly in	ı block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature	_		— <i>)</i>

AS BIOLOGY

Paper 2

Monday 4 June 2018

Afternoon

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.


Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the space provided. Do not write outside the box around each page or on blank pages.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 75.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
TOTAL	li de la companya de





		3		
		Answer all questions in the spaces provided.		
1.1	Struct	cures A to E are parts of a plant cell.		
	B Ch C Nu D Mi	ell Wall nloroplast ucleus itochondrion olgi apparatus		
	Comp	plete Table 1 by putting the correct letter, A , B , C , D or E in the ment.	box next to e	each
		, -, -,	[3 m	arks]
		Table 1		
		Statement	Letter	
		Has stacked membranes arranged in parallel and contains DNA.		
		Is made of polysaccharide.		
		Is an organelle and is not surrounded by two membranes.		
1.2	adapt conta	in breast milk is produced and secreted by gland cells. These gations that include many mitochondria and many Golgi vesicles ins a high concentration of protein. in the role of these cell adaptations in the production and secret milk.	. The milk	ave
	Diodo	· · · · · · · · · · · · · · · · · · ·	[2 m	arks



Do not	writ
outside	th
box	(

0 2.1	Describe how a peptide bond is formed between two amino acids to form a	dipeptide. [2 marks]
0 2 . 2	The secondary structure of a polypeptide is produced by bonds between an	nino acids.
	Describe how.	
		[2 marks]



0 2 . 3	Two proteins have the same number and type of amino acids but different tertiary structures.	Do not wr outside th box
	Explain why. [2 marks]	
		6

Turn over for the next question



0 3 .

1

6	
Describe the relationship between size and surface area to volu	ume ratio of organisms. [1 mark]
A scientist calculated the surface area of a large number of frog he mean surface area was 9.73 mm². Frog eggs are spherica	g eggs. He found that I.
The surface area of a sphere is calculated using this equation	
Surface area = $4\pi r^2$	
where r is the radius of a sphere	
$\pi = 3.14$	
Se this equation to calculate the mean diameter of a frog egg.	
Show your working.	[2 marks]
Diameter =	mm

Use this equation to calculate the mean diameter of a frog egg.
Show your working. [2 marks]
Diameter = mm



The scientist calculated the ratio of surface area to mass for eggs, tadpoles and frogs. He also determined the mean rate of oxygen uptake by tadpoles and frogs.

His results are shown in Table 2.

Table 2

Stage of frog development	Ratio of surface area to mass	Mean rate of oxygen uptake / μmol g ⁻¹ h ⁻¹
Egg	2904 : 1	no information
Tadpole	336 : 1	5.7
Adult	166 : 1	1.3

0 3.3	The scientist used units of μ mol $g^{-1} h^{-1}$ for the rate of oxygen uptake.
	Suggest why he used µmol in these units. [1 mark]
0 3.4	The scientist decided to use the ratio of surface area to mass, rather than the ratio of surface area to volume. He made this decision for practical reasons.
	Suggest one practical advantage of measuring the masses of frog eggs, tadpoles and adults, compared with measuring their volumes. [1 mark]



0 3.5	Explain why oxygen uptake is a measure of metabolic rate in organisms.	rk]
0 3 . 6	A student who looked at these results said that they could not make a conclusion about the relationship between stage of development and metabolic rate.	
	Use information in Table 2 to explain reasons why they were unable to make a	
	conclusion. [3 mark	ks]



Do not writ
outside the
box


0 4.1	Give two similarities in the movement of substances by diffusion and by osm	osis.
		[2 marks]
	1	
	2	
	Question 4 continues on the next page	



A scientist measured the rate of uptake of a monoglyceride and a monosaccharide by epithelial cells of the small intestine of mice. A monoglyceride is a molecule of glycerol with one fatty acid attached. She did this for different concentrations of monoglyceride and monosaccharide.

Her results are shown in Figure 1.

Figure 1





4 . 2	Use your knowledge of transport across membranes to explain the shape of the in Figure 1 for uptake of monosaccharides between concentrations:	e curve
	A and B	
	C and D	
4.3	Figure 1 is evidence for monoglycerides being lipid-soluble molecules.	
	Suggest how. [2	marks]



There are no questions on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



0 5.1	A student prepared a stained squash of cells from the tip of an onion root and observed it using an optical microscope.
	During the preparation of the slide, he:
	 cut the first 5 mm from the tip of an onion root and placed it on a glass slide covered this tip with a drop of stain solution and a cover slip warmed the glass slide pressed down firmly on the cover slip.
	He identified and counted nuclei in different stages of the cell cycle.
	Explain why the student: [2 marks]
	1. used only the first 5 mm from the tip of an onion root.
	2. pressed down firmly on the cover slip.
	Question 5 continues on the next page



Turn over ▶

Figure 2 shows the cells the student saw in one field of view. He used this field of view to calculate the length of time these onion cells spent in anaphase of mitosis.

Figure 2



Scientists have found the mean length of time spent by onion cells in anaphase of mitosis is 105 minutes. They also found the cell cycle of cells in the onion root shown in **Figure 2** takes 1080 minutes.

32 whole cells are shown in Figure 2.

Use this information and **Figure 2** to calculate the length of time the cells of this onion root are in anaphase **and** then calculate the percentage difference between your answer and the mean length of time found by the **scientists**.

Show your working.

[2 marks]

Answer =	_ %



	15	
0 5 . 3	Tick (✓) the name given to the division of cytoplasm during the cell cycle.	mark]
	Binary fission	
	Cytokinesis	
	Phagocytosis	
	Segregation	
0 5.4	Describe and explain what the student should have done when counting cells to sure that the mitotic index he obtained for this root tip was accurate. [2 m	make
	Question 5 continues on the next page	

Turn over ►



0	5		5	
U	5	•	5	

A scientist treated growing tips of onion roots with a chemical that stops roots growing. After 24 hours, he prepared a stained squash of these root tips.

Figure 3 is a drawing showing the chromosomes in a single cell observed in the squash of one of these root tips in anaphase. This cell was typical of other cells in anaphase in these root tips.

Figure 3



Use all of this information to suggest how the chemical stops the growth of roc	ots. 3 marks]

10



There are no questions on this page Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

Turn over ▶




0 6

Under the correct conditions, new roots grow from the cut end of a plant stem. A scientist investigated the effect of substance **X** on the growth of new roots.

She used a ringing experiment to investigate the movement of substance \mathbf{X} in stems taken from lemon plants. She cut out a length of stem from each plant. She then put a small block of agar on the top of each length of stem. Some agar blocks contained substance \mathbf{X} .

Figure 4 shows how she treated each length of stem.

Figure 4



She grew the lengths of stem in the same environmental conditions for 6 weeks, and then found the number of roots per length of stem. Roots grew at the other end of the stem from where the agar blocks were placed.

Table 3 shows the scientist's results.

Table 3

Treatment	Mean number of roots per length of stem
D	5
E	11
F	4
G	3



0 6 . 1	Treatment D is a control. Explain how the measurement obtained from this coused by the scientist.	Do not outside bo	de the
0 6.2	Using Figure 4 and Table 3 , what can you conclude from treatments D and I root growth?	E about [3 marks]	





Do not write
outside the
box

0 6 . 3	The mass flow hypothesis is used to explain the movement of substances through phloem.
	Evaluate whether the information from this investigation supports this hypothesis. Do not consider statistical analysis in the answer.
	[4 marks]



0 7.1	What is digestion? [2 marks]
	One species of fungus digests cellulose using two types of enzyme, endocellulases and exocellulases.
	Endocellulases act in the middle of the cellulose molecule and exocellulases act at the ends of the cellulose molecule.
0 7.2	Endocellulases and exocellulases act at different places on cellulose molecules.
	Suggest why. [2 marks]





A scientist prepared the following mixtures:

- 15 g cellulose with 0.2 mol dm⁻³ endocellulase
 15 g cellulose with 0.2 mol dm⁻³ exocellulase
 15 g cellulose with 0.2 mol dm⁻³ endocellulase and 0.2 mol dm⁻³ exocellulase.

The mixtures had identical total volumes. She determined the mass of cellulose remaining after 48 hours.

Her results are shown in Table 4.

Table 4

Mass of cellulose remaining / g		ning / g	
Time / hours	Endocellulase	Exocellulase	Endocellulase + exocellulase
48	11.9	14.8	9.2

0 7 . 3	Use information from Table 4 to calculate the rate of digestion of cellulose when both
	enzymes are present.

Give your answer in g min⁻¹ and in standard form. Show your working.

[2 marks]

	1
Answer =	a min-
Aliswei –	u IIIIII



	23	
0 7.4	The scientist used the same concentration of endocellulase and exocellulase in the mixtures. The rate of digestion of cellulose is greatest when both enzymes are present.	Do not write outside the box
	Suggest why. [2 marks]	
0 7 . 5	The scientist could have expressed her results as the percentage loss in mass of cellulose. In the space, write the equation for calculating the percentage loss in mass.	
	[1 mark]	9
	Turn over for the next question	



0 8.1	A student used a dilution series to investigate the number of cells present in a liq culture of bacteria.	uid
	Describe how he made a 1 in 10 dilution and then used this to make a 1 in 1000 dilution of the original liquid culture of bacteria.	
	[3 m	arks]
	Question 8 continues on the next page	



0 8 . 2	Using an optical microscope, the student determined there were 15 cells in 0.004 mm ³ of the 1 in 1000 dilution of the culture.
	Calculate the number of cells in 1cm³ of undiluted liquid culture. [2 marks]
	Answer = Number of cells
0 8.3	The student looked at cells in the 1 in 10 dilution during his preliminary work. He decided not to use this dilution to determine the number of cells in the undiluted liquid culture.
	Suggest an explanation for the student's decision.
	[2 marks]

Turn over ►



^	0		4
U	Ŏ	١.	4
_	_	-	-

On some farms, animals are routinely given antibiotics in their food.

Scientists investigated whether these farm animals had antibiotic-resistant bacteria in their intestines. They tested the bacteria for resistance to two antibiotics, tetracycline and streptomycin.

Their results are shown in Table 5.

Table 5

Antibiotic	Percentage of antibiotic-resistant bacteria
Tetracycline	29
Streptomycin	13

Suggest and explain one reason why bacteria resistant to tetracycline are more common than bacteria resistant to streptomycin in these farm animals.

[2 marks]



0 8 . 5	In recent years, these farm animals have not been given tetracycline in their food. Despite this, the percentage of bacteria resistant to tetracycline has remained constant.	Do not write outside the box
	Suggest one reason why. [1 mark]	
		10
	Turn over for the next question	





Do	not	writ
ou	tside	e th
	bo	X

Compare and contrast the DNA in eukaryotic cells with the DNA in prokaryot	ic cells. [5 marks]
	Compare and contrast the DNA in eukaryotic cells with the DNA in prokaryot



		1 -
9 . 2	Haemoglobins are chemically similar molecules found in many different species.	Do r outs
	Differences in the primary structure of haemoglobin molecules can provide evidence of phylogenetic (evolutionary) relationships between species.	
	Explain how.	
	[5 marks]	
		1 1

END OF QUESTIONS

There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



Do not write outside the box There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



There are no questions printed on this page DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.



Do not write outside the

box