



**GCE**

**Chemistry B**

**H433/02:** Scientific literacy in chemistry

Advanced GCE

**Mark Scheme for June 2019**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2019

Annotations available in RM Assessor

Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
	Benefit of doubt given
	Contradiction
	Rounding error
	Error in number of significant figures
	Error carried forward
	Level 1
	Level 2
	Level 3
	Benefit of doubt not given
	Noted but no credit given
	Ignore

Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

<b>Annotation</b>	<b>Meaning</b>
<b>DO NOT ALLOW</b>	Answers which are not worthy of credit
<b>IGNORE</b>	Statements which are irrelevant
<b>ALLOW</b>	Answers that can be accepted
( )	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument

## Subject-specific Marking Instructions

### INTRODUCTION


Your first task as an Examiner is to become thoroughly familiar with the material on which the examination depends. This material includes:


- the specification, especially the assessment objectives
- the question paper
- the mark scheme.

You should ensure that you have copies of these materials.

You should ensure also that you are familiar with the administrative procedures related to the marking process. These are set out in the OCR booklet **Instructions for Examiners**. If you are examining for the first time, please read carefully **Appendix 5 Introduction to Script Marking: Notes for New Examiners**.

Please ask for help or guidance whenever you need it. Your first point of contact is your Team Leader.

Question			Answer	Marks	AO element	Guidance
1	(a)		A: (di)acyl chloride ✓ B: (di)amine ✓	2	1.1 x 2	<b>IGNORE</b> arene/benzene/aromatic ring/secondary/ <b>DO NOT ALLOW</b> phenyl/amide/acyl on its own
1	(b)		Angle 120° ✓  three groups/sets of electrons/ 3 areas of electron density (around C) ✓  repel and get as far away as possible/minimise repulsion ✓	3	2.1  2.1  1.1	<b>ALLOW</b> 117 - 122 Mark separately (i.e. no ecf) <b>IGNORE</b> three (bonding) pairs
1	(c)		<b>FIRST CHECK THE ANSWER ON ANSWER LINE</b> <b>If answer = 26 (g) award 2 marks</b>  Amount benzene-1,4-dicarboxylic acid = 32/166 = 0.19 mol ✓  Mass compound <b>A</b> = 0.19 x 0.67 x 203 = 26 (g) (nearest whole number) ✓	2	2.4 x 2	<b>ALLOW</b> ecf from incorrect number of moles
1	(d)		step 1: (conc) ammonia/NH <sub>3</sub> ✓  step 2: Sn + <u>conc</u> HCl /names ✓	2	2.3  2.3	<b>IGNORE</b> heat/reflux/ethanolic but any other additional reagents is CON
1	(e)	(i)	hydrogen (bonds) ✓	1	1.1	
1	(e)	(ii)	 ✓	1	1.1	<b>BOTH</b> dotted lines required but not lone pairs or partial charges
1	(f)	(i)	Heat/ reflux with HCl / H <sub>2</sub> SO <sub>4</sub> / NaOH / acid / alkali / names ✓	1	1.2	<b>DO NOT ALLOW</b> conc. H <sub>2</sub> SO <sub>4</sub>
1	(f)	(ii)	Answer depends on catalysts chosen in (f)(i): alkaline hydrolysis:	2	1.2 x 2	<b>ALLOW</b> salts rather than cation/ anion <b>ALLOW</b> any unambiguous representation


Question	Answer	Marks	AO element	Guidance
	<p data-bbox="360 244 815 323"></p> <p data-bbox="360 331 875 451">acid hydrolysis:</p> <p data-bbox="360 363 875 451"></p>			<p data-bbox="1447 244 2112 379"><b>ALLOW</b> one mark for unionised diamine and dicarboxylic acid <b>IGNORE</b> ambiguous attachments eg OH attached through H</p>

Question			Answer	Marks	AO element	Guidance
2	(a)		<p>electrons raised/excited to higher energy levels (by heat) ✓</p> <p>fall and release energy/visible light/photon ✓</p> <p>frequency of energy/light/photon proportional to gap between energy levels / <math>(\Delta)E = hv</math> ✓</p>	3	1.2 x 3	<b>DO NOT ALLOW</b> answers where energy source is e/m radiation
2	(b)	(i)	<p><b>FIRST CHECK THE ANSWER ON ANSWER LINE</b> <b>If answer = 1260 (cm<sup>3</sup>) award 4 marks</b></p> <p>amount SrCO<sub>3</sub> (= 12.0/147.6) = 0.0813 (mol) ✓</p> <p><math>V = nRT/P</math> ✓</p> <p><math>V = 0.0813 \times 8.314 \times 290 \times 10^6/155000 = 1260</math> (cm<sup>3</sup>) (3 or more sf) ✓</p> <p>Answer to 3sf ✓</p>	4	2.8 x 4	<p><b>ALLOW ECF</b> <b>ALLOW</b> answers rounding to 1260 for 3 marks</p> <p>If values inserted into equation that clearly demonstrates use of MP2 this scores MP2</p> <p><b>ALLOW</b> sf mark for any calculated volume to 3 sf.</p>
2	(b)	(ii)	<p>strontium ions are larger (and attraction less) ✓</p> <p>strontium (ions) have lower charge density ✓</p> <p>they distort/polarise the carbonate (ions) less ✓</p> <p>thermal stability of strontium (carbonate) is higher ✓</p>	4	3.2 x 4	<p><b>ALLOW</b> ora throughout <b>DO NOT ALLOW</b> atomic radius</p> <p><b>ALLOW</b> thermal stability increases down the group.</p>
2	(c)	(i)	46 ✓	1	1.1	
2	(c)	(ii)	<p><b>FIRST CHECK THE ANSWER ON ANSWER LINE</b> <b>If answer = 87.71 award 2 marks</b></p> <p><math>(84 \times 0.56) + (86 \times 9.86) + (87 \times 7.00) + (88 \times 82.58)</math> ✓</p> <p>evaluated as percentage and expressed to 2 dp ✓</p>	2	1.2 x 2	If 2 marks not scored award max 1 mark for any calculated value between 86 and 88 to 2dp.
2	(d)*		<p>Please refer to the marking instructions on page 4 of this mark scheme for guidance on how to mark this question.</p> <p><b>Level 3 (5 – 6 marks)</b> Chooses an appropriate acid concentration.</p>	6	3.4 x 3 3.3 x 3	<p><b>Indicative scientific points include:</b></p> <p><b>Choice of acid concentration</b></p>

Question			Answer	Marks	AO element	Guidance
			<p><b>AND</b> Gives a detailed description, including some fine detail, of procedure. <b>AND</b> Describes how the result would be calculated.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured.</i></p> <p><b>Level 2 (3 – 4 marks)</b> Gives most of the key steps in the procedure, may include some fine detail <b>AND</b> describes how the result would be calculated. <b>OR</b> Addresses all three areas but lacks depth in any of them.</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p><b>Level 1 (1 – 2 marks)</b> A basic description of procedure. <b>OR</b> An attempt to describe the choice of acid concentration. <b>OR</b> An attempt to describe how the result would be calculated.</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b> No response or no response worthy of credit.</p>			<ul style="list-style-type: none"> <li>Calculates conc of <math>\text{Sr}(\text{OH})_{2(\text{aq})} = 0.08 \text{ mol dm}^{-3}</math>;</li> <li>use of reaction stoichiometry 2:1 to determine appropriate concentration of acid to be used (approx. <math>0.15 - 0.2 \text{ mol dm}^{-3}</math>)</li> </ul> <p><b>Practical details</b></p> <ul style="list-style-type: none"> <li>pipette <math>20/25 \text{ cm}^3</math> <math>\text{Sr}(\text{OH})_2/\text{HCl}</math> in a suitable flask;</li> <li>add indicator; (details not required)</li> <li>place acid/alkali in burette;</li> <li>titrate until colour change (details not required)</li> <li>repeat until concordant titres obtained</li> </ul> <p><b>Relevant fine detail</b></p> <ul style="list-style-type: none"> <li>Rinses pipette with solution to be delivered</li> <li>Rinses burette with solution to be delivered</li> <li>Performs a rough titration</li> <li>Add dropwise near to end point</li> </ul> <p><b>Final calculation</b></p> <ul style="list-style-type: none"> <li>Calculates average volume used</li> <li>Use of equation or mole ratio</li> <li>Gives example of suitable relationship to calculate actual concentration eg use of <math>c = n/v</math></li> </ul>
2	(e)	(i)	$\text{Sr}(\text{OH})_2(\text{s}) \rightleftharpoons \text{Sr}^{2+}(\text{aq}) + 2\text{OH}^{-}(\text{aq}) \checkmark$  $K_{\text{sp}} = [\text{Sr}^{2+}] [\text{OH}^{-}]^2 \checkmark$	2	2.2  1.1	Equilibrium can be either way round. Penalise incorrect charge on Sr ions once only
2	(e)	(ii)	<p><b>FIRST CHECK THE ANSWER ON ANSWER LINE</b> <b>If answer = <math>1.6 \times 10^{-4} \text{ mol}^3 \text{ dm}^{-9}</math> award 3 marks</b></p>	3	2.6 x 3	<b>If final answer does not = <math>1.6 \times 10^{-4} \text{ mol}^3 \text{ dm}^{-9}</math></b> <b>ALLOW ECF</b> from (i) provided only Sr and OH ions are involved

Question			Answer	Marks	AO element	Guidance
			$[\text{Sr}^{2+}] = 3.4 \times 10^{-2}$ <b>AND</b> $[\text{OH}^-] = 6.8 \times 10^{-2}$ ✓  $K_{\text{sp}} = 3.4 \times 10^{-2} \times (6.8 \times 10^{-2})^2 = 1.6 \times 10^{-4}$ ✓ units $\text{mol}^3 \text{dm}^{-9}$ ✓			<b>ALLOW ECF</b> from incorrect concentrations of Sr or OH ions, including units as appropriate  <b>ALLOW</b> 2 or more sf <b>ALLOW</b> units derived from an attempt at a worked calculation
2	(e)	(iii)	larger/increased concentration of $\text{OH}^-$ ✓  concentration of $\text{Sr}^{2+}$ reduces in order for $K_{\text{sp}}$ to remain constant <b>AND</b> solubility is lower ✓	2	3.2 x 2	<b>ALLOW</b> more hydroxide ions  <b>ALLOW</b> moves equilibrium to left <b>AND</b> solubility is lower Any reference to $K_{\text{sp}}$ changing is CON
2	(f)	(i)	s(-block) ✓	1	1.1	
2	(f)	(ii)	<b>Any two from:</b> $\text{Sr}^{2+}$ and $\text{Rb}^+$ / Sr loses 2 electrons and Rb loses 1 electron ✓  more (delocalised) electrons in Sr ✓  $\text{Sr}^{2+}$ attracts (more) electrons (in metallic structure) more strongly ✓	2	1.1 x 2	<b>DO NOT ALLOW</b> more outer shell electrons  <b>DO NOT ALLOW</b> references to Sr nuclei

Question			Answer	Marks	AO element	Guidance
3	(a)		$3\text{Cl}_2 + 6\text{OH}^- \rightarrow \text{ClO}_3^- + 5\text{Cl}^- + 3\text{H}_2\text{O}$ Numbers in front of chlorine species ✓ Numbers in front of $\text{OH}^-$ and $\text{H}_2\text{O}$ correct ✓	2	2.5 x 2	<b>ALLOW</b> '1' in front of $\text{ClO}_3^-$ / correct multiples
3	(b)	(i)	$\text{ClO}_3^-/\text{ClO}_2$ is less positive/ more negative than $\text{Cl}_2/\text{Cl}^-$ ✓	2	2.8 x 2	<b>IGNORE</b> 'larger'/'smaller'

			so $\text{ClO}_2$ is oxidised AND $\text{Cl}_2$ is reduced / electrons flow from $\text{ClO}_3^- / \text{ClO}_2$ (ora) / half equations are reversed ✓			<b>ALLOW</b> by reference to one species in either half equation.
3	(b)	(ii)	Larger $[\text{H}^+] / [\text{Cl}^-]$ ✓ Equilibrium / equation 3.1 moves to right ✓	2	3.1 x 2	<b>IGNORE</b> 'more' <b>ALLOW</b> $E^\ominus$ for $\text{ClO}_3^- / \text{ClO}_2$ more positive OR $E^\ominus \text{Cl}_2 / \text{Cl}^-$ becomes more negative
3	(c)	(i)	 <p>voltmeter and salt bridge ✓  <math>\text{Cu}^{2+}(\text{aq})</math> and <math>\text{Cu}(\text{s})</math> ✓  <math>\text{Cl}_2(\text{aq}) / \text{Cl}^-(\text{aq})</math> and Pt / C electrode ✓  solutions <math>1 \text{ mol dm}^{-3}</math> and <math>298 \text{ K}</math> ✓</p>	4	3.4 x 4	<b>IGNORE</b> description of makeup of salt bridge <b>IGNORE</b> '2' in front of ' $\text{Cl}^-$ '  <b>ALLOW</b> Cu and Pt/C without state symbols. <b>ALLOW</b> one mark for points 2 and 3 if all state symbols omitted <b>ALLOW</b> electrodes around the other way If no solution shown in either half cell MP2 <b>OR</b> MP3 cannot score If no solution shown in both half cells only penalise once.
3	(c)	(ii)	1.02 (V) ✓	1	2.8	<b>IGNORE</b> sign
3	(c)	(iii)	<u>in the wire</u> from Cu (ora) ✓	1	2.8	<b>ALLOW</b> movement of electrons correctly labelled on the diagram.
3	(c)	(iv)	$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$	1	1.2	<b>ALLOW</b> equation: •halved •with arrow •other way round

Question			Answer	Marks	AO element	Guidance
3	(c)	(v)	<p><b>FIRST CHECK THE ANSWER ON ANSWER LINE</b>  <b>If answer = +0.28 (V) award 2 marks</b></p> <p><math>\ln 0.01 = -4.6 \checkmark</math>  <math>E = +0.34 - (0.0128 \times 4.6) = +0.28 (V) \checkmark</math></p>	2	2.8 x 2	<p><b>ALLOW</b> 2 or more sf  <b>+ sign essential. (0.28 with no sign = 1)</b></p> <p>lg 0.01 answer is +0.31V for 1 mark only  <b>ALLOW</b> If MP1 not clearly stated then by implication it can be credited from a subsequent calculation eg;  <math>E_{\text{cell}} = 1.02</math> answer is +0.96V for 1 mark only</p>
3	(d)	(i)	$\text{Cl}_2 + 2\text{I}^- \rightarrow 2\text{Cl}^- + \text{I}_2$	1	1.2	<b>IGNORE</b> state symbols
3	(d)	(ii)	iodide(ion)	1	1.2	<b>IGNORE</b> formulae
3	(d)	(iii)	brown/orange/yellow (solution)	1	1.2	<b>ALLOW</b> these colours or any combination but no others. <b>IGNORE</b> reference to starting colour. PPT or (s) is CON
3	(d)	(iv)	Chlorine has a greater attraction for (AW) electrons (than iodine) (ora)	1	2.5	Reference to molecules is CON <b>IGNORE</b> references to electronegativity / attraction to valence electrons
3	(e)		<p>Test tube or flask containing Sodium Chloride and concentrated sulphuric acid <math>\checkmark</math></p> <p>Delivery tube for downward delivery into a test tube or boiling tube <math>\checkmark</math></p>	2	3.3 x 2	<p><b>ALLOW</b> formulae</p> <p>Collection over water, or into a sealed vessel            CONs MP2</p>

Question			Answer	Marks	AO element	Guidance
4	(a)		<p>Oxides of nitrogen/<math>\text{NO}_2</math> is recycled/regenerated/reformed <math>\checkmark</math>            reactions are <math>\text{NO}_2 + \text{O} \rightarrow \text{NO} + \text{O}_2</math> and <math>\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2 \checkmark</math></p>	2	3.1 x 2	
4	(b)	(i)		4	2.8 x 4	

Question			Answer	Marks	AO element	Guidance
			<p>axes round right way and labelled correctly ✓  scale to fill 2/3 of area ✓  plot with line of best fit ✓</p> <p>measurement of one half-life = 1400 s ± 100 ✓</p>			Should be a curve that touches at least 4 points. Mark half-life by answer given, no construction lines needed for <i>this</i> part.
4	(b)	(ii)	'Half lives constant' <b>AND</b> At least two half-lives constructed ✓	1	2.7	
	(c)		(k = $9.8 \times 10^{-5} / 0.210$ ⇒) $4.7 \times 10^{-4}$ ✓ units s <sup>-1</sup> ✓	2	2.4 x 2	<b>ALLOW</b> 2 or more sf Mark units separately
	(d)		<b>FIRST CHECK THE ANSWER ON ANSWER LINE</b> If answer = (+)100±10 (any sf) (kJ mol <sup>-1</sup> ) award 3 marks	3	2.6 x 3	<b>ALLOW</b> one or more sf <b>ALLOW ECF</b>
			slope = -12000±500 ✓  E <sub>a</sub> = 12000 x 8.314 = (+)99768 (J) ✓ = (+)99.8 (kJ mol <sup>-1</sup> ) ✓			MP1 is for calculating the gradient  MP2 is for multiplying by R and evaluating MP3 is for converting from J to kJ
	(e)		(this is a possible mechanism because) reactions add to overall equation / $2\text{N}_2\text{O}_5 \rightarrow 4\text{NO}_2 + \text{O}_2$ ✓	3	3.1 x 3	

Question			Answer	Marks	AO element	Guidance
			<p>step 1 could be rate determining because it uses <math>\text{N}_2\text{O}_5</math> as a reactant / <math>\text{N}_2\text{O}_5</math> decomposes ✓</p> <p>step 3 could be rate determining because it uses <math>\text{N}_2\text{O}_5</math> as a reactant / could be slow compared to steps 1 and 2 ✓</p>			<p><b>ALLOW</b> cannot be step 2 as <math>\text{N}_2\text{O}_5</math> does not appear in the equation for 1 mark if no reference made to either step 1 or step 3.</p> <p><b>ALLOW BOTH</b> step 1 and step 3 could be RDS with a reason scores 2 marks BOTH step 1 and step 3 with no reason scores 1 mark.</p>

Question		Answer	Marks	AO element	Guidance	
5	(a)	$\text{C}_2\text{H}_2(\text{g}) + 2.5\text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g}) + \text{H}_2\text{O}(\text{g/l})$ $\Delta_c H = 2\Delta_f H \text{CO}_2 + \Delta_f H \text{H}_2\text{O} - \Delta_f H \text{C}_2\text{H}_2 \checkmark$	3	2.1  2.1  2.1	<p>First mark for correct elements Second mark for correct <math>\Delta_f H</math> descriptions and top equation <b>ALLOW</b> <math>\Delta_f H \text{2CO}_2</math> <b>IGNORE</b> <math>\Delta_f H \text{O}_2</math></p> <p>Third mark for correct expression for <math>\Delta_c H</math> <i>Allow use of definitions/symbols from enthalpy cycle</i></p>	
5	(b)	<p><b>FIRST CHECK THE ANSWER ON ANSWER LINE</b> <b>If answer = 1.9 (times greater) award 4 marks</b></p> <p>Equation: <math>\text{C}_3\text{H}_8 + 5\text{O}_2 \rightarrow 3\text{CO}_2 + 4\text{H}_2\text{O} \checkmark</math>  <math>5/0.2 = 25</math> (moles 'air') <math>\checkmark</math>  12.5 moles 'air' for acetylene <math>\checkmark</math>  <math>26/13.5 = 1.9</math> (times greater) <math>\checkmark</math></p>	4	2.5 2.5 2.6 2.6	<p><b>ALLOW</b> 3 marks if mole fraction route not used i.e; Correct equation for propane <math>\checkmark</math> Scaled equation for acetylene so that moles of <math>\text{O}_2</math> are identical in both equations / acetylene needs 2.5 moles <math>\text{O}_2</math> and propane needs 5 moles <math>\text{O}_2 \checkmark</math> Ratio of acetylene to propane = 2 identified <math>\checkmark</math></p> <p><b>ALLOW</b> ECF from an incorrect equation</p>	
5	(c)	(i)	Carbon atoms contain 4 outer (shell) electrons $\checkmark$ $\text{sp}^2$ (orbitals) uses 3 electrons $\checkmark$	2	2.1 x 2	
		(ii)	ethene: form a $\pi$ bond $\checkmark$ naphthalene: delocalised/conjugated $\checkmark$	2	1.1 x 2	
5	(d)		Abstraction/removal of hydrogen from naphthalene $\checkmark$	1	2.5	<b>DO NOT ALLOW</b> steps before abstraction <b>IGNORE</b> any further steps that grow PAH
5	(e)	(i)	initiation <b>AND</b> radicals formed (from molecules) $\checkmark$	1	2.1	
		(ii)	Provide energy/break bonds by colliding $\checkmark$	1	3.2	<b>IGNORE</b> reference to catalyst
5	(f)		Please refer to the marking instructions on page 4 of this mark scheme for guidance on how to mark this question.	6	3.1 x 6	<b>Indicative scientific points include:</b> <b>Flame temp:</b>

Question	Answer	Marks	AO element	Guidance
	<p><b>Level 3 (5 – 6 marks)</b>            Gives a detailed account of controlling flame temperature, small molecule reactions and competing reactions, exemplified by the use of at least one appropriate equation.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured.</i></p> <p><b>Level 2 (3 – 4 marks)</b>            Gives an outline account of controlling flame temperature, small molecule reactions and competing reactions.</p> <p><b>OR</b>            Gives a detailed account of two of the following areas, controlling flame temperature, small molecule reactions, or competing reactions</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p><b>Level 1 (1 – 2 marks)</b>            Gives an outline account of two of the following areas, controlling flame temperature, small molecule reactions or competing reactions</p> <p><b>OR</b>            Gives a detailed account of one area</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b>  <i>No response or no response worthy of credit.</i></p>			<ul style="list-style-type: none"> <li>• saturated hydrocarbons need more oxygen per mole</li> <li>• fewer saturated hydrocarbon molecules in the same volume;</li> <li>• hence lower flame temp;</li> <li>• pure oxygen produces higher flame temperatures;</li> <li>• example comparison equations (Allow ORA for arguments in favour of unsaturated hydrocarbons)</li> </ul> <p><b>Small molecule reactions:</b></p> <ul style="list-style-type: none"> <li>• oxygen atoms/ molecules produce hydrogen atoms</li> <li>• use of appropriate equation(s), eg <math>\text{CH} + \text{O} \rightarrow \text{CO} + \text{H}</math> or <math>\text{CH}_2 + \text{O}_2 \rightarrow \text{CO}_2 + 2\text{H}</math></li> <li>• more hydrogen atoms abstracted from growing PAH</li> <li>• balance entropy AW</li> </ul> <p><b>Competing reactions:</b></p> <ul style="list-style-type: none"> <li>• production of acetylene vs <math>\text{CO}_2</math>;</li> <li>• saturated hydrocarbons produce more <math>\text{CO}_2</math>;</li> <li>• unsaturated hydrocarbons produce acetylene;</li> <li>• acetylene leads to soot formation</li> <li>• soot formation vs <math>\text{CO}_2</math> production</li> </ul>

**OCR (Oxford Cambridge and RSA Examinations)**  
**The Triangle Building**  
**Shaftesbury Road**  
**Cambridge**  
**CB2 8EA**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

**Oxford Cambridge and RSA Examinations**  
is a Company Limited by Guarantee  
Registered in England  
Registered Office; The Triangle Building, Shaftesbury Road, Cambridge, CB2 8EA  
Registered Company Number: 3484466  
OCR is an exempt Charity

**OCR (Oxford Cambridge and RSA Examinations)**  
Head office  
Telephone: 01223 552552  
Facsimile: 01223 552553

© OCR 2019

