

Please write clearly in	block capitals.		
Centre number		Candidate number	
Surname			
Forename(s)			
Candidate signature			

AS CHEMISTRY

Paper 2 Organic and Physical Chemistry

Thursday 23 May 2019

Morning

Time allowed: 1 hour 30 minutes

Materials

For this paper you must have:

- the Periodic Table/Data Sheet, provided as an insert (enclosed)
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer the questions in the spaces provided. Do **not** write outside the box around each page or on blank pages.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

Advice

You are advised to spend about 65 minutes on Section A and 25 minutes on Section B.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
Section B	
TOTAL	



Do not write
outside the
L

Section A

Answer all questions in this section.

1 The structures of three organic compounds A, B and C are shown.

These compounds can be distinguished by simple test-tube reactions.

For each pair of compounds in questions **01.1** and **01.2**, give a reagent (or combination of reagents) that could be added separately to each compound to distinguish between them.

State what is observed in each case.

0 1 . 1	Compounds A and B	[3 marks]
	Reagent	

Observation with A

Observation with **B**

0 1 . 2 Compounds A and C [3 marks]

Reagent _____
Observation with A _____

Observation with C

6



		Do not write
0 2	Bromoethane reacts with potassium cyanide to form compound D .	outside the box
	$CH_3CH_2Br + KCN \rightarrow CH_3CH_2CN + KBr$	
	Compound D	
0 2 . 1	Outline the mechanism for this reaction.	
	[2 marks]	
0 2.2	Give the IUPAC name of D . [1 mark]	
	[············	
0 2 . 3	Calculate the percentage atom economy for the formation of D in this reaction.	
	Give your answer to the appropriate number of significant figures.	
	[2 marks]	
	% atom economy	5
	70 atom coondiny	

Turn over ▶



0 3

This question is about enthalpy changes.

0 3 . 1

A student determined the enthalpy of combustion of cyclohexane (C₆H₁₂).

The student

- placed a pure sample of cyclohexane in a spirit burner
- placed the spirit burner under a beaker containing 50.0 g of water and ignited the cyclohexane
- extinguished the flame after a few minutes.

The results for the experiment are shown in **Table 1**.

Table 1

Initial temperature of the water / °C	19.1
Initial mass of spirit burner and cyclohexane / g	192.730
Final mass of spirit burner and cyclohexane / g	192.100

The student determined from this experiment that the enthalpy of combustion of cyclohexane is $-1216 \text{ kJ} \text{ mol}^{-1}$

Use the data to calculate the final temperature of the water in this experiment.

The specific heat capacity of water = 4.18 J K⁻¹ g⁻¹ The relative molecular mass (M_r) of cyclohexane = 84.0

[4 marks]



		Do not write outside the box
	Final temperature of the water°C	
0 3.2	A data book value for the enthalpy of combustion of cyclohexane is –3920 kJ mol ⁻¹ The student concluded that the temperature rise recorded in the experiment was smaller than it should have been. Suggest a practical reason for this.	
	[1 mark] - -
	Question 3 continues on the next page	



0 3 . 3

Table 2 gives some values of standard enthalpies of combustion ($\Delta_c H^e$).

Table 2

Substance	C(s)	H ₂ (g)	C ₆ H ₁₂ (I)
Standard enthalpy of combustion, $\Delta_{\rm c} H^{\rm e}$ / kJ mol ⁻¹	-394	-286	-3920

Use the data in **Table 2** to calculate the enthalpy change for the reaction represented by this equation

$$6C(s) + 6H_2(g) \rightarrow C_6H_{12}(I)$$

[3 marks]

Enthalpy change _____kJ mol⁻¹

8

This question is about fossil fuels.

The petrol fraction from crude oil contains octane (C₈H₁₈).

Give an equation for the complete combustion of octane.

[1 mark]

The combustion of petrol in car engines produces the pollutant nitrogen monoxide.

Give an equation for a reaction that removes nitrogen monoxide in a catalytic converter.

[1 mark]

Question 4 continues on the next page

Turn over ▶



Do not write
outside the

0	4		3
U	4	-	၁

Sulfur dioxide is produced in the combustion of fossil fuels. The total emissions of sulfur dioxide in the UK have fallen dramatically since 1970.

Sulfur dioxide is now removed from the flue gases in power stations by reaction with calcium oxide.

$$CaO + SO_2 \rightarrow CaSO_3$$

In 1970, the total UK emissions of sulfur dioxide were 6.49 million tonnes (1 tonne = 1000 kg).

Calculate the mass, in kilograms, of calcium oxide needed to react with this mass of sulfur dioxide.

Give your answer in standard form.

[2 marks]

Mass of calcium oxide	ka
mass or carolam shas	9



0 5	Methanol (CH₃OH) is an important alcohol with many uses.
0 5 . 1	Draw a diagram to show how two methanol molecules interact with each other through hydrogen bonding in the liquid phase.
	Include all partial charges and all lone pairs of electrons in your diagram. [3 marks]
0 5 . 2	The bond angle around the oxygen atom in methanol is slightly smaller than the regular tetrahedral angle of 109.5°
	Explain why this bond angle is smaller than 109.5° [1 mark]





0 5.3	Methanol is made by the reaction of carbon monoxide with hydrogen.	
	$CO + 2H_2 \rightleftharpoons CH_3OH$ $\Delta H = -91 \text{ kJ mol}^{-1}$	
	The reaction uses a copper-based catalyst, a pressure of 10 MPa and a temperature of 550 K	
	These conditions are used to provide a balance between equilibrium yield, reaction rate and cost.	
	Describe how the use of a catalyst, and changes in pressure and temperature each affect equilibrium yield, reaction rate and cost.	
	L'	6 marks]



-

10



0	6
---	---

Propene reacts with concentrated sulfuric acid to form two isomers, **E** and **F**.

The structure of **E** is shown.

$$CH_3 - CH - CH_3$$
 O
 $O = S = O$
 O
 O

0 6 . 1

Name and outline the mechanism for the formation of **E** in this reaction.

[5 marks]

Name of mechanism

Mechanism



0 6 . 2	Draw the structure of F .	Do not write outside the box
	[1 mark]	
0 6.3	Explain why more of isomer E than isomer F is formed in this reaction.	
	[2 marks]	
		8
	Turn over for the next question	

Turn over ▶



0 7	Propanedioic acid contains two carboxylic acid groups. It is a solid organic acid that is soluble in water.
0 7.1	Draw the skeletal formula of propanedioic acid. [1 mark]
0 7.2	Describe how to prepare 250 cm³ of an aqueous standard solution of propanedioic acid containing an accurately measured mass of the acid. Include essential practical details in your answer. [6 marks]



0 7.3	Calculate the mass, in mg, of propanedioic acid (<i>M_r</i> = 104.0) needed to prepare 250 cm ³ of a 0.00500 mol dm ⁻³ solution. [2 m	po not wroutside the box	ite
	Mass of propanedioic acid	_mg 9	

Turn over ▶



Do not write
outside the

0	8
---	---

Propanal can be prepared by the oxidation of propan-1-ol with acidified potassium dichromate(VI).

An ionic equation for this reaction is

 $3CH_3CH_2CH_2OH + Cr_2O_7^{2-} + 8H^+ \rightarrow 3CH_3CH_2CHO + 2Cr^{3+} + 7H_2O$

0 8 . 1

Calculate the minimum volume, in cm³, of 0.40 mol dm⁻³ potassium dichromate(VI) solution needed to oxidise 6.0 cm³ of propan-1-ol to propanal.

 $M_{\rm r}$ of propan-1-ol = 60.0 Density of propan-1-ol = 0.80 g cm⁻³

[3 marks]

Minimum volume cm³



0 8 . 2 The reaction is done in a pear-shaped flask.

Complete the diagram to show the assembled apparatus needed to prepare propanal from propan-1-ol in this way.

Label the diagram.

[3 marks]

Do not write outside the




Turn over for the next question

Turn over ▶



IB/G/Jun19/7404/2


6





|--|

The compound 1,2-dichlorotetrafluoroethane is a CFC that was previously used in refrigerators as a coolant.



0 9

Molecules of 1,2-dichlorotetrafluoroethane can break down in the upper atmosphere to form chlorine radicals.

Give an equation to show the breakdown of one molecule of 1,2-dichlorotetrafluoroethane to form one chlorine radical and one other species.

[1 mark]

0	9	2

Give **two** equations to show how chlorine radicals catalyse the decomposition of ozone.

[2 marks]

Question 9 continues on the next page

Turn over ▶



Do not write
outside the
hav

0	9	. 3	Butane can be used as a replacement for CFCs in refrigerators.

During its use, the butane is repeatedly converted from liquid to gas and then back to liquid. Liquid butane expands as it turns into a gas.

- Calculate the volume, in cm³, of 38.8 g of butane gas at 272 K and 101 kPa (the gas constant R = 8.31 J K⁻¹ mol⁻¹) (M_r of butane = 58.0)
- Calculate the volume, in cm³, of 38.8 g of liquid butane. (density of liquid butane = 0.60 g cm⁻³)
- Use your answers to calculate the factor by which butane expands in volume when it changes from a liquid to a gas.

Show your working.

[6 marks]

Volume of butane gas cm³




		Do not write outside the box
Volume of liquid butane	_cm³	
		9
Expansion factor	_	





Se	cti	on	В
-	~	•••	_

Answer all questions in this section. Only **one** answer per question is allowed. For each answer completely fill in the circle alongside the appropriate answer. CORRECT METHOD WRONG METHODS **∞** ● 👄 If you want to change your answer you must cross out your original answer as shown. If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown. You may do your working in the blank space around each question but this will not be marked. Do **not** use additional sheets for this working. 1 0 A 'drink-driving' offence is committed if the blood alcohol level of a driver is over 80 mg of ethanol per 100 cm³ of blood. What is the concentration, in mol dm⁻³, of ethanol if there are 80 mg of ethanol $(M_r = 46.0)$ per 100 cm³ of blood? [1 mark] **A** 0.00017 **B** 0.0017 **C** 0.017 **D** 1.7





1 1	Which statement is correct for the distribution curve of molecular energy	gies in a	a gas? [1 mark]
	A The curve is symmetrical about the maximum.	0	
	B There are always some molecules with zero energy.	0	
	$\boldsymbol{c}\ $ The position of the maximum of the curve is not dependent on the temperature.	0	
	The mean energy of the molecules is greater than the most probable energy of the molecules.	0	
1 2	When one mole of ammonia is heated to a given temperature, 50 % of and the following equilibrium is established.	it disso	ociates
	$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) + \frac{3}{2}H_2(g)$		
	What is the total amount, in moles, of gas in this equilibrium mixture?		[1 mark]
	A 1.5	0	
	B 2.0	0	
	C 2.5	0	
	D 3.0	0	
1 3	Which compound is not an isomer of the following compound?		
	0		[1 mark]
	A CH₃CH₂COCH₃	0	
	B CH ₃ CH=CHCH ₂ OH	0	
	C (CH ₃) ₂ CHCHO	0	
	D CH ₂ =CHCH ₂ CHO	0	

Turn over ▶



1 4	How many isomers are there of C₃H₃N?		[1 mark]
	A 2	0	
	B 3	0	
	C 4	0	
	D 5	0	
1 5	Which equation represents a propagation step?		[1 mark]
	$\mathbf{A} \cdot \mathbf{CH_2Cl} + \mathbf{Cl} \cdot \rightarrow \mathbf{CH_2Cl_2}$	0	
	$\mathbf{B} \cdot \mathbf{CH}_3 + \mathbf{CH}_3 \rightarrow \mathbf{C}_2\mathbf{H}_6$	0	
	\mathbf{C} $\mathrm{Cl}_2 \to \mathrm{Cl} \bullet + \mathrm{Cl} \bullet$	0	
	D CH ₃ Cl + Cl• → •CH ₂ Cl + HCl	0	
1 6	Which compound can react with ammonia to produce propylamine?		
			[1 mark]
	A CH ₃ CH=CH ₂	0	
	B CH ₃ CH ₂ CH ₂ OH	0	
	C CH ₃ CH ₂ CH ₂ Br	0	
	D CH ₃ CH ₂ CH ₃	0	



1 7 Which statement is not correct about CH ₂ =C(CH ₃)CH ₂ Br?	[1 mark]	Do not write outside the box
A It displays <i>E-Z</i> isomerism.	0	
B It forms an addition polymer.	0	
C It reacts with electrophiles.	0	
D It decolourises bromine water.	0	
1 8 Which compound can be oxidised to form (CH ₃) ₂ CHCOCH ₃ ?	[1 mark]	
A 2-methylpropan-1-ol	0	
B 2,2-dimethylpropanol	0	
C 2-methylbutan-2-ol	0	
D 3-methylbutan-2-ol	0	
1 9 Which species can act as a nucleophile?	[1 mark]	
A NH ₄ ⁺	0	
B CH₃OH	0	
C CH ₄	0	
D H ⁺	0	

Turn over ►



				Do not write
2 0	Which alcohol forms a mixture of alkenes when dehydrated?		[1 mark]	outside the box
	A propan-1-ol	0		
	B propan-2-ol	0		
	C pentan-1-ol	0		
	D pentan-2-ol	0		
2 1	Which compound has the highest boiling point?		[1 mark]	
	A CH ₃ CH ₂ CH ₂ Br	0		
	B CH ₃ CH ₂ CH ₂ F	0		
	C CH ₃ CH ₂ CHO	0		
	D CH ₃ CH ₂ COOH	0		
2 2	Which compound could not be produced by reacting 2-bromo-3-meth sodium hydroxide?		e with	
	A 2-methylbut-1-ene	0		
	B 3-methylbut-1-ene	0		
	C 2-methylbut-2-ene	0		
	D 3-methylbutan-2-ol	0		



2 3 The infrared spectrum of an organic compound is shown. 100 Transmittance 50 / % 3000 500 2000 1500 1000 4000 Wavenumber / cm⁻¹ Which compound produces this spectrum? [1 mark] A ethanoic acid 0 **B** 4-hydroxybutanone C propan-1-ol D prop-2-en-1-ol

Turn over for the next question

2 7

Turn over ▶

2 4

The heat released when 1.00 g of ethanol (M_r = 46.0) undergoes complete combustion is 29.8 kJ

Do not write outside the

15

What is the heat released by each molecule, in joules, when ethanol undergoes complete combustion?

(the Avogadro constant $L = 6.022 \times 10^{23} \text{ mol}^{-1}$)

[1 mark]

A $2.28 \times 10^{-18} \,\mathrm{J}$

0

B $4.95 \times 10^{-20} \text{ J}$

0

C $2.28 \times 10^{-21} \text{ J}$

0

D $4.95 \times 10^{-23} \text{ J}$

0

END OF QUESTIONS

Copyright Information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.





IB/G/Jun19/7404/2