

Please write clearly in block capitals.	
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	

A-level **MATHEMATICS**

Paper 1

Wednesday 5 June 2019

Morning

Time allowed: 2 hours

Materials

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer each question in the space provided for that question.
 If you require extra space, use an AQA supplementary answer book; do not use the space provided for a different question.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
TOTAL	



Answer all questions in the spaces provided.

1 Given that a > 0, determine which of these expressions is **not** equivalent to the others.

Circle your answer.

[1 mark]

$$-2\log_{10}\left(\frac{1}{a}\right)$$
 $2\log_{10}(a)$ $\log_{10}(a^2)$ $-4\log_{10}(\sqrt{a})$


Given $y = e^{kx}$, where k is a constant, find $\frac{dy}{dx}$ 2

Circle your answer.

[1 mark]

$$\frac{dy}{dx} = e^{kx} \qquad \qquad \frac{dy}{dx} = ke^{kx} \qquad \qquad \frac{dy}{dx} = kxe^{kx-1} \qquad \qquad \frac{dy}{dx} = \frac{e^{kx}}{k}$$

3 The diagram below shows a sector of a circle.



The radius of the circle is 4 cm and $\theta = 0.8$ radians.

Find the area of the sector.

Circle your answer.

[1 mark]

$$1.28\,\mathrm{cm}^2$$
 $3.2\,\mathrm{cm}^2$ $6.4\,\mathrm{cm}^2$ $12.8\,\mathrm{cm}^2$

$$3.2 \, \text{cm}^2$$



Do	not	writ
ou	tside	e the

4	The point A has coordinates $(-1, a)$ and the point B has coordinates $(3, b)$	
	The line AB has equation $5x + 4y = 17$	
	Find the equation of the perpendicular bisector of the points A and B.	[4 marks]

Turn over for the next question



Turn over ▶

Do n	ot	writ
outs	ide	the
	box	(

5	An arithmetic sequence has first term a and common difference d .	
	The sum of the first 16 terms of the sequence is 260	
5 (a)	Show that $4a + 30d = 65$	[2 marks]
5 (b)	Given that the sum of the first 60 terms is 315, find the sum of the first 41 t	erms. [3 marks]



Do	not	writ
ou	tside	e the
	ho	Y

S_n is the sum of the first n terms of the sequence.	
Explain why the value you found in part (b) is the maximum value of \mathcal{S}_n	[2 marks]
	
	
Turn over for the next question	



Do	not	write
ou	tside	e the
	L -	

6	The function f is defined by	
	$f(x) = \frac{1}{2}(x^2 + 1), x \ge 0$	
6 (a)	Find the range of f.	[1 mark]
6 (b) (i)	Find $f^{-1}(x)$	[3 marks]
6 (b) (ii)	State the range of $f^{-1}(x)$	[1 mark]



$y = f^{-1}(x)$	[
Find the coordinates of the point of intersection of the graphs of $y = f^{-1}(x)$	y = f(x) and
	[2
Turn over for the next question	




7 (a) By sketching the graphs of $y = \frac{1}{x}$ and $y = \sec 2x$ on the axes below, show that the equation

$$\frac{1}{x} = \sec 2x$$

has exactly one solution for x > 0

[3 marks]



7 (b) By considering a suitable change of sign, show that the solution to the equation lies between 0.4 and 0.6

[2	marks]
L	ılıal kəj

7 (c) Show that the equation can be rearranged to give

$$x = \frac{1}{2}\cos^{-1}x$$

[2 marks]



Do	not	writ
ou	tside	e the
	ho	~

7 (d) (i) Use the iterative formula


$$x_{n+1} = \frac{1}{2} \cos^{-1} x_n$$

with $x_1 = 0.4$, to find x_2 , x_3 and x_4 , giving your answers to four decimal places.

[2 marks]

7 (d) (ii) On the graph below, draw a cobweb or staircase diagram to show how convergence takes place, indicating the positions of x_2 , x_3 and x_4 .

[2 marks]







8	$P(n) = \sum_{k=0}^{n} k^3 - \sum_{k=0}^{n-1} k^3$ where <i>n</i> is a positive integer.	
8 (a)	Find P(3) and P(10)	[2 marks]
0 /h)	Solve the equation $D(n) = 4.25 \times 408$	
8 (b)	Solve the equation $P(n) = 1.25 \times 10^8$	[2 marks]



Do not v	vrite
outside	the
box	

9	Prove that the sum of a rational number and an irrational number is always irrational. [5 marks]
	Turn over for the next question



Turn over ▶

Do	not	write
ou	tside	e the
	L	

Show that the rate of to r^2		· •	•	
	Volume of a sp	where $=\frac{4}{3}\pi r^3$		
		3		[-




Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

Turn over ▶

Jodie is attempting to use differentiation from first principles to prove that the gradient of $y = \sin x$ is zero when $x = \frac{\pi}{2}$

Jodie's teacher tells her that she has made mistakes starting in Step 4 of her working. Her working is shown below.



Step 1 Gradient of chord
$$AB = \frac{\sin\left(\frac{\pi}{2} + h\right) - \sin\left(\frac{\pi}{2}\right)}{h}$$

Step 2
$$= \frac{\sin\left(\frac{\pi}{2}\right)\cos\left(h\right) + \cos\left(\frac{\pi}{2}\right)\sin\left(h\right) - \sin\left(\frac{\pi}{2}\right)}{h}$$

Step 3
$$= \sin\left(\frac{\pi}{2}\right) \left(\frac{\cos\left(h\right) - 1}{h}\right) + \cos\left(\frac{\pi}{2}\right) \frac{\sin\left(h\right)}{h}$$

Step 4 For gradient of curve at A,

let h = 0 then

$$\frac{\cos(h)-1}{h}=0 \text{ and } \frac{\sin(h)}{h}=0$$

Step 5 Hence the gradient of the curve at A is given by

$$\text{sin}\Big(\frac{\pi}{2}\Big)\times 0 + \text{cos}\Big(\frac{\pi}{2}\Big)\times 0 = 0$$

Do	not	writ	(
ou	tside	e the	9
	ho	x	

Complete Steps 4	4 and 5 of Jodie's working below, to correct her proof.	[4
Step 4	For gradient of curve at A,	[4 marl
Step 5	Hence the gradient of the curve at A is given by	
отер о	Therice the gradient of the curve at A is given by	
	Turn over for the next question	



Do not write
outside the
have

12 (a)	Show that the equation	
	$2\cot^2 x + 2\csc^2 x = 1 + 4\csc x$	
	can be written in the form	
	$a\csc^2 x + b\csc x + c = 0$	[2 marks]



Do	not	write
ou	tside	the
	ho	·

12 (b)	Hence, given x is obtuse and	
	$2\cot^2 x + 2\csc^2 x = 1 + 4\csc x$	
	find the exact value of $tan x$	
	Fully justify your answer.	[5] o.ul.o.1
		[5 marks]
	Turn over for the next question	



Do	not	write
ou	tside	the
	ho	~

13	A curve, C, has equation	
	$y = \frac{e^{3x-5}}{x^2}$	
	Show that C has exactly one stationary point.	
	Fully justify your answer.	[7 marks]




Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



Turn over ▶

The graph of $y = \frac{2x^3}{x^2 + 1}$ is shown for $0 \le x \le 4$



Caroline is attempting to approximate the shaded area, A, under the curve using the trapezium rule by splitting the area into n trapezia.

14 (a) When n = 4

14 (a) (i) State the number of ordinates that Caroline uses.

[1 mark]

14 (a) (ii) Calculate the area that Caroline should obtain using this method.

Give your answer correct to two decimal places.

[3 marks]

		Do not write
		outside the box
	-	
	·	
14 (b)	Show that the exact area of A is	
	16 — In 17	
	Fully justify your analysis	
	Fully justify your answer. [5 marks]	
	[3 marks]	
	Overtion 44 continues on the most very	
	Question 14 continues on the next page	



14 (c)	Explain what would happen to Caroline's answer to part (a)(ii) as $n \to \infty$	[1 mark]



Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



Turn over ▶

Do n	ot	write
outs	ide	the
	hav	,

15 (a)	At time t hours after a high tide , the height, h metres, of the tide and the velocity, v knots, of the tidal flow can be modelled using the parametric equations
	$v = 4 - \left(\frac{2t}{3} - 2\right)^2$
	$h=3-2\sqrt[3]{t-3}$
	High tides and low tides occur alternately when the velocity of the tidal flow is zero.
	A high tide occurs at 2 am.
15 (a) (i)	Use the model to find the height of this high tide. [1 mark]
15 (a) (ii)	Find the time of the first low tide after 2 am. [3 marks]

15 (a) (iii) Find the height	t of this low tide.
------------------------------	---------------------

[1 mark]



15 (b)	Use the model to find the height of the tide when it is flowing with maximum	velocity. [3 marks]
4= ()		
15 (c)	Comment on the validity of the model.	[2 marks]
	Turn over for the next question	




Do	not	writ
ou	tside	e the
	ho	~

$y = e^{-x}(\sin x + \cos x)$	$\operatorname{os} x$)	
Find $\frac{\mathrm{d}y}{\mathrm{d}x}$		
Simplify your answ	wer.	[3
		ုဒ
Hence, show that		
Hence, show that	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	
Hence, show that where a is a ratio	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2
	$\int e^{-x} \sin x dx = ae^{-x} (\sin x + \cos x) + c$	[2



16 (c) A sketch of the graph of $y = e^{-x} \sin x$ for $x \ge 0$ is shown below.

The areas of the finite regions bounded by the curve and the x-axis are denoted by $A_1,\,A_2,\,...,\,A_n,\,...$



16 (c) (i) Find the exact value of the area A_1

[3 marks]

Question 16 continues on the next page





Do not v	vrit
outside	the
hov	

16 (c) (ii)	Show that		
		A2 -	
		$\frac{A_2}{A_1} = e^{-\pi}$	
		'	[4 marks]
	,		



16 (c) (iii) Given to	that
-----------------------	------

$$\frac{A_{n+1}}{A_n} = e^{-\pi}$$


show that the exact value of the total area enclosed between the curve and the x-axis is

$$\frac{1+e^\pi}{2(e^\pi-1)}$$


[4 marks]

END OF QUESTIONS











There are no questions printed on this page

Do not write outside the box



Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.



