

Please write clearly in block capitals.	
Centre number	Candidate number
Surname	
Forename(s)	
Candidate signature	

# A-level **MATHEMATICS**

Paper 3

Friday 14 June 2019

Afternoon

Time allowed: 2 hours

#### **Materials**

- You must have the AQA Formulae for A-level Mathematics booklet.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

#### Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer all questions.
- You must answer each question in the space provided for that question.
   If you require extra space, use an AQA supplementary answer book; do not use the space provided for a different question.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

## Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

#### Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use		
Question	Mark	
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
12		
13		
14		
15		
16		
17		
TOTAL		



## **Section A**

Answer all questions in the spaces provided.

1  $f(x) = \arcsin x$ 

State the maximum possible domain of f

Tick (✓) one box.

[1 mark]

$$\{x \in \mathbb{R}: -1 \le x \le 1\}$$



$$\left\{x \in \mathbb{R}: -\frac{\pi}{2} \le x \le \frac{\pi}{2}\right\}$$



$${x \in \mathbb{R} : -\pi \le x \le \pi}$$



$${x \in \mathbb{R} : -90 \le x \le 90}$$



Find the value of  $\frac{100!}{98! \times 3!}$ 2

Circle your answer.

[1 mark]

3 Given  $u_1 = 1$ , determine which one of the formulae below defines an increasing sequence for  $n \ge 1$ 

Circle your answer.

[1 mark]

$$u_{n+1}=1+\frac{1}{u_n}$$

$$u_{n+1} = 1 + \frac{1}{u_n}$$
  $u_n = 2 - 0.9^{n-1}$   $u_{n+1} = -1 + 0.5u_n$   $u_n = 0.9^{n-1}$ 

$$u_{n+1} = -1 + 0.5u_n$$

$$u_n=0.9^{n-1}$$




4 Sketch the region defined by the inequalities

$$y \le (1 - 2x)(x + 3)$$
 and  $y - x \le 3$ 

Clearly indicate your region by shading it in and labelling it R.

[3 marks]



Turn over for the next question




Turn over ▶

**5** A circle has equation  $x^2 + y^2 - 6x - 8y = 264$ 

AB is a chord of the circle.

The angle at the centre of the circle, subtended by AB, is 0.9 radians, as shown in the diagram below.



Find the area of the minor segment shaded on the diagram.

Give your answer to three significant figures.

	[5 marks]
·	



Do not write
outside the
box

6	The three sides of a right-angled triangle have lengths $a, b$ and $c$ , where $a, b, c \in \mathbb{Z}$	
	C	
	a	
	b	
• ( )		
6 (a)	State an example where $a,\ b$ and $c$ are all even. [1 mark]	
	[1 mark]	
6 (b)	Prove that it is <b>not</b> possible for all of $a$ , $b$ and $c$ to be odd.	
0 (b)	[3 marks]	
	Turn over for the next question	



Do	not	write
ou	tside	e the
	L	

7 (a)	Express $\frac{4x+3}{(x-1)^2}$ in the form $\frac{A}{x-1} + \frac{B}{(x-1)^2}$	[3 marks]
		[o marko]



Do	not	write
ou	tside	the t
	40	

7 (b)	Show that	
	$\int_{3}^{4} \frac{4x+3}{(x-1)^{2}}  \mathrm{d}x = p + \ln q$	
	where $p$ and $q$ are rational numbers.	[5 marks]
	Turn over for the next question	
	Tam ever to the next queetion	



Turn over ▶

Do	not)	writ
οι	ıtside	e the
	ho	

8	A student is conducting an experiment in a laboratory to investigate how quickly liquids cool to room temperature.
	A beaker containing a hot liquid at an initial temperature of 75 °C cools so that the temperature, $\theta$ °C, of the liquid at time $t$ minutes can be modelled by the equation
	$\theta = 5(4 + \lambda e^{-kt})$
	where $\lambda$ and $k$ are constants.
	After 2 minutes the temperature falls to 68 °C.
8 (a)	Find the temperature of the liquid after 15 minutes.
	Give your answer to three significant figures.  [7 marks]



Do	not	writ
οι	ıtside	e the

8 (b) (i)	Find the room temperature of the laboratory, giving a reason for your answer	er. [2 marks]
8 (b) (ii)	Find the time taken in minutes for the liquid to cool to 1 °C above the room	
	temperature of the laboratory.	[2 marks]
8 (c)	Explain why the model might need to be changed if the experiment was cor a different place.	nducted in
	a different place.	[1 mark]
	Turne Assau San Alan manda mara di su	
	Turn over for the next question	



ı	Do	not	write
	ou	tside	e the
		ho	~

9	A curve has equation	
	$x^2y^2 + xy^4 = 12$	
9 (a)	Prove that the curve does not intersect the coordinate axes.	[2 marks]
9 (b) (i)	Show that $\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{2xy + y^3}{2x^2 + 4xy^2}$	[5 marks]



(b) (ii)	Prove that the curve has no stationary points.  [4 marks	Do no outsi b
		-
		-
		-
		-
		-
	,	-
		-
		-
		_
		-
b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - 5]
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - <b>3]</b>
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - - - -
(b) (iii)	In the case when $x>0$ , find the equation of the tangent to the curve when $y=1$ [4 marks	- - - - - - - - -



Turn over ▶

	Section B	Do not write outside the box						
	Answer <b>all</b> questions in the spaces provided.							
10	Which of the options below best describes the correlation shown in the diagram below?							
	y × ×							
	× × ×							
	Tick ( $\checkmark$ ) one box. [1 mark]							
	moderate positive							
	strong positive							
	moderate negative							
	strong negative							



Lenny is one of a team of people interviewing shoppers in a town centre.

He is asked to survey 50 women between the ages of 18 and 29

Identify the name of this type of sampling.

Circle your answer.

[1 mark]

simple random stratified quota systematic

Turn over for the next question



12	Amelia	decides	to	analyse	the	heights	of	members	of	her	school	rowing	club.
----	--------	---------	----	---------	-----	---------	----	---------	----	-----	--------	--------	-------

The heights of a random sample of 10 rowers are shown in the table below.

Rower	Jess	Nell	Liv	Neve	Ann	Tori	Maya	Kath	Darcy	Jen
Height (cm)	162	169	172	156	146	161	159	164	157	160

12 (a)	Any value more than 2 standard deviations from the mean may be regarded as an outlier.
	Verify that Ann's height is an outlier.
	Fully justify your answer.  [4 marks]



Do	not	write
ou	tside	the
	ho	~

12 (b)	Amelia thinks she may have written down Ann's height incorrectly.
	If Ann's height were discarded, state with a reason what, if any, difference this would
	make to the mean and standard deviation.  [2 marks]
	[2 marks
	Turn over for the next question



13	Patrick is practising his skateboarding skills. On each day, he has 30 attempts at performing a difficult trick.
	Every time he attempts the trick, there is a probability of 0.2 that he will fall off his skateboard.
	Assume that the number of times he falls off on any given day may be modelled by a binomial distribution.
13 (a) (i)	Find the mean number of times he falls off in a day.  [1 mark]
13 (a) (ii)	Find the variance of the number of times he falls off in a day.  [1 mark]
13 (b) (i)	Find the probability that, on a particular day, he falls off exactly 10 times.  [2 marks]



13 (b) (ii)	Find the probability that, on a particular day, he falls off 5 or more times.  [3 ma	irks]
13 (c)	Patrick has 30 attempts to perform the trick on each of 5 consecutive days.	
13 (c) (i)	Calculate the probability that he will fall off his skateboard at least 5 times on each the 5 days.	
	[2 ma	
13 (c) (ii)	Explain why it may be unrealistic to use the same value of 0.2 for the probability of falling off for all 5 days.  [1 m	
	Turn over for the next question	



The survey recorded whether or not they had suffered from a range of four health issues in the past year.

In addition, their physical exercise level was categorised as low, medium or high.

50 teachers had a low exercise level, 40 teachers had a medium exercise level and 30 teachers had a high exercise level.

The results of the survey are shown in the table below.

	Low exercise	Medium exercise	High exercise
Back trouble	14	7	10
Stress	38	14	5
Depression	9	2	1
Headache/Migraine	4	5	5

**14 (a)** Find the probability that a randomly selected teacher:

14 (a) (i)	suffers from back trouble and has a high exercise level;	[1 mark]
14 (a) (ii)	suffers from depression.	[2 marks]



14	(a) (iii)	suffers from stress, given that they have a low exercise level.	[2 marks]
14	(b)	For teachers in the survey with a low exercise level, explain why the events from back trouble' and 'suffers from stress' are <b>not</b> mutually exclusive.	'suffers
		ment sask heaste and canere hem chees are net mataan, executioner	[2 marks]
		Turn over for the next question	



Jamal, a farmer, claims that the larger the rainfall, the greater the yield of wheat from his farm.

He decides to investigate his claim, at the 5% level of significance.

He measures the rainfall in centimetres and the yield in kilograms for a random sample of ten years.

He correctly calculates the product moment correlation coefficient between rainfall and yield for his sample to be 0.567

The table below shows the critical values for correlation coefficients for a sample size of 10 for different significance levels, for both 1- and 2-tailed tests.

1-tailed test significance level	5%	2.5%	1%	0.5%
2-tailed test significance level	10%	5%	2%	1%
Critical value	0.549	0.632	0.716	0.765

Determine what Jamal's conclusion to his investigation should be, justifying your answer.


		[3 marks]
		······································
		······································



Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

Turn over ▶

16 (a) The graph below shows the amount of salt, in grams, purchased per person per week in England between 2001–02 and 2014, based upon the Large Data Set.



Meera and Gemma are arguing about what this graph shows.

Meera believes that the amount of salt consumed by people decreased greatly during this period.

Gemma says that this is not the case.

Using your knowledge of the Large Data Set, give **two** reasons why Gemma may be correct.

		[2 marks



Do not write
outside the
box

	Assume that the survey data is a random sample taken from a normal distribution and that the standard deviation has remained the same.  [6 marks]
	Investigate, at the 5% level of significance, whether the mean amount of sugar purchased per person in England has changed between 2014 and 2018.
	In 2018, a sample of 918 people had a mean of 80.4 grams of sugar purchased per person.
16 (b)	It is known that the mean amount of sugar purchased per person in England in 2014 was 78.9 grams, with a standard deviation of 25.0 grams.



16 (c)	Another test is performed to determine whether the mean amount of fat purchased				
- (-)	per person has changed between 2014 and 2018.				
	At the 10% significance level, the null hypothesis is rejected.				
	With reference to the 10% significance level, explain why it is not necessarily true that there has been a change.				
	[2 marks]				



Turn over for the next question DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED



Turn over ▶

Е	Elizabeth's Bakery makes brownies.
lt d	is known that the mass, $\boldsymbol{X}$ grams, of a brownie may be modelled by a normal listribution.
1	0% of the brownies have a mass less than 30 grams.
8	0% of the brownies have a mass greater than 32.5 grams.
F	Find the mean and standard deviation of $X$ . $ \begin{tabular}{ll} \hline \end{tabular}  \begin{tabular}{ll} \hline \end{tabular}$
_	
_	
_	
_	
_	
_	
_	
_	
_	
_	
_	
_	



Do	not	writ
οι	ıtside	e the

17 (b) (i)	Find $P(X \neq 35)$ [1 mark]
17 (b) (ii)	Find $P(X < 35)$ [2 marks]
17 (c)	Brownies are baked in batches of 13.
	Calculate the probability that, in a batch of brownies, no more than 3 brownies are less than 35 grams.
	You may assume that the masses of brownies are independent of each other.  [2 marks]
	END OF QUESTIONS



There are no questions printed on this page

Do not write outside the

# DO NOT WRITE ON THIS PAGE ANSWER IN THE SPACES PROVIDED

#### Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.



