



GCE

Chemistry B

H433/02: Scientific literacy in chemistry

Advanced GCE

Mark Scheme for November 2020

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.




© OCR 2020

Annotations

Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
	Benefit of doubt given
	Contradiction
	Rounding error
	Error in number of significant figures
	Error carried forward
	Level 1
	Level 2
	Level 3
	Benefit of doubt not given
	Noted but no credit given
	Ignore


Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

Annotation	Meaning
DO NOT ALLOW	Answers which are not worthy of credit
IGNORE	Statements which are irrelevant
ALLOW	Answers that can be accepted
()	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
ECF	Error carried forward
AW	Alternative wording
ORA	Or reverse argument


Question			Answer	Mark	AO element	Guidance
1	a		 <p>✓</p>	1	1.1	ALLOW correct C–N bond circled
1	b	i	 <p>Correct 3D structure around chiral centre ✓ correct mirror image ✓</p>	2	2.1 x 2	ALLOW any object and correct mirror image scores 1 mark ALLOW dotted wedges instead of dotted lines. ALLOW wedges for bonds in front of the plane of the paper ALLOW two ordinary lines as long as they are not 180° to each other.
1	b	ii	 <p>✓</p>	1	1.1	DO NOT ALLOW NH ³⁽⁺⁾ IGNORE extra correct details on skeletal structure
1	b	iii	<p>H₂NCH(COOH)CH₂COOH also formed ✓ also CH₃OH ✓ hydrolysis of ester ✓</p>	3	3.2 x 3	ALLOW any unambiguous structures ALLOW NH ₃ ⁺ in formula of amino acid
1	c			1	1.1	ALLOW circles round NH ₂ , NH and OH rather than the atoms separately ALL indicated atoms MUST be circled to score the mark

Question			Answer	Mark	AO element	Guidance
1	d		Instantaneous (dipole)-induced dipole/id-id ✓	1	2.5	ALLOW Van der Waals or London
1	e	i	place in solvent and allow solvent to rise up paper ✓ solvent below level of spot ✓ (dry and) use ninhydrin/locating agent/UV (light) ✓ more than one spot indicates hydrolysis ✓	4	1.2 3.3 1.2 3.4	ALLOW labelled diagrams to score MP1, 2 & 4 IGNORE use of water as solvent ALLOW matching spots from individual amino acids
1	e	ii	enzyme would be denatured /tertiary structure broken down so no reaction (AW)✓	1	3.1	
1	e	iii	No difference to rate (AW)✓ Aspartame/substrate zero order (when in excess) ✓	2	3.1 x 2	ALLOW initially rate increases AND then becomes constant(AW) ALLOW all active sites occupied(AW) when aspartame in excess/at high concentration'


Question			Answer	Mark	AO element	Guidance
2	a	i	salt bridge ✓ filter paper and potassium nitrate (solution) ✓	2	3.4 x 2	ALLOW ion bridge ALLOW any Gr I /II nitrate solution
2	a	ii	298 K/ 25°C ✓ concentrations of solutions 1 mol dm ⁻³ ✓	2	1.2 x 2	
2	b	i	silver <u>ions</u> / Ag ⁺ ✓	1	1.2	
2	b	ii	From Cu to Ag AND Cu electrode potential is more negative/less positive ✓	1	2.3	ALLOW Ag electrode is more positive/less negative ALLOW Cu is oxidised/loses electrons ALLOW Ag ⁺ ions are reduced/accept electrons
2	c		FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 6.8 x 10⁻⁷ (mol dm⁻³) award 4 marks $E^{\ominus} = 0.46$ ✓ Rearrange equation: $0.06 \log[\text{Ag}^+] = E - E^{\ominus}$ ✓ $\log [\text{Ag}^+] = -6.17$ ✓ $[\text{Ag}^+] = 6.8 \times 10^{-7} \text{ (mol dm}^{-3}\text{)}$ ✓	4	2.8 x 4	ALLOW 1 or more sf (0.06 in equation) ALLOW ecf throughout
2	d	(i)	Equation showing that IO ₃ ⁻ reacts with Cu ✓ $2\text{IO}_3^- + 12\text{H}^+ + 5\text{Cu} \rightarrow \text{I}_2 + 6\text{H}_2\text{O} + 5\text{Cu}^{2+}$ ✓	2	2.8 x 2	Mark separately Second mark can be scored for reverse equation. ALLOW equilibrium sign IGNORE state symbols
2	d	(ii)		2	3.3 x 2	




Question			Answer	Mark	AO element	Guidance
			Pt/graphite electrode ✓ IO ₃ ⁻ and I ₂ and H ⁺ in solution ✓			
2	d	(iii)	FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 0.4 (mol dm⁻³) award 3 marks [I ₂] = 0.04 ✓ √0.04 = 0.2 ✓ [IO ₃ ⁻] = 0.4 (mol dm ⁻³) ✓	3	2.8 x 3	0.28.... (omission of factors of 2) scores 1 mark 0.56 and 0.2 score 2 marks ALLOW ecf from any statement that begins "[I ₂] ="

Question		Answer	Mark	AO element	Guidance	
3	a	 <p>ten electrons around P atom / eight electrons around P atom if clear that lone pair from P are used to form a dative covalent bond to one O atom ✓</p> <p>rest correct ✓</p>	2	2.5 x 2	IGNORE minus sign ALLOW 'extra' electron on single bonded O to be the same symbol or different from the rest of the electrons on that O. Must be clear that only five electrons are from P atom.	
3	b	<p>mol P = $52 \times 2/142 = 0.73$ AND mol K = $34 \times 2/94.2 = 0.72$ ✓</p> <p>Ratio is about 1:1 as in KH_2PO_4 ✓</p>	2	2.6 x 2	ALLOW $52/142 = 0.365$ AND $34/94.2 = 0.36$ for MP1	
3	c	i	HPO_4^{2-} AND proton acceptor ✓	1	1.1	
3	c	ii	7.2..... ✓	1	1.2	
3	c	iii	<p>FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 5.1.... award 2 marks</p> <p>$[\text{H}^+] (= \sqrt{(6.2 \times 10^{-8} \times 1.0 \times 10^{-3})}) = 7.87 \times 10^{-6} (\text{mol dm}^{-3})$ ✓</p> <p>pH = 5.1... ✓</p>	2	2.2 x 2	ALLOW ecf provided ' $[\text{H}^+] =$ ' or ' $\text{H}^+ =$ ' is shown AND value is $> 10^{-7}$ and $< 10^{-4}$
3	d	<p>$\text{NH}_4^+ \rightleftharpoons \text{NH}_3 + \text{H}^+$ OR $\text{NH}_4^+ + \text{H}_2\text{O} \rightleftharpoons \text{NH}_3 + \text{H}_3\text{O}^+$ ✓</p> <p>Acid/ H^+/ H_3O^+ (from KH_2PO_4) moves equilibrium (position) to left ✓</p>	2	2.7 x 2	ALLOW equilibrium the other way round (and the corresponding 'right' in MP2)	

Question		Answer	Mark	AO element	Guidance	
3	e	<p>FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 69 (g) award 4 marks</p> <p>Value of $[H^+]$: 3.16×10^{-7} ✓</p> <p>Rearrangement of eqn: $[H_2PO_4^-] = [H^+] [HPO_4^{2-}] / K_a$ ✓</p> <p>$[H_2PO_4^-] = 3.16 \times 10^{-7} \times 0.1 / 6.2 \times 10^{-8}$ OR 0.51 ✓</p> <p>mass $KH_2PO_4 = (0.51 \times 136) = 69$ (g) ✓</p>	4	2.8 x 4	<p>ALLOW 2 or more sf</p> <p>ALLOW ecf throughout</p>	
3	f	i	Equilibrium will move to left (AW) ✓	2	3.1 x 2	
			$[H^+]$ only restored/unchanged (AW) if $[H_2PO_4^-]$ is large ✓			
3	f	ii	(Student is correct (AW) AND) HCl strong (acid) so all reacts/dissociates/ionises ✓	1	3.1	
	f	iii	<p>FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 0.7 award 2 marks</p> <p>$[H^+] (= 0.05 \times 0.01 / 1000) = 5 \times 10^{-7}$ ✓</p> <p>pH change = $7 - 6.3 = 0.7$ ✓</p>	2	2.8 x 2	ALLOW ecf

Question			Answer	Mark	AO element	Guidance
4	a	i	(C=C) decolorises bromine water ✓ (COOH/acid) will fizz with carbonate/appropriate colour with (named) indicator ✓	2	1.2 x 2	ALLOW gas produced turns limewater cloudy(AW) if carbonate test used
4	a	ii	FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 5 (cm³) award 3 marks conc maleic acid = (2.32/116) OR 0.02 (mol dm ⁻³) ✓ amt NaOH = 2 x 250 x 0.02/1000 OR 0.01 (mol) ✓ vol NaOH = 1000 x 0.01/2 = 5 ✓	3	2.4 x 3	2.5 scores 2 marks ALLOW ecf Allow wrong answers (after ecf) to 2 or more sig figs.
4	b	i	CHO ✓	1	1.2	ALLOW elements in any order
	b	ii	Z ✓ butenedioic acid ✓	2	1.1 x 2	ALLOW 'butendioic acid'
	c		Description of groups being spatially different across C=C ✓ Lack of free rotation of C=C /groups in a fixed position ✓	2	1.2 x 2	ALLOW they are stereoisomers.
	d		Loss of COOH/CO ₂ H OR C ₃ H ₃ O ₂ (⁺)/CHCHCOOH ✓	1	2.6	
	e	i	 one for each arrow ✓✓	2	2.5 x 2	One arrow must start) on the double bond and end pointing to H ⁺ (or to the bond being formed) The other arrow must start and end on the bonds shown. Other arrows are CON
	e	ii	(Electrophilic) Elimination	1	1.1	

Question		Answer	Mark	AO element	Guidance
e	iii	No, either of the Hs on the left C (of carbocation) could leave ✓ some $^2\text{H/D}$ would be incorporated ✓	2	3.2 x 2	
f			1	2.1	
g	i	Atom economies $(98/170) = 58\%$ AND $(98/222) = 44\%$ ✓ Identifies butane/reaction 1 should be used because it has the larger atom economy/produces less waste ✓	2	2.6 3.1	ALLOW 0.58 AND 0.44
g	ii	Benzene toxic/ more expensive/butane more available from cracking /doesn't produce CO_2 (ORA for reaction 2) ✓	1	3.1	ALLOW ecf from (g)(i) eg butane more flammable
g	iii	FIRST CHECK THE ANSWER ON ANSWER LINE If answer = 61 (kg) award 2 marks amt butane = $15000/24 = 625 \text{ mol}$ ✓ mass maleic anh = $625 \times 98/1000 = 61(\text{kg})$ (2sf) ✓	2	2.8 x 2	61.3 or 61.2 score 1 Correct use of gas equation can score both marks

Question		Answer	Mark	AO element	Guidance
	h*	<p><i>Refer to marking instructions on page 5 of mark scheme for guidance on marking this question.</i></p> <p>Level 3 (5 – 6 marks) Correct structure/identity deduced and detailed evidence related to the structure is provided from each spectrum.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p>Level 2 (3 – 4 marks) Correct structure/identity deduced with detailed evidence from at least one spectrum. OR Structure/identity attempted with some correct evidence from a minimum of two spectra.</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p>Level 1 (1 – 2 marks) Correct structure/identity deduced with no relevant evidence. OR Structure/identity not given or incorrect, but some correct evidence from at least one spectrum.</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p>0 marks No response or nothing of value</p>	6	3.1 x 4 3.2 x 2	<p>Indicative scientific points may include:</p> <p>AO3.1 Analysis: IR:</p> <ul style="list-style-type: none"> • C=O at 1700 • Broad OH 2500-3300 • COOH <p>HNMR:</p> <ul style="list-style-type: none"> • Three H environments • 1.2: Hs on C connected to C-H • 2.3: H on C connected to Cs with many Hs • 11.8 H in COOH <p>CNMR:</p> <ul style="list-style-type: none"> • Three C environments • two Cs in same environment • one C=O (at 183ish) • two C-C <p>AO3.2 Evaluation: Compound A identified as 2-methylpropanoic acid Structure $\text{CH}_3\text{CH}(\text{CH}_3)\text{COOH}$ presented in any unambiguous way</p>

Question			Answer	Mark	AO element	Guidance
5	a	i	Two Cls/chlorines/two Cl-/ chloride ligands are each -1 ✓	1	1.1	
5	a	ii	5d ⁸ ✓	1	1.1	
5	a	iii	Yes, it has an ion with an incomplete d subshell (in a compound) ✓	1	1.1	
5	b		<p>correct identification ✓</p> <p>full structural (with O⁻ for first two) ✓</p>	2	1.1 x 2	
5	c	i	guanine ✓	1	1.1	
5	c	ii	hydrogen (bond) ✓	1	2.1	
5	c	iii	(dative) covalent/ co-ordinate (bond) ✓ (lone) pair on nitrogen donated to Pt ✓	2	2.1 x 2	
5	d	i	octahedral ✓	1	2.1	
		ii	6 ✓	1	2.1	

Question			Answer	Mark	AO element	Guidance
5	d	iii	<p>Satraplatin also has ethanoate/ester/C=O groups/(4) O atoms (in addition to NH₃ groups that cisplatin also has) ✓</p> <p>(These form more) H-bonds with water(ORA for cisplatin) ✓</p> <p>and (more than) compensate for 'insoluble' (AW) ring OR H-bonds broken in water ✓</p>	3	3.1 x 3	Oxygen MOLECULES is CON
5	e*		<p><i>Refer to marking instructions on page 5 of mark scheme for guidance on marking this question.</i></p> <p>Level 3 (5 – 6 marks) Considers disadvantages of cisplatin, with most examples of both toxicity and cell resistance. AND Identifies most examples of how new drugs attempt to overcome the disadvantages of cisplatin.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p>Level 2 (3 – 4 marks) Considers disadvantages of cisplatin, with some examples of both toxicity and cell resistance. OR Gives a detailed treatment of either toxicity OR cell resistance</p> <p>AND Attempts to provide examples of the ways new drugs attempt to overcome the disadvantages of cisplatin.</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p>	6	3.1 x 3 3.2x 3	<p>Indicative scientific points include:</p> <p>Disadvantages:</p> <p>toxicity</p> <ul style="list-style-type: none"> damaging to the kidneys/ gastrointestinal tract / nervous system. targets all cells difficulty of administration <p>cell resistance</p> <ul style="list-style-type: none"> influx slower/method of entry slowed/prevented efflux faster/method of exit accelerated removed before it can act on DNA improved cell repair mechanism cancer cells evolve improved methods of DNA repair <p>New Drugs attempting to overcome disadvantages</p> <ul style="list-style-type: none"> carboplatin less toxic satraplatin/picoplatin different structure and overcame resistance formulae of some of above bulky group in picoplatin helps overcome resistance satraplatin soluble/ administered orally

Question	Answer	Mark	AO element	Guidance
	<p>Level 1 (1 – 2 marks) Considers disadvantages of cisplatin, with examples of toxicity OR cell resistance. OR Attempts to provide examples of the ways new drugs attempt to overcome the disadvantages of cisplatin</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant and correct.</i></p> <p>0 marks No response or nothing of value</p>			

OCR (Oxford Cambridge and RSA Examinations)
The Triangle Building
Shaftesbury Road
Cambridge
CB2 8EA

OCR Customer Contact Centre

Education and Learning

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored