



Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

---

Forename(s)

---

Candidate signature

---

I declare this is my own work.

# A-level FURTHER MATHEMATICS

## Paper 3 Statistics

Thursday 11 June 2020

Afternoon

Time allowed: 2 hours

### Materials

- You must have the AQA formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)
- You must ensure you have the other optional Question Paper/Answer Book for which you are entered (**either** Discrete **or** Mechanics). You will have 2 hours to complete **both** papers.

### Instructions

- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 50.

### Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
<b>TOTAL</b>	



J U N 2 0 7 3 6 7 3 S 0 1

PB/Jun20/E8

**7367/3S**

Answer **all** questions in the spaces provided.

- 1** The continuous random variable  $X$  has probability density function

$$f(x) = \begin{cases} \frac{1}{5} & 1 \leq x \leq 6 \\ 0 & \text{otherwise} \end{cases}$$

Find  $P(X \geq 3)$

Circle your answer.

[1 mark]

$\frac{1}{5}$

$\frac{2}{5}$

$\frac{3}{5}$

$\frac{4}{5}$

- 2** Jamie is conducting a hypothesis test on a random variable which has a normal distribution with standard deviation 1

The hypotheses are

$$H_0 : \mu = 5$$

$$H_1 : \mu > 5$$

He takes a random sample of size 4

The mean of his sample is 6

He uses a 5% level of significance.

Before Jamie conducted the test, what was the probability that he would make a Type I error?

Circle your answer.

[1 mark]

0.0228

0.0456

0.0500

0.1587



**3** The mass of male giraffes is assumed to have a normal distribution.

Duncan takes a random sample of 600 male giraffes.

The mean mass of the sample is 1196 kilograms.

The standard deviation of the sample is 98 kilograms.

**3 (a)** Construct a 94% confidence interval for the mean mass of male giraffes, giving your values to one decimal place.

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

**3 (b)** Explain whether or not your answer to part (a) would change if a sample of size 5 was taken with the same mean and standard deviation.

**[1 mark]**

---

---

---

**Turn over for the next question**

**Turn over ►**





**4 (b)** A spinning toy can land on one of four values: 2, 4, 6 or 8

Using a discrete uniform distribution, find the probability that the next value the toy lands on is greater than 2

**[1 mark]**

---

---

---

**4 (c)** State an assumption that is required for the discrete uniform distribution used in part **(b)** to be valid.

**[1 mark]**

---

---

---

**Turn over for the next question**

**Turn over ►**



**5** Emily claims that the average number of runners per minute passing a shop during a long distance run is 8

Emily conducts a hypothesis test to investigate her claim.

**5 (a)** State the hypotheses for Emily's test.

**[1 mark]**

---

---

---

---

**5 (b)** Emily counts the number of runners,  $X$ , passing the shop in a randomly chosen minute.

The critical region for Emily's test is  $X \leq 2$  or  $X \geq 14$

During a randomly chosen minute, Emily counts 3 runners passing the shop.

Determine the outcome of Emily's hypothesis test.

**[3 marks]**

---

---

---

---

---

---

---

---

---

---



**5 (c)** The actual average number of runners per minute passing the shop is 7

Find the power of Emily's hypothesis test, giving your answer to three significant figures.

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

**Turn over for the next question**

**Turn over ►**





**6 (b)** Find the probability density function of the random variable  $X$ .

**[2 marks]**

---

---

---

---

---

---

---

**6 (c)** After investigation, the distances between successive breaks in water pipes are found to have a standard deviation of 5 metres.

Explain whether or not the use of an exponential model in parts **(a)** and **(b)** is appropriate.

**[2 marks]**

---

---

---

---

---

---

---

**Turn over for the next question**

**Turn over ►**






**7 (b)** For the test carried out in part (a), state in context the meaning of a Type II error. **[1 mark]**

---

---

---

**7 (c)** The distribution of rainfall per day in February in the town over 10 years is shown in the histogram.



Explain whether or not the assumption that rainfall per day in February has a normal distribution is appropriate.

**[1 mark]**

---

---

---

**Turn over for the next question**

**Turn over ►**



**8** Ray is conducting a hypothesis test with the hypotheses

$H_0$  : There is no association between time of day and number of snacks eaten

$H_1$  : There is an association between time of day and number of snacks eaten

He calculates **expected** frequencies correct to two decimal places, which are given in the following table.

		Number of snacks eaten		
		0	1	2 or more
Time of Day	Day	23.68	21.05	5.26
	Night	21.32	18.95	4.74

Ray calculates his test statistic using  $\sum \frac{(O - E)^2}{E}$

**8 (a)** State, with a reason, the error Ray has made and describe any changes Ray will need to make to his test.

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



**8 (b)** Having made the necessary corrections as described in part (a), the correct value of the test statistic is 8.74

Complete Ray's hypothesis test using a 1% level of significance.

**[3 marks]**

---

---

---

---

---

---

---

---

---

---

**Turn over for the next question**

**Turn over ►**





*Do not write  
outside the  
box*

---

---

---

---

---

---

---

---

---

---

**END OF QUESTIONS**



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**









