



**GCE**

**Chemistry A**

**H432/02: Synthesis and analytical techniques**

Advanced GCE

**Mark Scheme for Autumn 2021**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.

© OCR 2021

## 1. Annotations





Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
	Benefit of doubt given
	Contradiction
	Rounding error
	Error in number of significant figures
	Error carried forward
	Level 1
	Level 2
	Level 3
	Benefit of doubt not given
	Noted but no credit given
	Ignore

2. Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).



<b>Annotation</b>	<b>Meaning</b>
<b>DO NOT ALLOW</b>	Answers which are not worthy of credit
<b>IGNORE</b>	Statements which are irrelevant
<b>ALLOW</b>	Answers that can be accepted
( )	Words which are not essential to gain credit
—	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument


Question	Answer	Marks	AO element	Guidance
1	C	1	AO2.1	<b>ALLOW</b> 4 (This is the number of structural isomers)
2	B	1	AO1.2	
3	C	1	AO2.2	
4	C	1	AO2.6	
5	D	1	AO2.1	
6	B	1	AO1.2	
7	A	1	AO1.2	
8	C	1	AO2.1	
9	C	1	AO1.2	
10	A	1	AO2.1	
11	D	1	AO2.5	
12	B	1	AO2.1	
13	B	1	AO2.1	
14	C	1	AO1.1	
15	A	1	AO1.2	
	<b>Total</b>	<b>15</b>		





Question		Answer	Marks	AO element	Guidance
16	(a) (i)	<p><math>\sigma</math>-bond: Overlap of <b>orbitals</b> between (bonding) atoms ✓</p> <p><math>\pi</math>-bond: <b>Sideways</b> overlap of (adjacent) <b>p-orbitals</b> ✓</p>	2	AO1.1 ×2	<p><b>ALLOW</b> labelled diagrams</p> <p><b>IGNORE</b> the type of orbital for <math>\sigma</math>-bond</p> <p><b>DO NOT ALLOW</b> pi-orbital</p>
	(ii)	<p><math>\sigma</math>-bonds: 9 ✓</p> <p><math>\pi</math>-bonds: 2 ✓</p>	2	AO1.2 ×2	
	(b) (i)	<p>Curly arrow from C=C bond to H of H-Br ✓ <b>DO NOT ALLOW</b> partial charge on C=C</p> <p>Correct dipole shown on H-Br <b>AND</b> curly arrow showing breaking of H-Br bond ✓</p>	4	AO1.2 ×2  AO2.5 ×2	<p><b>NOTE:</b> curly arrows can be straight, snake like, etc. but <b>NOT</b> double headed or half headed arrows</p> <p><b>1st curly arrow must</b></p> <ul style="list-style-type: none"> <li>go to the H atom of H-Br</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>start from, <b>OR</b> be traced back to <b>any point across width</b> of C=C</li> </ul> <p><b>2nd curly arrow must</b></p> <ul style="list-style-type: none"> <li>start from, <b>OR</b> be traced back to <b>any part of</b> <math>\delta^+H-Br\delta^-</math> bond</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>go to <math>Br\delta^-</math></li> </ul>





Question	Answer	Marks	AO element	Guidance
	<p>Correct carbocation  <b>AND</b> curly arrow from Br<sup>-</sup> to C<sup>+</sup> of carbocation ✓  <b>DO NOT ALLOW</b> δ<sup>+</sup> on C of carbocation</p>  <p>Correct product ✓</p> 			<p><b>3rd curly arrow must</b></p> <ul style="list-style-type: none"> <li>go to the C<sup>+</sup> of carbocation  <b>AND</b></li> <li>start from, <b>OR</b> be traced back to any point across width of lone pair on :Br<sup>-</sup></li> <li><b>OR</b> start from – charge of Br<sup>-</sup> ion</li> </ul>  <p>(Lone pair <b>NOT</b> needed if curly arrow shown from – charge of Br<sup>-</sup> ion)</p> <p><b>ALLOW ECF</b> for product from incorrect carbocation, i.e.</p>  <p><b>IF</b> Br<sub>2</sub> is used instead of HBr contact your Team Leader</p>


Question		Answer	Marks	AO element	Guidance
	(ii)	(major product forms from) most/more stable intermediate/carbocation ✓  (major product forms from a) secondary carbocation <b>OR</b> carbocation bonded to more C atoms / more alkyl groups <b>OR</b> carbocation bonded to fewer H atoms ✓	2	AO1.1  AO1.2	For carbocation, <b>ALLOW</b> carbonium ion or cation  <b>IGNORE</b> descriptions of the major/minor product in terms of Markownikoff's rule e.g. H atom joins to C with most H  <b>IGNORE</b> references to stability of the product ----- <b>ALLOW ORA</b> , i.e. (minor product forms from) least/less stable intermediate/carbocation ✓  (minor product forms from a) primary carbocation <b>OR</b> carbocation bonded to less C atoms / less alkyl groups <b>OR</b> carbocation bonded to more H atoms ✓
	(iii)	3 ✓	1	AO1.2	
(c)	(i)	Same structural formula <b>AND</b> Different arrangement (of atoms) in space <b>OR</b> different spatial arrangement (of atoms) ✓	1	AO1.1	<b>ALLOW</b> structure/displayed/skeletal formula  <b>DO NOT ALLOW</b> same empirical formula <b>OR</b> same general formula  <b>IGNORE</b> same molecular formula  Reference to <i>E/Z</i> isomerism or optical isomerism is <b>not</b> sufficient
	(ii)	Student is <b>not</b> correct <b>AND</b> 2 groups on one carbon atom (of C=C) are the same <b>OR</b> C–C bond can rotate ✓	1	AO3.1	<b>DO NOT ALLOW</b> one side of C=C




Question		Answer	Marks	AO element	Guidance
(d)	(i)	 <p>1 mark for each curly arrow ✓✓</p>	2	AO2.5 x2	<p><b>IGNORE</b> any dipoles shown</p> <p><b>NOTE:</b> curly arrows can be straight, snake-like, etc. but <b>NOT</b> half headed or double headed arrows</p> <p><b>Curly arrow</b> from C=C bond must start from, <b>OR</b> be traced back to,</p> <p><b>Lower left:</b> any part of C=C bond and go to C-C</p> <p><b>Upper left:</b> any part of C=C bond and go to gap between C=C and C=C</p>
	(ii)		2	AO3.2 x2	
<b>Total</b>			<b>17</b>		

Question		Answer	Marks	AO element	Guidance
17	(a)	<p><b>Formation of Cl•</b>  <math>\text{CClF}_3 \rightarrow \text{CF}_3\cdot + \text{Cl}\cdot \checkmark</math></p> <p><b>Breakdown of O<sub>3</sub></b>  <math>\text{Cl}\cdot + \text{O}_3 \rightarrow \cdot\text{ClO} + \text{O}_2 \checkmark</math></p> <p><math>\cdot\text{ClO} + \text{O} \rightarrow \text{Cl}\cdot + \text{O}_2 \checkmark</math></p>	3	AO2.5  AO1.1 ×2	<p><b>IGNORE</b> dots for formation Cl•, i.e. <b>ALLOW</b> <math>\text{CClF}_3 \rightarrow \text{CF}_3 + \text{Cl}</math></p> <p><b>DO NOT ALLOW</b> ECF Dots <b>required</b> in this equation</p> <p><b>IGNORE</b> <math>\text{O} + \text{O}_3 \rightarrow 2\text{O}_2</math></p> <p><b>ALLOW 1 mark</b> if both equations are correct by atom but dot(s) missing or incorrect</p>
	(b) (i)		1	AO2.5	<p><b>ALLOW</b> any combination of skeletal <b>OR</b> structural <b>OR</b> displayed formula as long as unambiguous</p> <p>'End bonds' <b>MUST</b> be shown</p> <p><b>DO NOT ALLOW</b> more than 1 repeat unit</p> <p><b>IGNORE</b> brackets</p> <p><b>IGNORE</b> <i>n</i></p>
	(ii)	<p>More points of contact / more surface interaction (between molecules)  <b>AND</b>            Stronger/more dipole(-dipole) interactions <math>\checkmark</math></p> <p>More energy needed to break the intermolecular forces <math>\checkmark</math></p>	2	AO2.1 ×2	<p><b>Both answers need to be a comparison</b></p> <p><b>IGNORE</b> surface area  <b>ALLOW</b> more electrons</p> <p><b>ALLOW</b> induced/permanent dipole interactions  <b>ALLOW</b> London forces  <b>ALLOW</b> van der Waals' forces (as permanent dipole-dipole and induced dipole-dipole interactions are present within this polymer)  <b>IGNORE</b> IDID</p>

Question	Answer	Marks	AO element	Guidance
(c)	<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>✓</p> </div> <div style="text-align: center;">  <p>✓</p> </div> </div> <hr/> <div style="text-align: center;">  <p>Amide link: ✓</p> <p>1 repeat unit of correct polymer: ✓</p> </div>	4	<p>AO2.5 ×2</p> <p>AO1.2</p> <p>AO2.5</p>	<p><b>For polymer, DO NOT ALLOW &gt; 1 repeat unit</b></p> <p>'End bonds' <b>MUST</b> be shown (do not have to be dotted)</p> <p><b>ALLOW</b> -NH- at either end i.e.</p> <div style="text-align: center;">  </div> <p><b>IGNORE</b> brackets</p> <p><b>IGNORE</b> <i>n</i></p>
	<b>Total</b>	<b>10</b>		

Question			Answer	Marks	AO element	Guidance
18	(a)	(i)	Non-superimposable mirror images (about a chiral centre) ✓	1	AO1.1	
		(ii)	<p>Correct groups attached to chiral C of alanine seen once e.g.</p>  <p>Two <b>3D structures</b> of alanine that are mirror images <b>AND</b> correct connectivity in both i.e.</p> 	2	AO2.1 × 2	<p>Each structure must have four central bonds with <b>at least two wedges</b>. For bond into paper accept:</p>  <p><b>ALLOW</b> two 3D structures with 2 groups swapped e.g.</p>  <p><b>IF</b> CH<sub>3</sub> is shown as 'R' <b>ALLOW</b> 1 mark for two 3D structures with correct connectivity that are mirror images e.g.</p> 




Question	Answer	Marks	AO element	Guidance
(iii)	4 ✓	1	AO2.2	
(b)	<p>  </p> <p>ester C<sub>7</sub>H<sub>14</sub>O<sub>3</sub></p> <p>H<sup>+</sup>/H<sub>2</sub>O OR H<sup>+</sup>(aq) OR HCl(aq)</p> <p>C<sub>5</sub>H<sub>10</sub>O<sub>3</sub></p> <p>H<sub>2</sub>O</p> <p>acyl chloride C<sub>5</sub>H<sub>9</sub>O<sub>2</sub>Cl</p> <p>NaBr/Br<sup>-</sup> AND H<sub>2</sub>SO<sub>4</sub>/H<sup>+</sup></p> <p>haloalkane C<sub>5</sub>H<sub>9</sub>BrO<sub>2</sub></p> <p>NH<sub>3</sub> AND ethanol OR excess NH<sub>3</sub></p> <p>valine</p>	7	AO1.2 × 4  AO2.5 × 3	<p><b>ALLOW</b> any combination of skeletal <b>OR</b> structural <b>OR</b> displayed formula as long as unambiguous</p> <p><b>ALLOW</b> names of reagents</p> <p><b>DO NOT ALLOW</b> OH<sup>-</sup> for HO<sup>-</sup> but <b>ALLOW ECF</b> for subsequent use in (b)</p> <p>For hydrolysis, <b>ALLOW</b> dilute acid <b>ALLOW</b> alkaline conditions followed by protonation of carboxylate i.e. NaOH(aq)/OH<sup>-</sup>(aq) <b>AND</b> H<sup>+</sup>(aq)/HCl(aq)</p> <p><b>ALLOW</b> HBr for NaBr/H<sub>2</sub>SO<sub>4</sub></p>





Question		Answer	Marks	AO element	Guidance
	(c) (i)	$C_{13}H_{18}O_2$ ✓	1	AO2.1	<b>ALLOW</b> C, H and O in any order
	(ii)	<p><b>FIRST CHECK ANSWER ON THE ANSWER LINE</b>  <b>If answer = <math>1.17 \times 10^{21}</math> award 3 marks</b></p> <p><math>M(\text{ibuprofen}) = 206</math> ✓</p> <p><math>n(\text{ibuprofen}) = \frac{400 \div 1000}{206} = 1.94 \times 10^{-3} \text{ (mol)}</math> ✓</p> <p>Number of molecules = <math>1.94 \times 10^{-3} \times 6.02 \times 10^{23}</math>  = <math>1.17 \times 10^{21}</math> to <b>3 SF</b> ✓</p>	3	AO2.2 × 3	<p><b>ALLOW ECF</b> from (c)(i)</p> <p>Calculator: <math>1.941747573 \times 10^{-3}</math></p> <p><b>ALLOW ECF</b> from <math>n(\text{ibuprofen})</math>  <b>3 SF essential</b></p>
(d)	(i)	 	2	AO3.2 × 2	<p><b>IGNORE</b> small slip in carbon chains</p> <p><b>ALLOW</b></p> 
	(ii)	More soluble in water ✓	1	AO3.1	<p><b>Answer must be a comparison</b>  <b>ALLOW</b> dissolve faster/quicker  <b>IGNORE</b> absorbed more quickly (given in question)</p>
<b>Total</b>			<b>18</b>		




Question			Answer	Marks	AO element	Guidance
19	(a)	(i)	3-methylbut-2-enal ✓	1	AO1.2	<b>IGNORE</b> lack of hyphens, or addition of commas
		(ii)	<p>Reaction scheme for question 19(a)(ii):</p> <p>Starting material: prenal (3-methylbut-2-enal)</p> <p>Pathway 1 (Left):</p> <ul style="list-style-type: none"> <li>Reduction: <math>\text{NaBH}_4</math> → 3-methylbut-2-en-1-ol ✓</li> <li>Oxidation: <math>\text{Cr}_2\text{O}_7^{2-}</math> AND <math>\text{H}^+</math> ✓ → 3-methylbut-2-enoic acid ✓</li> <li>Esterification: <math>(\text{CH}_3)_2\text{CHOH}</math> ✓ → 3-methylbut-2-enoic isopropyl ester</li> </ul> <p>Pathway 2 (Right):</p> <ul style="list-style-type: none"> <li>Hydrogenation: <math>\text{H}_2</math> AND <math>\text{Ni}</math> ✓ → 3-methylbutan-1-ol ✓</li> <li>Esterification: <math>\text{CH}_3\text{COOH}</math> ✓ → 3-methylbutyl acetate</li> </ul>	7	AO1.2 ×4  AO2.5 ×3	<p><b>ALLOW</b> any combination of skeletal <b>OR</b> structural <b>OR</b> displayed formula as long as unambiguous</p> <p><b>ALLOW</b> names of reagents and catalyst</p> <p>For oxidation, <b>ALLOW</b> <math>\text{K}_2\text{Cr}_2\text{O}_7</math> for <math>\text{Cr}_2\text{O}_7^{2-}</math> <b>ALLOW</b> <math>\text{H}_2\text{SO}_4</math> for <math>\text{H}^+</math></p> <p>For left hand side esterification <b>IGNORE</b> <math>\text{C}_3\text{H}_7\text{OH}</math></p> <p><b>IF</b> esterification is given instead of hydrogenation contact your Team Leader</p>



Question	Answer	Marks	AO element	Guidance
(b)*	<p><i>Refer to marking instructions on page 5 of mark scheme for guidance on marking this question.</i></p> <p><b>Level 3 (5-6 marks)</b>            Correct calculation of the mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl  <b>AND</b>            Planned synthesis to form the intermediate C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN followed by hydrolysis to form <b>A</b> with most of the reagents identified and equations are mostly correct.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p><b>Level 2 (3-4 marks)</b>            Correct calculation of the mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl  <b>AND</b>            Planned synthesis to form the intermediate C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN with most of the reagents identified and equation is mostly correct  <b>OR</b>            Calculation of the mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl is partly correct  <b>AND</b>            Planned synthesis includes formation of the intermediate C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN followed by hydrolysis to form <b>A</b> with some of the reagents identified  <b>OR</b>            Attempts to calculate mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl but makes little progress  <b>AND</b>            Planned synthesis includes formation of the intermediate C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN followed by hydrolysis to form <b>A</b> with most of the reagents identified and equations are mostly correct</p>	6	AO2.4 ×2  AO2.7 ×2  AO3.3 ×2	<p><b>Indicative scientific points may include:</b></p> <p><b>Calculation of mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl</b></p> <p><b>Using moles</b></p> <ul style="list-style-type: none"> <li><math>n(\mathbf{A}) = \frac{5.44}{136}</math>  <math>= 0.04(00)</math> (mol)</li> <li><math>n(\text{C}_6\text{H}_5\text{CH}_2\text{Cl}) = 0.0400 \times \frac{100}{25}</math>  <math>= 0.16(0)</math> (mol)</li> <li>Mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl = 126.5 × 0.16  <math>= 20.2(4)</math> g</li> </ul> <p><b>Using mass</b></p> <ul style="list-style-type: none"> <li>Theoretical mass of ester = <math>5.44 \times \frac{100}{25}</math>  <math>= 21.76</math> (g)</li> <li>Theoretical <math>n(\text{C}_6\text{H}_5\text{CH}_2\text{Cl}) = \frac{21.76}{136}</math>  <math>= 0.16(0)</math> (mol)</li> <li>Mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl = 126.5 × 0.160  <math>= 20.2(4)</math> g</li> </ul> <p><b>ALLOW</b> small slip/rounding errors such as errors in <i>M<sub>r</sub></i>, e.g. use of 137 instead of 136 for C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>COOH</p> <p>-----</p> <p><i>Examples of partly correct calculations</i></p> <p>Mass = 1.265 g from <math>0.0400 \times \frac{25}{100} \times 126.5</math>            (% yield inverted)</p> <p>Mass = 5.06 g from <math>0.0400 \times 126.5</math>            (% yield omitted)</p>



Question	Answer	Marks	AO element	Guidance
	<p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p><b>Level 1 (1-2 marks)</b>            Calculation of the mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl is partly correct  <b>OR</b>            Attempts to calculate mass of C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>Cl but makes little progress  <b>AND</b>            Planned synthesis includes formation of the intermediate C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN with the reagent identified  <b>OR</b>            Planned synthesis includes both steps with some of the reagents identified  <b>OR</b>            Attempts equations for both steps but these may contain errors  <b>OR</b>            Describes one step of the synthesis with reagent(s) and equation mostly correct</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b>            No response or no response worthy of credit.</p>			<p><b><u>Synthesis: reagents and conditions</u></b></p> <p><b>Stage 1:</b> Formation of intermediate, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>CN</p> <ul style="list-style-type: none"> <li>• Reagents: CN<sup>-</sup>(/ethanol)</li> <li>• Equation:  <math display="block">\text{C}_6\text{H}_5\text{CH}_2\text{Cl} + \text{CN}^- \rightarrow \text{C}_6\text{H}_5\text{CH}_2\text{CN} + \text{Cl}^-</math> <b>OR</b> <math display="block">\text{C}_6\text{H}_5\text{CH}_2\text{Cl} + \text{NaCN} \rightarrow \text{C}_6\text{H}_5\text{CH}_2\text{CN} + \text{NaCl}</math> <b>(OR use of KCN)</b></li> </ul> <p><b>Stage 2:</b> Formation of <b>A</b>, C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>COOH</p> <ul style="list-style-type: none"> <li>• Reagents: H<sup>+</sup>/H<sub>2</sub>O (<b>ALLOW</b> 'acid hydrolysis')</li> <li>• Equation:  <math display="block">\text{C}_6\text{H}_5\text{CH}_2\text{CN} + 2\text{H}_2\text{O} + \text{H}^+ \rightarrow \text{C}_6\text{H}_5\text{CH}_2\text{COOH} + \text{NH}_4^+</math> <b>OR</b> <math display="block">\text{C}_6\text{H}_5\text{CH}_2\text{CN} + 2\text{H}_2\text{O} + \text{HCl} \rightarrow \text{C}_6\text{H}_5\text{CH}_2\text{COOH} + \text{NH}_4\text{Cl}</math></li> </ul>
	<b>Total</b>	<b>18</b>		


Question			Answer	Marks	AO element	Guidance
20	(a)	(i)	<p><b>Stage 1</b></p>  <p>1 mark for each curly arrow as shown.</p>	6	AO1.1 AO1.2 AO2.5	<p><b>ANNOTATE WITH TICKS AND CROSSES</b></p> <p><b>NOTE:</b> curly arrows can be straight, snake-like, etc. but <b>NOT</b> double headed or half headed arrows</p> <p><b>Curly arrow</b> from OH<sup>-</sup> must</p> <ul style="list-style-type: none"> <li>go to the H of O-H</li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>start from, <b>OR</b> be traced back to <b>any point across width</b> of lone pair on O of OH<sup>-</sup></li> </ul>  <ul style="list-style-type: none"> <li><b>OR</b> start from - charge<sup>-</sup>OH ion</li> </ul>  <p><b>Curly arrow</b> from O-H bond must start from, <b>OR</b> be traced back to, <b>any part of</b> O-H bond and go to O</p> <p><b>IGNORE</b> dipoles on O-H bond</p> <p><b>IGNORE</b> Na<sup>+</sup></p>

Question	Answer	Marks	AO element	Guidance
	<p><b>Stage 2</b></p> <p>Curly arrow from <math>\pi</math>-ring to C in <math>\text{CO}_2</math>  <b>AND</b>            curly arrow from the C=O bond to O atom ✓</p>  <p>Correct intermediate ✓</p> <p>Curly arrow from C–H bond to reform <math>\pi</math>-ring  <b>AND</b> <math>\text{H}^+</math> formed ✓</p>		<p>AO2.5</p> <p>AO2.5</p> <p>AO1.2</p>	<p><b>1st curly arrow</b> must</p> <ul style="list-style-type: none"> <li>go to the C of <math>\text{CO}_2</math></li> </ul> <p><b>AND</b></p> <ul style="list-style-type: none"> <li>start from, <b>OR</b> close to <b>circle of benzene ring</b></li> </ul>  <p><b>2nd curly arrow</b> must start from, <b>OR</b> be traced back to, <b>any part of</b> C=O bond and go to O</p>  <p><b>ALLOW 2nd</b> curly arrow from C=O to any O in <math>\text{CO}_2</math></p> <p><b>DO NOT ALLOW</b> the following intermediate:</p>  <p><math>\pi</math>-ring must cover more than half of the benzene ring structure  <b>AND</b>            the correct orientation, <i>i.e.</i> gap towards C with <math>\text{CO}_2^-</math></p> <p><b>ALLOW</b> + sign anywhere inside the 'hexagon' of the intermediate.</p>

Question	Answer	Marks	AO element	Guidance
				<p><b>DO NOT ALLOW</b> mark for intermediate if phenolic O<sup>-</sup> is missing</p> <p><b>curly arrow</b> must start from, <b>OR</b> be traced back to, <b>any part of C-H bond</b> and go inside the 'hexagon'</p> 
(ii)	<p>OH<sup>-</sup>: base ✓</p> <p>CO<sub>2</sub>: electrophile <b>OR</b> electron pair acceptor ✓</p>	2	AO2.1 ×2	<p><b>ALLOW</b> alkali</p> <p><b>IGNORE</b> 'nucleophile', 'donates electron pair'</p> <p><b>IGNORE</b> lone pair acceptor (<i>No lone pair involved</i>)</p>
(iii)	 <p>One ester link in organic product ✓</p> <p>Correct structure of organic product ✓</p> <p>Correct equation <b>AND</b> balanced ✓</p>	3	AO3.1 AO3.2 AO2.6	

Question		Answer	Marks	AO element	Guidance
	(b) (i)	Dissolve in <b>hot</b> water/solvent ✓  <b>Minimum</b> amount of solvent ✓  Cool <b>AND</b> Filter <b>AND</b> (leave to) dry ✓ <i>All three needed</i>	3	AO3.3 ×3	<b>ALLOW</b> any solvent  <b>IGNORE</b> <ul style="list-style-type: none"> <li>Initial filtering</li> <li>hot filtration to remove insoluble impurities</li> </ul> <b>DO NOT ALLOW</b> adding of a drying agent (e.g. MgSO <sub>4</sub> )
	(ii)	C : H : N : O 31.44/12 : 1.31/1 : 18.34/14 : 48.91/16 <b>OR</b> 2.62 : 1.31 : 1.31 : 3.06 ✓  6:3:3:7 <b>OR</b> C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub> ✓  Molecular formula = C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub> <b>AND</b> use of <i>M</i> = 229.0 (directly linked to molecular formula) ✓  Any trisubstituted –NO <sub>2</sub> substituted phenol that is consistent with <i>M</i> = 229.0 ✓  <b>Evidence for substitution</b> 2,4,6 <b>OR</b> 3,4,5 substituted phenol <b>AND</b> 4 peaks/ C environments from <sup>13</sup> C NMR ✓  2,4,6 substituted phenol <b>AND</b> directing effects of –OH ✓	6	AO1.2 × 2  AO3.1  AO3.2  AO3.1 ×2	<b>ALLOW</b> alternative approach for empirical formula and evidence that 229 is equal to C <sub>6</sub> H <sub>3</sub> N <sub>3</sub> O <sub>7</sub>  <b>DO NOT ALLOW</b> ECF from the empirical formula with the wrong molar ratio    
<b>Total</b>			<b>20</b>		

Question	Answer	Marks	AO element	Guidance
21*	<p>Refer to marking instructions on page 5 of mark scheme for guidance on marking this question.</p> <p><b>Level 3 (5–6 marks)</b> Compounds <b>D, E AND F</b> correctly identified <b>AND</b> Most of the observations and NMR data analysed.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p><b>Level 2 (3–4 marks)</b> Most of compounds <b>D, E AND F</b> correctly identified <b>AND</b> Some of the observations and NMR data analysed.</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p><b>Level 1 (1–2 marks)</b> Most of compounds <b>D, E AND F</b> correctly identified <b>OR</b> Some of compounds <b>D, E AND F</b> correctly identified <b>AND</b> Analyses some of the observations or NMR data <b>OR</b> Analyses most of the observations from the test-tube tests. <b>OR</b> Analyses most of the NMR data. <b>OR</b> Analyses some of the observations and NMR data</p>	6	AO3.1 ×4  AO3.2 ×2	<p><b>Indicative scientific points may include:</b> <b><u>Observations from Test-tube tests</u></b></p> <p>2,4 DNP    <b>D</b> has no C=O                   <b>E</b> and <b>F</b> have C=O present</p> <p>H<sup>+</sup>/Cr<sub>2</sub>O<sub>7</sub><sup>2-</sup> <b>D</b> is primary <b>OR</b> secondary alcohol                   <b>E</b> and <b>F</b> are ketones                   <i>(negative test shows not aldehydes)</i></p> <p>Br<sub>2</sub>            <b>D, E</b> and <b>F</b> have no C=C/are saturated</p> <p><b><u><sup>13</sup>C NMR analysis</u></b></p> <p><b>D:</b></p> <ul style="list-style-type: none"> <li>• 3 carbon environments/types of C</li> <li>• δ = 24, 36 ppm                    <b>C–C</b></li> <li>• δ = 73 ppm,                        <b>C–O</b></li> </ul> <p><b><u><sup>1</sup>H NMR analysis</u></b></p> <p><b>E:</b></p> <ul style="list-style-type: none"> <li>• δ = 2.4 ppm, quartet            <b>CH<sub>3</sub>–CH<sub>2</sub>–C=O</b></li> <li>• δ = 1.1 ppm, triplet            <b>CH<sub>3</sub>–CH<sub>2</sub>–</b></li> </ul> <p><b>F:</b></p> <ul style="list-style-type: none"> <li>• δ = 2.6 ppm, heptet/multiplet <b>(CH<sub>3</sub>)<sub>2</sub>–CH–C=O</b></li> <li>• δ = 2.1 ppm, singlet,            <b>CH<sub>3</sub>–C=O</b></li> <li>• δ = 1.1 ppm, doublet            <b>CH<sub>3</sub>–CH–</b></li> </ul> <p><b><u>Structures</u></b> <b>ALLOW</b> any combination of skeletal <b>OR</b> structural <b>OR</b> displayed formula as long as unambiguous</p> <div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center;">  <p><b>D</b></p> </div> <div style="margin: 0 20px;"><b>OR</b></div> <div style="text-align: center;">  </div> </div>

Question			Answer	Marks	AO element	Guidance
			<p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>0 marks</b> No response or no response worthy of credit.</p>			 <p>E                      F</p>
<b>Total</b>				<b>6</b>		

**OCR (Oxford Cambridge and RSA Examinations)**  
**The Triangle Building**  
**Shaftesbury Road**  
**Cambridge**  
**CB2 8EA**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored