



---

# Level 2 Certificate

# FURTHER MATHEMATICS

# 8365/2

Paper 2 Calculator

---

Mark scheme

June 2021

---

Version: 1.0 Final



2 1 6 G 8 3 6 5 / 2 / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [aqa.org.uk](http://aqa.org.uk)

#### **Copyright information**

AQA retains the copyright on all its publications. However, registered schools/colleges for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2021 AQA and its licensors. All rights reserved.

## Glossary for Mark Schemes

GCSE examinations are marked in such a way as to award positive achievement wherever possible. Thus, for GCSE Mathematics papers, marks are awarded under various categories.

If a student uses a method which is not explicitly covered by the mark scheme the same principles of marking should be applied. Credit should be given to any valid methods. Examiners should seek advice from their senior examiner if in any doubt.

|                |  |
|----------------|--|
| <b>M</b>       | Method marks are awarded for a correct method which could lead to a correct answer.  |
| <b>M dep</b>   | A method mark dependent on a previous method mark being awarded.   |
| <b>A</b>       | Accuracy marks are awarded when following on from a correct method. It is not necessary to always see the method. This can be implied. |
| <b>B</b>       | Marks awarded independent of method.   |
| <b>B dep</b>   | A mark that can only be awarded if a previous independent mark has been awarded.   |
| <b>ft</b>      | Follow through marks. Marks awarded following a mistake in an earlier step.  |
| <b>SC</b>      | Special case. Marks awarded within the scheme for a common misinterpretation which has some mathematical worth.                        |
| <b>oe</b>      | Or equivalent. Accept answers that are equivalent.<br><br>eg accept 0.5 as well as $\frac{1}{2}$                                       |
| <b>[a, b]</b>  | Accept values between $a$ and $b$ inclusive.   |
| <b>3.14...</b> | Accept answers which begin 3.14 eg 3.14, 3.142, 3.1416   |

Examiners should consistently apply the following principles.

**Diagrams**

Diagrams that have working on them should be treated like normal responses. If a diagram has been written on but the correct response is within the answer space, the work within the answer space should be marked. Working on diagrams that contradicts work within the answer space is not to be considered as choice but as working, and is not, therefore, penalised.

**Responses which appear to come from incorrect methods**

Whenever there is doubt as to whether a candidate has used an incorrect method to obtain an answer, as a general principle, the benefit of doubt must be given to the candidate. In cases where there is no doubt that the answer has come from incorrect working then the candidate should be penalised.

**Questions which ask candidates to show working**

Instructions on marking will be given but usually marks are not awarded to candidates who show no working.

**Questions which do not ask candidates to show working**

As a general principle, a correct response is awarded full marks.

**Misread or miscopy**

Candidates often copy values from a question incorrectly. If the examiner thinks that the candidate has made a genuine misread, then only the accuracy marks (A or B marks), up to a maximum of 2 marks are penalised. The method marks can still be awarded.

**Further work**

Once the correct answer has been seen, further working may be ignored unless it goes on to contradict the correct answer.

**Choice**

When a choice of answers and/or methods is given, mark each attempt. If both methods are valid then M marks can be awarded but any incorrect answer or method would result in marks being lost.

**Work not replaced**

Erased or crossed out work that is still legible should be marked.

**Work replaced**

Erased or crossed out work that has been replaced is not awarded marks.

**Premature approximation**

Rounding off too early can lead to inaccuracy in the final answer. This should be penalised by 1 mark unless instructed otherwise.

**Continental notation**

Accept a comma used instead of a decimal point (for example, in measurements or currency), provided that it is clear to the examiner that the candidate intended it to be a decimal point.

| Q | Answer  | Mark  | Comments                                       |
|---|---|-------|--|
| 1 | $10x - 5$ or $44 - 4x$                                      | M1    | may be seen in a grid                          |
|   | $10x - 5 + 44 - 4x$ or $6x + 39$<br>or $3 \times (2x + 13)$ | M1dep | oe<br>all terms correct in a single expression |
|   | $3(2x + 13)$  | A1    |  |
|   | <b>Additional Guidance</b>                                  |       |  |
|   |   |       |  |

| Q    | Answer  | Mark | Comments   |
|------|---|------|--|
| 2(a) | $5m \times (1 - 0.4)$ or $5m \times 0.6$<br>or $3m$ | M1   | oe eg $5m - 0.4 \times 5m$ or $5m - 2m$<br>may be seen in an equation<br>eg $3m = m + 1$ |
|      | $\frac{1}{2}$ or 0.5                                | A1   |  |
|      | <b>Additional Guidance</b>                          |      |  |
|      | $2m$ only   |      | M0   |
|      | $2m = 1$  |      | M1   |

| Q    | Answer  | Mark | Comments          |
|------|---|------|-------------------|
| 2(b) | $2w - 10 = 18^3$<br>or<br>$2w - 10 = 5832$<br>or<br>$\frac{18^3 + 10}{2}$ | M1   | oe eg $2w = 5842$ |
|      | 2921  | A1   |                   |
|      | <b>Additional Guidance</b>  |      |                   |
|      |   |      |                   |

| Q | Answer   | Mark  | Comments   |
|---|--|-------|--|
| 3 | $12de \times 2d$ or $24d^2e$<br>or $\frac{1}{2} \times 8e^2 \times 9d$ or $36de^2$ | M1    | oe eg $24ed^2$ or $36e^2d$   |
|   | $24d = 36e$  | M1dep | oe equation with squared terms eliminated<br>eg $2d = 3e$ or $\frac{2d}{e} = 3$ or $\frac{e}{d} = \frac{12}{18}$ |
|   | $\frac{3}{2}$ or $1\frac{1}{2}$ or 1.5   | A1    |  |
|   | <b>Additional Guidance</b>   |       |  |
|   | Equivalent fraction to $\frac{3}{2}$ or $1\frac{1}{2}$ with no incorrect working   |       | M2A0   |

| Q  | Answer   | Mark  | Comments  |
|--|--|-------|---|
| 4  | $\pi \times 100 (\div 4)$ or $100\pi (\div 4)$<br>or $25\pi$<br>or<br>$\pi \times 36 (\div 4)$ or $36\pi (\div 4)$ or $9\pi$ | M1    | oe  |
|  | $\pi \times 100 \div 4 - \pi \times 36 \div 4$   | M1dep | oe eg $\frac{100\pi - 36\pi}{4}$ or $\frac{64\pi}{4}$ |
|  | $16\pi$  | A1    | SC2 $2176\pi$   |
|  | <b>Additional Guidance</b>   |       |   |
|  | Use of circumference instead of area throughout  |       | M0M0A0  |
|  | Allow substitution of $\pi = [3.14, 3.142]$ for M marks  |       |   |
|  | $16\pi$ in working with eg 50.3 on answer line   |       | M2A0  |
|  | SC2 is for using radii of 100 and 36   |       |   |
| Omission of $\pi$ in working must be recovered |  |       |   |

| Q                                    | Answer  | Mark  | Comments  |
|--------------------------------------|---|-------|---|
| 5                                    | $k - 1 = 15 - 6$<br>or<br>$\tan 45 = \frac{k - 1}{15 - 6}$  | M1    | oe equation<br>eg $k - 1 = 9$<br>or $\frac{k - 1}{9} = 1$                                       |
|                                      | $(k =) 15 - 6 + 1$<br>or<br>$(k =) (15 - 6) \tan 45 + 1$<br>or<br>$(k =) 10$  | M1dep | oe eg $(k =) 9 + 1$<br>may be seen on diagram   |
|                                      | (5.5, 8)  | A1    | oe eg $\left(5\frac{1}{2}, 8\right)$ or $\left(\frac{11}{2}, 8\right)$<br>SC1 answer (....., 8) |
|                                      | <b>Additional Guidance</b>  |       |   |
|                                      | First M1 can be scored using Pythagoras' theorem and cosine rule<br>eg $(PR^2 =) 9^2 + (15 - 6)^2$ or 162<br>and<br>$(PQ =) 6 - 1$ or 5<br>and<br>$(\text{angle } QPR =) 180 - 45$ or 135<br>and<br>$(k - 1)^2 + (15 - 1)^2 = \text{their } 162 + (\text{their } 5)^2$<br>$- 2 \times \sqrt{\text{their } 162} \times \text{their } 5 \times \cos \text{ their } 135$ |       | M1  |
| 10 but not seen or implied to be $k$ |   | M0M0  |   |

| Q                                       | Answer  | Mark  | Comments |
|---|---|-------|----------|
| <b>6</b>                                | <b>Alternative method 1</b>   |       |          |
|   | $y^2 = \frac{x+2w}{3}$  | M1    |          |
|   | $3y^2 - x = 2w$<br>or<br>$\frac{3y^2 - x}{2}$ or $\frac{3y^2}{2} - \frac{x}{2}$   | M1dep |          |
|   | $w = \frac{3y^2 - x}{2}$<br>or<br>$w = \frac{3y^2}{2} - \frac{x}{2}$  | A1    |          |
|   | <b>Alternative method 2</b>   |       |          |
|   | $y^2 = \frac{x}{3} + \frac{2w}{3}$  | M1    |          |
|   | $y^2 - \frac{x}{3} = \frac{2w}{3}$<br>or<br>$\frac{3}{2} \left( y^2 - \frac{x}{3} \right)$ or $\frac{3y^2}{2} - \frac{3x}{6}$ | M1dep |          |
|   | $w = \frac{3}{2} \left( y^2 - \frac{x}{3} \right)$<br>or<br>$w = \frac{3y^2}{2} - \frac{3x}{6}$                               | A1    |          |
|   | <b>Additional Guidance</b>  |       |          |
|   | Condone eg $w = \frac{3y^2 - x}{2}$ seen in working with $\frac{3y^2 - x}{2}$ on answer line                                  |       | M2A1     |
| $w = \frac{3}{2}y^2 - \frac{1}{2}x$ etc |   | M2A1  |          |

| Q    | Answer   | Mark | Comments               |
|------|--|------|------------------------|
| 7(a) | $a^{4m}$ or $a^{10m}$<br>or<br>$4m = 10m$            | M1   | oe eg $a^{4 \times m}$ |
|      | 0  | A1   |                        |
|      | <b>Additional Guidance</b>                           |      |                        |
|      | Allow $a$ to be replaced by any value greater than 1 |      |                        |

| Q    | Answer   | Mark | Comments   |
|------|--|------|--|
| 7(b) | $w^{13}x^7 \div w^3x^2$ or $w^{10}x^5$<br>or<br>$x^2y^5 = w^{10}x^7$ or $y^5 = \frac{w^{10}x^7}{x^2}$<br>or<br>$w^3y^5 = w^{13}x^5$ or $y^5 = \frac{w^{13}x^5}{w^3}$ | M1   | oe eg $\frac{w^{13}x^7}{w^3x^2}$<br><br>may be embedded eg $\sqrt[5]{w^{10}x^5}$ |
|      | $w^2x^{(1)}$   | A1   | oe eg $xw^2$   |
|      | <b>Additional Guidance</b>   |      |  |
|      | $y = w^{10}x^5$  |      | M1A0   |

| Q | Answer  | Mark | Comments  |  |
|---|---|------|---|--|
| 8 | -3 2 6 14<br>with no other solutions  | B4   | B3 three correct with at most one incorrect<br>B2 two correct with at most two incorrect<br>B1 one correct with at most three incorrect<br>SC2 -3 2 6 14 with no other values seen<br>SC1 Two or three of -3 2 6 14 with no other values seen |  |
|   | <b>Additional Guidance</b>  |      |   |  |
|   | Solutions may be in any order<br>eg1 -3 14 6 2<br>eg2 14 -3                           |      | B4<br>B2  |  |
|   | $x < -3$ $2 < x < 6$ $x > 14$   |      | SC2   |  |
|   | $2 \leq x \leq 6$   |      | SC1   |  |
|   | -3 2 6 14 seen in working with no other values and<br>answer line $-3 \leq x \leq 14$ |      | SC2   |  |

| Q    | Answer                     | Mark | Comments |  |
|------|----------------------------|------|----------|--|
| 9(a) | $\frac{b+a}{ab}$           | B1   |          |  |
|      | <b>Additional Guidance</b> |      |          |  |
|      |                            |      |          |  |

| Q    | Answer  | Mark | Comments  |  |
|------|---|------|---|--|
| 9(b) | $\frac{c^3}{6c+1}$                                    | B3   | B2 $c^3(6c-1)$ and $(6c+1)(6c-1)$<br>B1 $c^3(6c-1)$ or $(6c+1)(6c-1)$ |  |
|      | <b>Additional Guidance</b>                            |      |   |  |
|      | $\frac{c^3}{6c+1}$ followed by incorrect further work |      | B2  |  |

| Q         | Answer   | Mark  | Comments   |
|-----------|--|-------|--|
| <b>10</b> | <b>Alternative method 1</b>  |       |  |
|           | $\frac{4}{3} \times \pi \times \frac{27k^3}{8}$ or $\frac{9\pi k^3}{2}$                                  | M1    | oe   |
|           | $k^3 = 972 \times 2 \div 9$<br>or $k^3 = 216$<br>or $\sqrt[3]{216}$                                      | M1dep | oe eg $k^3 = 972\pi \times 2 \div 9\pi$<br>must have $k^3$ seen or implied |
|           | 6  | A1    |  |
|           | <b>Alternative method 2</b>  |       |  |
|           | $972\pi \div \frac{4}{3}\pi$ or 729<br>or $972\pi \div \frac{4}{3}\pi$ or 729<br>or $\sqrt[3]{729}$ or 9 | M1    | oe eg ( $r^3 =$ ) $972 \div \frac{4}{3}$                                   |
|           | $\frac{3k}{2} = \sqrt[3]{\text{their } 729}$<br>or $\frac{3k}{2} = 9$<br>or $9 \times 2 \div 3$          | M1dep | oe<br>must have $\frac{3k}{2}$ seen or implied                             |
|           | 6  | A1    |  |
|           | <b>Additional Guidance</b>   |       |  |
|           | Alt 1 $\frac{4}{3} \times \frac{27k^3}{8}$ is M0 unless recovered  |       |  |
|           | Alt 2 $972\pi \div \frac{4}{3}$ is M0 unless recovered   |       |  |
|           | Alt 1 $\frac{4}{3} \times \pi \times \left(\frac{3k}{2}\right)^3$ not subsequently simplified correctly  |       | M0   |

| Q   | Answer  | Mark | Comments  |  |
|---|---|------|---|--|
| 11  | 3 terms from<br>$20x^2 - 5xy^2 (+)12xy^2 - 3y^4$                        | M1   | may be seen in a grid   |  |
|   | $20x^2 - 5xy^2 + 12xy^2 - 3y^4$   | A1   | four correct terms in any order<br>may be seen in a grid<br>implied by correct answer |  |
|   | $20x^2 + 7xy^2 - 3y^4$  | A1   | terms may be in any order   |  |
|   | <b>Additional Guidance</b>  |      |   |  |
|   | Terms seen in a grid must have the correct signs                        |      |   |  |
|   | Terms must be fully processed eg do not allow $4x3y^2$ unless recovered |      |   |  |
|   | $xy^2$ may be $y^2x$ throughout   |      |   |  |
| $20x^2 + 7xy^2 - 3y^4$ followed by incorrect further work |   |      | M1A1A0  |  |

| Q   | Answer   | Mark  | Comments   |
|---|--|-------|--|
| 12  | 10   | B1    | y-coordinate of C<br>may be seen on the graph  |
|   | $(-)\frac{\text{their } 10}{5}$ or $(-)\ 2$                                | M1    | $\pm$ their gradient of L  |
|   | $(y =) -\frac{\text{their } 10}{5}x + \text{their } 10$                    | M1dep | oe eg $y - 0 = -\frac{\text{their } 10}{5}(x - 5)$<br>or $y - \text{their } 10 = -\frac{\text{their } 10}{5}(x - 0)$<br>must use a negative gradient |
|   | $-\frac{\text{their } 10}{5}x + \text{their } 10 = 3x + 2$<br>or $5x = 8$  | M1dep | oe   |
|   | 1.6  | A1ft  | oe eg $\frac{8}{5}$<br>ft B0M3   |
|   | <b>Additional Guidance</b>   |       |  |
|   | A1ft values must be exact or rounded to 1 decimal place or better          |       |  |
|   | Ignore any y-coordinate of B calculated after working out the x-coordinate |       |  |
|   | Assuming the lines are perpendicular can score a maximum of B1             |       |  |
|   | y-coordinate of C = 8  |       | B0   |
| gradient L = $-\frac{8}{5}$   |  | M1    |  |
| $y = -\frac{8}{5}x + 8$   |  | M1    |  |
| $-\frac{8}{5}x + 8 = 3x + 2$  |  | M1    |  |
| 1.3   |  | A1ft  |  |
| $\left( \text{Note that the exact value is } \frac{30}{23} \right)$ |  |       |  |

| Q  | Answer  | Mark  | Comments   |            |
|----|---|-------|--|------------|
| 13 | $3ax^2$ or $20x$  | M1    | oe eg $3 \times ax^{3-1}$ or $2 \times 10x^{2-1}$  |            |
|    | $3a \times 2^2 + 20 \times 2$ or $12a + 40$                               | M1    | ft substitution of $x = 2$ into their derivative<br>must have attempted differentiation<br>and have two terms with one involving $a$<br>may be seen in a denominator |            |
|    | their $(12a + 40) = -1 \div -\frac{1}{4}$<br>or<br>their $(12a + 40) = 4$ | M1dep | oe eg $-\frac{1}{\text{their } (12a + 40)} = -\frac{1}{4}$<br>dep on 2nd M1  |            |
|    | $-3$  | A1    |  |            |
|    | <b>Additional Guidance</b>  |       |  |            |
|    | Only substituting $x = 2$ into $y$  |       |  | Zero       |
|    | $ax^2 + 10x$<br>$4a + 20 = 4$   |       |  | M0<br>M1M1 |
|    | $3x^2 + 20x$<br>$12 + 20$   |       |  | M1<br>M0M0 |

| Q     | Answer   | Mark | Comments |          |
|-------|--|------|----------|----------|
| 14(a) | Reflection in the $x$ -axis<br>or<br>reflection in $y = 0$   | B1   |          |          |
|       | <b>Additional Guidance</b>   |      |          |          |
|       | Reflect(ed) in the $x$ -axis   |      |          | B1       |
|       | Do not allow if there is additional incorrect information<br>eg1 Reflection in the $x$ -axis about the origin<br>eg2 Reflection in the $x$ -axis and rotated |      |          | B0<br>B0 |
|       | Reflection   |      |          | B0       |

| Q                                 | Answer   | Mark | Comments  |                |
|-----------------------------------|--|------|---|----------------|
| 14(b)                             | Rotation through 180° centre the origin<br>or<br>enlargement scale factor –1 centre the origin   | B2   | B1 $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$<br>or<br>enlargement scale factor – 1<br>or<br>rotation through 180°<br>or<br>indication that <b>B</b> represents rotation through 270° (anticlockwise centre the origin)<br>or<br>indication that <b>B</b> represents rotation through 90° clockwise (centre the origin) |                |
|                                   | <b>Additional Guidance</b>   |      |   |                |
|                                   | For B2 ignore any reference to clockwise or anticlockwise rotation   |      |   |                |
|                                   | Condone omission of degrees symbol throughout<br>eg B is rotation through 270  |      |   | B1             |
|                                   | Mark intention<br>eg1 Rotate(d) 180 about <i>O</i><br>eg2 Enlarge(d) sf –1   |      |   | B2<br>B1       |
|                                   | Allow rotation through 540 centre the origin   |      |   | B2             |
|                                   | Do not allow if there is additional information that is incorrect<br>eg1 Rotation through 180° and a reflection<br>eg2 Enlargement sf –1 rotated through 90°<br>eg3 Rotation through 180° centre the origin so the shape turns |      |   | B0<br>B0<br>B2 |
|                                   | Rotation   |      |   | B0             |
|                                   | Enlargement  |      |   | B0             |
|                                   | Do not allow turn for rotation   |      |   |                |
|                                   | Do not allow eg half turn for 180°   |      |   |                |
| Do not allow negative enlargement |  |      |   |                |

| Q  | Answer  | Mark  | Comments   |
|----|---|-------|--|
| 15 | <b>Alternative method 1</b>   |       |  |
|    | angle $ABO = x$   | M1    | may be seen on diagram<br>implied by angle $AOB = 180 - 2x$  |
|    | angle $ACB = 180 - w$   | M1    | oe eg angle $ACB + w = 180$<br>may be seen on diagram  |
|    | angle $AOB = 2 \times (180 - w)$<br>or angle $AOB = 360 - 2w$             | M1dep | may be seen on diagram<br>dep on 2nd M1<br>angle $AOB$ may be seen as $180 - 2x$   |
|    | $x + x + 2 \times (180 - w) = 180$  | M1dep | oe eg $2(180 - w) = 180 - 2x$<br>dep on M3   |
|    | $w = x + 90$ with M4<br>and<br>all reasons given                          | A1    | eg of reasons<br>isosceles triangle<br>and angles on a straight line<br>and angle at centre<br>and angle sum of triangle |
|    | <b>Alternative method 2</b>   |       |  |
|    | angle $ABO = x$   | M1    | may be seen on diagram<br>implied by angle $AOB = 180 - 2x$  |
|    | angle $AOB = 180 - x - x$<br>or angle $AOB = 180 - 2x$                    | M1dep | oe eg $2x + \text{angle } AOB = 180$<br>may be seen on diagram   |
|    | angle $ACB = \frac{1}{2} \times (180 - x - x)$<br>or angle $ACB = 90 - x$ | M1dep | oe eg angle $ACB = \frac{1}{2} \times (180 - 2x)$<br>may be seen on diagram<br>angle $ACB$ may be seen as $180 - w$      |
|    | $\frac{1}{2} \times (180 - x - x) + w = 180$                              | M1dep | oe eg $w = 180 - (90 - x)$   |
|    | $w = x + 90$ with M4<br>and<br>all reasons given                          | A1    | eg of reasons<br>isosceles triangle<br>and angle sum of triangle<br>and angle at centre<br>and angles on a straight line |

**Mark scheme and Additional Guidance continues on the next two pages**

| Q                  | Answer  | Mark  | Comments   |
|--------------------|---|-------|--|
| <b>15<br/>cont</b> | <b>Alternative method 3</b> Draws tangent (eg $PQ$ ) at $A$ |       |  |
|                    | angle $QAB = 90 - x$  | M1    | oe eg $x + \text{angle } QAB = 90$<br>may be seen on diagram   |
|                    | angle $ACB = 180 - w$                                       | M1    | oe eg angle $ACB + w = 180$<br>may be seen on diagram  |
|                    | angle $QAB = \text{angle } ACB$                             | M1    | may be seen on diagram<br>eg both angles labelled $y$  |
|                    | $90 - x = 180 - w$  | M1dep | oe eg $90 - x + w = 180$<br>dep on M3  |
|                    | $w = x + 90$ with M4<br>and<br>all reasons given            | A1    | eg of reasons<br>radius perpendicular to tangent<br>and angles on a straight line<br>and alternate segment |

**Additional Guidance is on the next page**

|                    |  | <b>Additional Guidance</b>  |  |
|--------------------|--|---|--|
| <b>15<br/>cont</b> |  | Allow angle $BCD$ for $w$ throughout  |  |
|                    |  | 3rd M1 and 4th M1 may be seen in one line of working<br>eg1 Alt 1<br>angle $ABO = x$<br>angle $ACB = 180 - w$<br>$180 - 2x = 2 \times (180 - w)$<br><br>eg2 Alt 2<br>angle $ABO = x$<br>angle $AOB = 180 - 2x$<br>$180 - w = \frac{1}{2} \times (180 - 2x)$ | M1<br>M1<br>M1M1<br><br>M1<br>M1<br>M1M1 |
|                    |  | Condone slips in notation only if angles are marked in correct position on the diagram<br>eg1 Do not allow angle $C = 180 - w$ unless marked in correct position on the diagram<br>eg2 Allow $ACB$ for angle $ACB$  |  |
|                    |  | For reasons, allow if the intention is clear<br>eg1 Allow isos triangle for isosceles triangle<br>eg2 Allow angles in a triangle for angle sum of a triangle<br>eg3 Allow angles on a line for angles on a straight line                                    |  |
|                    |  | For reasons do not allow incorrect statements<br>eg do not allow angles in a triangle add to 360  |  |

| Q   | Answer   | Mark  | Comments  |
|---|--|-------|---|
| 16  | $15 \times 2^4$ or $15 \times 16$ or 240   | M1    | oe eg $\binom{6}{4}2^4$ or $2^6 \times \frac{6 \times 5}{2} \times \left(\frac{1}{2}\right)^2$<br>may include $a^2$ and/or $x^4$<br>allow embedded eg ${}^6C_4 a^2(2x)^4$ |
|   | $240a^2 = 1500$ or $a^2 = \frac{1500}{240}$<br>or $(\pm)\sqrt{\frac{1500}{240}}$<br>or $\frac{5}{2}$ or $-\frac{5}{2}$ | M1dep | must evaluate $\binom{6}{4}$<br>oe eg $15 \times 2^4 a^2 = 1500$ or $(\pm)\sqrt{\frac{1500}{15 \times 2^4}}$<br>may include $x^4$ on both sides of an equation            |
|   | $\frac{5}{2}$ and $-\frac{5}{2}$<br>with no other values   | A1    | oe eg 2.5 and -2.5<br>SC2 [2.236, 2.24] and [-2.24, -2.236]<br>SC1 [2.236, 2.24] or [-2.24, -2.236]   |
|   | <b>Additional Guidance</b>   |       |   |
|   | The relevant term must be selected from a full expansion but the other terms can be ignored                            |       |   |
|   | Allow $\binom{6}{4}$ to be $\binom{6}{2}$  |       |   |
|   | $240a^2x^4 = 1500x^4$  |       | M1M1  |
| $240a^2x^4 = 1500$ recovered to $(\pm)\sqrt{\frac{1500}{240}}$ oe     |  | M1M1  |   |
| $240a^2x^4 = 1500$ not recovered to $(\pm)\sqrt{\frac{1500}{240}}$ oe |  | M1M0  |   |

| Q            | Answer   | Mark  | Comments   |
|--------------|--|-------|--|
| 17           | <b>Alternative method 1</b> Works out $MD$ and $BD$ and uses $\tan MBD$  |       |  |
|              | $\tan 28 = \frac{GN}{32}$ or $32 \tan 28$<br>or [17, 17.015]   | M1    | oe eg $\frac{32}{\tan(90 - 28)}$<br>working out $GN$ or $HM$   |
|              | $32 - 32 \tan 28$ or [14.985, 15]  | M1dep | oe<br>working out $NC$ or $MD$   |
|              | $\sqrt{32^2 + 32^2}$ or $\sqrt{2048}$<br>or [45.2, 45.3]   | M1    | oe eg $32\sqrt{2}$<br>working out $BD$   |
|              | $\tan MBD = \frac{\text{their [14.985, 15]}}{\text{their [45.2, 45.3]}}$   | M1dep | oe eg $\tan^{-1} \frac{\text{their [14.985, 15]}}{\text{their [45.2, 45.3]}}$<br>dep on M3                         |
|              | [18.3, 18,4]   | A1    |  |
|              | <b>Alternative method 2</b> Works out $BD$ and $MB$ and uses $\cos MBD$  |       |  |
|              | $\tan 28 = \frac{GN}{32}$ or $32 \tan 28$<br>or [17, 17.015]   | M1    | oe eg $\frac{32}{\tan(90 - 28)}$<br>working out $GN$ or $HM$   |
|              | $32 - 32 \tan 28$ or [14.985, 15]  | M1dep | oe<br>working out $NC$ or $MD$   |
|              | $\sqrt{32^2 + 32^2}$ or $\sqrt{2048}$<br>or [45.2, 45.3]<br>or<br>$\sqrt{32^2 + 32^2 + \text{their [14.985, 15]}^2}$<br>or [47.67, 47.7] | M1    | oe eg $32\sqrt{2}$<br>working out $BD$ or $MB$<br>if awarding this mark for working out $MB$ it is dependent on M2 |
|              | $\cos MBD =$<br>$\frac{\sqrt{32^2 + 32^2}}{\sqrt{32^2 + 32^2 + \text{their [14.985, 15]}^2}}$  | M1dep | oe<br>eg $\cos^{-1} \frac{\sqrt{32^2 + 32^2}}{\sqrt{32^2 + 32^2 + \text{their [14.985, 15]}^2}}$<br>dep on M3      |
| [18.3, 18,4] | A1   |       |  |

Mark scheme and Additional Guidance continue on the next page

|  |  |       |   |
|--|--|-------|---|
| <b>17<br/>cont</b>   | <b>Alternative method 3</b> Works out $MD$ and $MB$ and uses $\sin MBD$  |       |   |
|  | $\tan 28 = \frac{GN}{32}$ or $32 \tan 28$<br>or $[17, 17.015]$   | M1    | oe eg $\frac{32}{\tan(90 - 28)}$<br>working out $GN$ or $HM$  |
|  | $32 - 32 \tan 28$ or $[14.985, 15]$  | M1dep | oe<br>working out $NC$ or $MD$  |
|  | $\sqrt{32^2 + 32^2}$ + their $[14.985, 15]^2$<br>or $[47.67, 47.7]$  | M1dep | oe<br>working out $MB$  |
|  | $\sin MBD =$<br>$\frac{\text{their } [14.985, 15]}{\sqrt{32^2 + 32^2 + \text{their } [14.985, 15]^2}}$                             | M1dep | oe<br>eg $\sin^{-1} \frac{\text{their } [14.985, 15]}{\sqrt{32^2 + 32^2 + \text{their } [14.985, 15]^2}}$ |
|  | $[18.3, 18.4]$   | A1    |   |
|  | <b>Additional Guidance</b>   |       |   |
|  | 1st M1 $GN$ may be seen as a letter, eg $x$ , but do not award if subsequently used as the length of an incorrect side (eg $MN$ )  |       |   |
|  | 4th M1 $MBD$ may be seen as a letter, eg $y$ , but do not award if subsequently used as the size of an incorrect angle (eg $DMB$ ) |       |   |
|  | Alt 1 or Alt 2 $32\sqrt{1^2 + 1^2}$  |       | 3rd M1  |
| Alt 1 $\tan MBD = \frac{32(1 - \tan 28)}{32\sqrt{2}}$ or $\tan MBD = \frac{(1 - \tan 28)}{\sqrt{2}}$ |  | M4    |   |

| Q  | Answer   | Mark  | Comments  |
|----|--|-------|---|
| 18 | <b>Alternative method 1</b>  |       |   |
|    | 12 or $-3x^{-2}$   | M1    | oe eg $12x^0$ or $3 \times -1x^{-1-1}$ or $-\frac{3}{x^2}$  |
|    | 12 and $-3x^{-2}$  | M1dep | oe eg $12 - \frac{3}{x^2}$<br>or $12x^0$ and $3 \times -1x^{-1-1}$  |
|    | $12 - 3x^{-2} = 0$ and $x = 0.5$<br>or<br>$12 - 3 \times 0.5^{-2} = 0$ | M1dep | oe<br>$= 0$ must be seen<br>condone inclusion of $x = -0.5$   |
|    | $6x^{-3}$  | M1    | oe eg $-2 \times -3x^{-2-1}$<br>ft differentiation of their first derivative if it involves a negative power of $x$ |
|    | M4 and $6 \times 0.5^{-3} (= 48)$ which is positive (so minimum)       | A1    | oe<br>do not allow if $\frac{6}{0.5^3}$ is evaluated incorrectly  |

**Mark scheme and Additional Guidance continue on the next page**

|                    |   |       |  |
|--------------------|---|-------|--|
| <b>18<br/>cont</b> | <b>Alternative method 2</b>   |       |  |
|                    | 12 or $-3x^{-2}$  | M1    | oe eg $12x^0$ or $3 \times -1x^{-1-1}$   |
|                    | 12 and $-3x^{-2}$   | M1dep | oe eg $12 - \frac{3}{x^2}$<br>or $12x^0$ and $3 \times -1x^{-1-1}$   |
|                    | $12 - 3x^{-2} = 0$ and $x = 0.5$<br>or<br>$12 - 3 \times 0.5^{-2} = 0$  | M1dep | oe<br>= 0 must be seen<br>condone inclusion of $x = -0.5$  |
|                    | Substitutes one $x$ value in range<br>(0, 0.5) into $12 - 3x^{-2}$<br>and<br>substitutes one $x$ value $> 0.5$<br>into $12 - 3x^{-2}$                 | M1    | eg $12 - 3 \times 0.25^{-2}$<br>and<br>$12 - 3 \times 1^{-2}$<br>ft substitution into their first derivative if it<br>involves a negative power of $x$   |
|                    | M4 and two correct evaluations<br>(so minimum)<br>or<br>M4 and two correct signs shown<br>with no incorrect evaluations (so<br>minimum)               | A1    | eg M4 and $12 - 3 \times 0.25^{-2} = -36$<br>and<br>$12 - 3 \times 1^{-2} = 9$ (so minimum)<br>or<br>M4 and $12 - 3 \times 0.25^{-2}$ is negative<br>and $12 - 3 \times 1^{-2}$ is positive (so minimum) |
|                    | <b>Additional Guidance</b>  |       |  |
|                    | Alt 1<br>$12 + 3x^{-2} = 0$<br>$-6x^{-3}$   |       | M1M0M0<br>M1A0   |
|                    | Alt 2<br>$12 - 3x^{-2}$<br>$6x^{-3}$<br>$12 - 3 \times 0.25^{-2} = -36$ $12 - 3 \times 1^{-2} = 9$ so minimum<br>(A1 only possible after awarding M4) |       | M1M1M0<br>M1<br>A0   |
|                    | Ignore any testing of the stationary point at $x = -0.5$  |       |  |

| Q     | Answer                     | Mark | Comments |
|-------|----------------------------|------|----------|
| 19(a) | $x^4$                      | B1   |          |
|       | <b>Additional Guidance</b> |      |          |
|       |                            |      |          |

| Q     | Answer   | Mark | Comments  |
|-------|--|------|---|
| 19(b) | $2x^2 + 10$ or $2(x^2 + 5)$                    | B2   | B1 $k(x) = 2x$ or $(k(x))^2 = 4x^2$<br>or $h(2x) = 4x^2 + 5$<br>or $(2x)^2 + 5$ |
|       | <b>Additional Guidance</b>                     |      |   |
|       | $2(x^2 + 5)$ in working with answer $2x^2 + 5$ |      | B1  |

| Q  | Answer  | Mark  | Comments   |
|----|---|-------|--|
| 20 | <b>Alternative method 1</b> Uses a common denominator of $\sin x$   |       |  |
|    | $\frac{2 \sin x + \cos x}{\sin x} - \frac{1}{\cos x}$ or $\frac{\cos x(2 \sin x + \cos x)}{\sin x} - \frac{1}{\sin x}$  | M1    | implied by 2nd M1<br>condone omission of $-\frac{1}{\sin x}$ |
|    | $\frac{2 \sin x \cos x + \cos^2 x}{\sin x} - \frac{1}{\sin x}$ or $2 \cos x + \frac{\cos^2 x}{\sin x} - \frac{1}{\sin x}$   | M1dep | condone omission of $-\frac{1}{\sin x}$                      |
|    | $\frac{2 \sin x \cos x + \cos^2 x - \cos^2 x - \sin^2 x}{\sin x}$ or $\frac{2 \sin x \cos x + 1 - \sin^2 x - 1}{\sin x}$ or $\frac{2 \sin x \cos x - \sin^2 x}{\sin x}$ or $2 \cos x - \frac{\sin^2 x}{\sin x}$ | M1dep |  |
|    | $2 \cos x - \sin x$ with M3   | A1    |  |

Mark scheme and Additional Guidance continues on the next two pages

|                    |   |       |   |
|--------------------|---|-------|---|
| <b>20<br/>cont</b> | <b>Alternative method 2</b> Uses a common denominator of $\tan x \sin x$  |       |   |
|                    | $\frac{\sin x (2 \sin x + \cos x) - \tan x}{\tan x \sin x}$ or $\frac{2 \sin^2 x + \cos x \sin x - \tan x}{\tan x \sin x}$  | M1    | $\tan x$ may be seen as $\frac{\sin x}{\cos x}$   |
|                    | $\frac{2 \sin^2 x \cos x + \cos^2 x \sin x - \sin x}{\sin^2 x}$   | M1dep |   |
|                    | $\frac{2 \sin x \cos x + \cos^2 x - \cos^2 x - \sin^2 x}{\sin x}$ or $\frac{2 \sin x \cos x + 1 - \sin^2 x - 1}{\sin x}$ or $\frac{2 \sin x \cos x - \sin^2 x}{\sin x}$ | M1dep | allow the fractions with denominator $\sin^2 x$<br>eg $\frac{2 \sin^2 x \cos x + \sin x - \sin^3 x - \sin x}{\sin^2 x}$ |
|                    | $2 \cos x - \sin x$ with M3   | A1    |   |

**Additional Guidance is on the next page**

| <b>Additional Guidance</b>  |   |
|---|---|
| $2 \cos x - \sin x$ with no method  | Zero  |
| Allow eg $1 \sin x$ for $\sin x$  |   |
| Condone $x$ missing for M marks and award A mark if recovered   |   |
| <p>Fractions may be seen as separate fractions/terms</p> <p>eg1 Alt 1</p> $\frac{2 \sin x}{\sin x} + \frac{\cos x}{\sin x} - \frac{1}{\sin x}$ $\frac{2 \sin x}{\cos x} + \frac{\cos x}{\cos x} - \frac{1}{\sin x}$ $2 \cos x + \frac{\cos^2 x}{\sin x} - \frac{1}{\sin x}$ $2 \cos x + \frac{1}{\sin x} - \frac{\sin^2 x}{\sin x} - \frac{1}{\sin x}$ <p>eg2 Alt 2</p> $\frac{2 \sin^2 x + \cos x \sin x}{\tan x \sin x} - \frac{\tan x}{\tan x \sin x}$ $\frac{2 \sin^2 x \cos x + \cos^2 x \sin x}{\sin^2 x} - \frac{\sin x}{\sin^2 x}$ $\frac{2 \sin x \cos x + \cos^2 x}{\sin x} - \frac{\cos^2 x + \sin^2 x}{\sin x}$ | <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> |

| Q   | Answer  | Mark  | Comments  |
|---|---|-------|---|
| <b>21</b>   | <b>Alternative method 1</b>   |       |   |
|   | $3(x^2 + ax + ax + a^2) \dots$<br>or $3(x^2 + 2ax + a^2) \dots$<br>or $3\left(x + \frac{b}{3}\right)^2 \dots$<br>or $2b = 6a$<br>or $8a = 3a^2 + b + 2$ | M1    | oe eg $3x^2 + 6ax + 3a^2 \dots$<br>or $\frac{b}{3} = a$<br>or $b + 2 = -3\left(\frac{b}{3}\right)^2 + 8a$ |
|   | $2b = 6a$<br>and $8a = 3a^2 + b + 2$  | M1dep | oe equations<br>eg $\frac{b}{3} = a$ and $b + 2 = -3\left(\frac{b}{3}\right)^2 + 8a$                      |
|   | $3a^2 + 3a - 8a + 2 (= 0)$<br>or $3a^2 - 5a + 2 (= 0)$  | M1dep | oe quadratic equation in $a$  |
|   | $(3a - 2)(a - 1)$<br>or $\frac{-5 \pm \sqrt{(-5)^2 - 4 \times 3 \times 2}}{2 \times 3}$   | M1    | oe eg $\frac{5}{6} \pm \sqrt{\frac{25}{36} - \frac{2}{3}}$<br>ft their 3-term quadratic                   |
|   | $a = \frac{2}{3}$ and $a = 1$<br>or<br>$a = \frac{2}{3}$ and $b = 2$<br>or<br>$a = 1$ and $b = 3$   | A1    |   |
| $a = \frac{2}{3}$ and $b = 2$<br>and<br>$a = 1$ and $b = 3$ | A1  |       |   |

**Mark scheme and Additional Guidance continue on the next page**

|   |   |       |   |
|---|---|-------|---|
| <b>21<br/>cont</b>  | <b>Alternative method 2</b>   |       |   |
|   | $3(x^2 + ax + ax + a^2) \dots$<br>or $3(x^2 + 2ax + a^2) \dots$<br>or $3\left(x + \frac{b}{3}\right)^2 \dots$<br>or $2b = 6a$<br>or $8a = 3a^2 + b + 2$ | M1    | oe eg $3x^2 + 6ax + 3a^2 \dots$<br>or $\frac{b}{3} = a$<br>or $b + 2 = -3\left(\frac{b}{3}\right)^2 + 8a$ |
|   | $2b = 6a$<br>and $8a = 3a^2 + b + 2$  | M1dep | oe equations<br>eg $\frac{b}{3} = a$ and $b + 2 = -3\left(\frac{b}{3}\right)^2 + 8a$                      |
|   | $\frac{8b}{3} = 3\left(\frac{b}{3}\right)^2 + b + 2$<br>or $b^2 - 5b + 6 (= 0)$   | M1dep | oe quadratic equation in $b$  |
|   | $(b - 2)(b - 3)$<br>or $\frac{-5 \pm \sqrt{(-5)^2 - 4 \times 1 \times 6}}{2 \times 1}$  | M1    | oe eg $\frac{5}{2} \pm \sqrt{\frac{25}{4} - 6}$<br>ft their 3-term quadratic                              |
|   | $b = 2$ and $b = 3$<br>or<br>$a = \frac{2}{3}$ and $b = 2$<br>or<br>$a = 1$ and $b = 3$   | A1    |   |
|   | $a = \frac{2}{3}$ and $b = 2$<br>and<br>$a = 1$ and $b = 3$   | A1    |   |
|   | <b>Additional Guidance</b>  |       |   |
|   | Allow 0.6 for $\frac{2}{3}$   |       |   |
|   | Allow 0.67 for $\frac{2}{3}$ for first A1   |       |   |
| In quadratic formula allow $5^2$ for $(-5)^2$ but use of $-5^2$ must be recovered |   |       |   |