AQA

Please write clearly in block capitals.

Centre number ☐☐☐☐☐   Candidate number ☐☐☐☐

Surname _____

Forename(s) _____

Candidate signature _____
I declare this is my own work.

A-level
FURTHER MATHEMATICS

Paper 3 Mechanics

Time allowed: 2 hours

Materials
- You must have the AQA Formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a scientific calculator that meets the requirements of the specification. (You may use a graphical calculator.)
- You must ensure you have the other optional Question Paper/Answer Book for which you are entered (**either** Discrete **or** Statistics). You will have 2 hours to complete **both** papers.

Instructions
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 50.

Advice
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
TOTAL	

Answer **all** questions in the spaces provided.

1   A spring of natural length 50 cm and modulus of elasticity λ newtons has an elastic potential energy of 4 J when compressed by 5 cm.

Find the value of λ

Circle your answer.

[1 mark]

   8   16   800   1600

3

2 A force of magnitude 7 N acts at each end of a rod of length 20 cm, forming a couple.

The forces act at right angles to the rod, as shown in the diagram below.

7 N

20 cm

7 N

Find the magnitude of the resultant moment of the couple.

Circle your answer.

[1 mark]

1.4 N m   2.8 N m   140 N m   280 N m

Turn over for the next question

3 A ball has mass 0.4 kg and is hit by a wooden bat.

The speed of the ball just before it is hit by the bat is $6\,\text{m s}^{-1}$

The velocity of the ball immediately after being hit by the bat is perpendicular to its initial velocity.

The speed of the ball just after it is hit by the bat is $8\,\text{m s}^{-1}$

Show that the impulse on the ball has magnitude $4\,\text{N s}$

[3 marks]

4   A spring has stiffness k

4 (a)   Determine the dimensions of k

[1 mark]

_____

_____

4 (b)   One end of the spring is attached to a fixed point. A particle of mass m kg is attached to the other end of the spring.

The particle is set into vertical motion and moves up and down, taking t seconds to complete one oscillation.

A possible model for t is

$$t = pm^a g^b k^c$$

where p is a dimensionless constant and $g\,\mathrm{m\,s}^{-2}$ is the acceleration due to gravity.

Find the values of a, b and c for this model to be dimensionally consistent.

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

Turn over ▶

5   A uniform lamina has the shape of the region enclosed by the curve $y = x^2 + 1$ and the lines $x = 0$, $x = 4$ and $y = 0$

The diagram below shows the lamina.

5 (a)   Find the coordinates of the centre of mass of the lamina, giving your answer in exact form.

[4 marks]

_____

_____

_____

_____

_____

_____

_____

_____

5 (b) The lamina is suspended from the point where the curve intersects the line $x = 4$ and hangs in equilibrium.

Find the angle between the vertical and the longest straight edge of the lamina, giving your answer correct to the nearest degree.

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

Turn over ▶

There are no questions printed on this page

6   A ball of mass m kg is held at rest at a height h metres above a horizontal surface.

The ball is released and bounces on the surface.

The coefficient of restitution between the ball and the surface is e

Prove that the kinetic energy lost during the first bounce is given by

$$mgh(1 - e^2)$$

[4 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

Turn over ▶

7 A light string has length 1.5 metres.

A small sphere is attached to one end of the string.

The other end of the string is attached to a fixed point *O*

A thin horizontal bar is positioned 0.9 metres directly below *O*

The bar is perpendicular to the plane in which the sphere moves.

The sphere is released from rest with the string taut and at an angle α to the downward vertical through *O*

The string becomes slack when the angle between the two sections of the string is 60°

Ben draws the diagram below to show the initial position of the sphere, the bar and the path of the sphere.

7 (a) State two reasons why Ben's diagram is not a good representation of the situation.

[2 marks]

Reason 1 _____

_____

_____

Reason 2 _____

_____

_____

7 (b)   Using your answer to part **(a)**, sketch an improved diagram.

[1 mark]

O

Bar ●

Question 7 continues on the next page

12

7 (c)   Find α, giving your answer to the nearest degree.

[6 marks]

Turn over for the next question

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Turn over ▶

8 **In this question use $g = 9.8\,\mathrm{m\,s^{-2}}$**

A lift is used to raise a crate of mass 250 kg

The lift exerts an upward force of magnitude P newtons on the crate.

When the crate is at a height of x metres above its initial position

$$P = k(x + 1)(12 - x) + 2450$$

where k is a constant.

The crate is initially at rest, at the point where $x = 0$

8 (a) Show that the work done by the upward force as the crate rises to a height of 12 metres is given by

$$29\,400 + 360k$$

[3 marks]

_____

_____

_____

_____

_____

_____

_____

8 (b) The speed of the crate is $3\,\mathrm{m\,s^{-1}}$ when it has risen to a height of 12 metres.

Find the speed of the crate when it has risen to a height of 15 metres.

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

8 (c)   Find the height of the crate when its speed becomes zero.

[2 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

8 (d)   Air resistance has been ignored.

Explain why this is reasonable in this context.

[1 mark]

_____

_____

Turn over ▶

16

9   **In this question use $g = 9.81\,\mathrm{m\,s}^{-2}$**

A conical pendulum is made from an elastic string and a sphere of mass 0.2 kg

The string has natural length 1.6 metres and modulus of elasticity 200 N

The sphere describes a horizontal circle of radius 0.5 metres at a speed of $v\,\mathrm{m\,s}^{-1}$

The angle between the elastic string and the vertical is α

9 (a)   Show that

$$62.5 - 200\sin\alpha = 1.962\tan\alpha$$

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

9 (b) Use your calculator to find α

[1 mark]

_____

_____

_____

9 (c) Find the value of v

[4 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

END OF QUESTIONS

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

Question number	Additional page, if required. Write the question numbers in the left-hand margin.

Question number	Additional page, if required. Write the question numbers in the left-hand margin.
2 0	