



Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname _____

Forename(s) _____

Candidate signature _____

I declare this is my own work.

A-level PHYSICS

Paper 3 Section B Astrophysics

Materials

For this paper you must have:

- a pencil and a ruler
- a scientific calculator
- a Data and Formulae Booklet
- a protractor.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.
- Show all your working.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 35.
- You are expected to use a scientific calculator where appropriate.
- A Data and Formulae Booklet is provided as a loose insert.

Time allowed: The total time for both sections of this paper is 2 hours. You are advised to spend approximately 50 minutes on this section.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
TOTAL	



Section B

Answer **all** questions in this section.

0	1
---	---

 .

1

 Draw a ray diagram to show how a converging lens can cause spherical aberration. **[1 mark]**

_____ principal axis

0	1
---	---

 .

2

 Draw a labelled ray diagram for an astronomical refracting telescope in normal adjustment. **[3 marks]**

Show **three** non-axial rays passing through both lenses.
Label the principal foci of the lenses.

_____ principal axis



There are no questions printed on this page

*Do not write
outside the
box*


**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



0 2

U Cephei is an eclipsing binary system consisting of two stars that orbit their common centre of mass.
The primary star is class B; the secondary star is class G.

Figure 1 shows the variation of apparent magnitude of U Cephei with time as observed from Earth.



0 2 . 1

Explain the shape of the graph in **Figure 1**.

[2 marks]

Question 2 continues on the next page

Turn over ►



A particular spectral line has a wavelength of 486.136 nm when measured from a source in the laboratory.
This line is also present in the absorption spectrum of the primary star of U Cephei.
When observed from Earth, the wavelength of the primary star’s absorption line varies as shown in **Table 1**.

Table 1

	Wavelength / nm
maximum value	486.498
minimum value	485.672

0 2 . 2 State why the average of the values in **Table 1** is different from the laboratory value. **[1 mark]**

0 2 . 3 Show that the orbital speed of the primary star is about 250 km s⁻¹. **[3 marks]**



Do not write
outside the
box

0 2 . 4 Calculate the orbital radius of the primary star.

[2 marks]

orbital radius = _____ m

0 2 . 5 Which absorption lines would be most prominent in the spectrum of the primary star?
Tick (✓) **one** box.

[1 mark]

- hydrogen
- hydrogen and helium
- ionised metals
- neutral metals

0 2 . 6 A different eclipsing binary star system is thought to consist of a white dwarf star and a neutron star.

Discuss how astronomers could confirm this.

[2 marks]



03

3C 273 was the first quasar to be discovered.
IC 1101 is one of the largest galaxies known.
Table 2 shows some information about these objects.

Table 2

	Absolute magnitude	Apparent magnitude	Distance / Mpc
quasar 3C 273	X	12.8	760
galaxy IC 1101	−22.8	14.7	320

03.1

State the property of the quasar that led to its discovery.

[1 mark]

03.2

Show that the absolute magnitude **X** of quasar 3C 273 is about −27

[2 marks]



Do not write
outside the
box

03.3

Assume that the quasar and the galaxy are both viewed from the same distance.

Explain which would be the brighter object.

Go on to calculate the ratio $\frac{\text{brightness of brighter object}}{\text{brightness of dimmer object}}$.

[3 marks]

ratio = _____

03.4

The black hole at the centre of IC 1101 has a mass of $7.1 \times 10^{11} M_{\text{S}}$ where M_{S} is the mass of the Sun.

Calculate the average density within the event horizon of the black hole.

[3 marks]

average density = _____ kg m^{-3}

9


Turn over ►



0 4

In the middle of the 20th century, there were two competing theories of the Universe. In 1964, electromagnetic radiation was observed coming from all directions in space. **Figure 2** shows the distribution of this radiation as observed from Earth.

Figure 2



The graph provides evidence for one of these theories of the Universe.

Discuss the main features of this theory of the Universe.

In your answer, you should include:

- the main predictions and evidence for the theory, and
- a suitable calculation.

[6 marks]



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



Do not write
outside the
box

Question number	Additional page, if required. Write the question numbers in the left-hand margin.



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from www.aqa.org.uk.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2022 AQA and its licensors. All rights reserved.



1 6



2 2 6 A 7 4 0 8 / 3 B A

IB/M/Jun22/7408/3BA