

Friday 26 May 2023 – Afternoon

AS Level Further Mathematics A

Y534/01 Discrete Mathematics

Time allowed: 1 hour 15 minutes

You must have:

- the Printed Answer Booklet
- the Formulae Booklet for AS Level Further Mathematics A
- · a scientific or graphical calculator






- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided in the Printed Answer
 Booklet. If you need extra space use the lined pages at the end of the Printed Answer
 Booklet. The question numbers must be clearly shown.
- Fill in the boxes on the front of the Printed Answer Booklet.
- Answer all the guestions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.
- Give non-exact numerical answers correct to **3** significant figures unless a different degree of accuracy is specified in the question.
- The acceleration due to gravity is denoted by $g \, \text{m} \, \text{s}^{-2}$. When a numerical value is needed use g = 9.8 unless a different value is specified in the question.
- Do not send this Question Paper for marking. Keep it in the centre or recycle it.

INFORMATION

- The total mark for this paper is **60**.
- The marks for each question are shown in brackets [].
- This document has 8 pages.

ADVICE

Read each question carefully before you start your answer.



1 Jane wants to travel from home to the local town.


Jane can do this by train, by bus or by both train and bus.

(a) Give an example of a problem that Jane could be answering that would give a construction problem. [1]

A website gives Jane all the possible buses and trains that she could use. Jane finds 7 possible ways to make the journey.

- 2 of the 7 journeys involve travelling by train for at least part of the journey
- 6 of the 7 journeys involve travelling by bus for at least part of the journey
- (b) Use the inclusion-exclusion principle to find how many of the 7 journeys involve travelling by both train and bus. [2]

2 A network is shown below.



- (a) Use an appropriate algorithm to find the least weight (shortest) path from A to D. [3]
- (b) Use Kruskal's algorithm to find a minimum spanning tree for the network. [3]

3

The list of numbers below is to be sorted into increasing order.											
23	10	18	7	62	54	31	82				
(a)	(a) Sort the list using bubble sort. You do not need to show intermediate working.										
	(i)	Record the	e list th	at resul	ts at the	end o	f each pass	5.			[3]
	(ii) Record the number of swaps used in each pass.									[2	
(b)	(b) Now sort the original list using shuttle sort. You do not need to show intermediate working.										
	(i)	Record the	e list th	at resul	ts at the	end o	f each pass	3.			[3]
	(ii)	Record the	e numb	er of sw	aps use	ed in e	ach pass.				[2
(c)	(c) Using the total number of comparisons plus the total number of swaps as a measure of efficiency, explain why shuttle sort is more efficient than bubble sort for sorting this particular list.							[1]			
Bubble sort and shuttle sort are both $O(n^2)$.											
(d)	Explain what this means for the run-time of the algorithms when the length of the list being sorted changes from 1000 to 3000.							ing [1]			

- 4 Graph G is a simply connected Eulerian graph with 4 vertices.
 - (a) (i) Explain why graph G cannot be a complete graph. [2]
 - (ii) Determine the number of arcs in graph G, explaining your reasoning. [2]
 - (iii) Show that graph G is a bipartite graph. [1]

Graph H is a digraph with 4 vertices and no undirected arcs. The adjacency matrix below shows the number of arcs that directly connect each pair of vertices in digraph H.

		To			
		A	В	C	D
From	A	0	1	0	1
	В	0	0	2	0
	С	2	1	0	1
	D	0	1	1	0
	С	0 2 0	0 1 1		0 1 0

- (b) (i) Write down a feature of the adjacency matrix that shows that H has no loops. [1]
 - (ii) Find the number of arcs in H. [1]
 - (iii) Draw a possible digraph H. [1]
 - (iv) Show that digraph H is semi-Eulerian by writing down a suitable trail. [2]

5 Hiro has been asked to organise a quiz.

The table below shows the activities involved, together with the immediate predecessors and the duration of each activity in hours.

	Activity	Immediate predecessors	Duration (hours)
A	Choose the topics	_	0.5
В	Find questions for round 1	A	2
С	Check answers for round 1	В	2.5
D	Find questions for round 2	A	2
Е	Check answers for round 2	D	2.5
F	Choose pictures for picture round	A	1
G	Get permission to use pictures	F	1.5
Н	Choose music for music round	A	2
I	Get permission to use music	Н	1.5
J	Produce answer sheets	G	0.5

(a) A sketch of the activity network is provided in the Printed Answer Booklet.

Apply a forward pass to determine the minimum project completion time. [2]

- (b) Use a backward pass to determine the critical activities.

 You can show your working on the activity network from part (a).
- (c) Give the total float for each non-critical activity. [3]

Hiro decides that there should be a final check of the answers which he will include as activity L.

Activity L needs to be done after checking the answers for rounds 1 and 2 and also after getting permission to use the pictures and music but before producing the answer sheets.

- (d) (i) Complete the activity network provided in the Printed Answer Booklet to show the new precedences, with the final check of the answers included as activity L. [2]
 - (ii) As a result of including L, the minimum project completion time found in part (a) increases by 2.5 hours.

Determine the duration of L. [2]

- 6 Ryan and Casey are playing a card game in which they each have four cards.
 - Ryan's cards have the letters A, B, C and D.
 - Casey's cards have the letters W, X, Y and Z.

Each player chooses one of their four cards and they simultaneously reveal their choices.

The table shows the number of points won by Ryan for each combination of strategies.

		Casey				
		W	X	Y	Z	
Ryan	A	4	0	2	1	_
	В	0	2	-3	4	
	C	1	4	-4	5	
	D	6	-1	5	0	

For example, if Ryan chooses A and Casey chooses W then Ryan wins 4 points (and Casey loses 4 points).

Both Ryan and Casey are trying to win as many points as possible.

- (a) Use dominance to reduce the 4×4 table for the zero-sum game above to a 4×2 table. [2]
- (b) Determine an optimal mixed strategy for Casey. [4]

7

7 A linear programming problem is

Maximise
$$P = 4x + y$$

subject to $3x - y \le 30$
 $x + y \le 15$
 $x - 3y \le 6$
and $x \ge 0$, $y \ge 0$

(a) Use a graphical method to find the optimal value of P, and the corresponding values of x and y.

An additional constraint is introduced.

This constraint means that the value of y must be at least k times the value of x, where k is a positive constant.

- (b) (i) Determine the set of values of k for which the optimal value of P found in part (a) is unchanged. [2]
 - (ii) Determine, in terms of k, the values of x, y and P in the cases when the optimal solution is different from that found in part (a). [4]

END OF QUESTION PAPER



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.