![AQA logo]

Please write clearly in block capitals.

Centre number ☐☐☐☐☐   Candidate number ☐☐☐☐

Surname _____

Forename(s) _____

Candidate signature _____
I declare this is my own work.

A-level
FURTHER MATHEMATICS

Paper 2

Monday 5 June 2023   Afternoon   Time allowed: 2 hours

Materials
- You must have the AQA Formulae and statistical tables booklet for A-level Mathematics and A-level Further Mathematics.
- You should have a graphical or scientific calculator that meets the requirements of the specification.

Instructions
- Use black ink or black ball-point pen. Pencil should only be used for drawing.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer each question in the space provided for that question. If you require extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do **not** write outside the box around each page or on blank pages.
- Show all necessary working; otherwise marks for method may be lost.
- Do all rough work in this book. Cross through any work that you do not want to be marked.

Information
- The marks for questions are shown in brackets.
- The maximum mark for this paper is 100.

Advice
- Unless stated otherwise, you may quote formulae, without proof, from the booklet.
- You do not necessarily need to use all the space provided.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
TOTAL	

2

Answer **all** questions in the spaces provided.

1 Given that $y = \sin x + \sinh x$, find $\dfrac{\mathrm{d}^2 y}{\mathrm{d}x^2} + y$

Circle your answer.

[1 mark]

$$2\sin x \qquad\qquad -2\sin x \qquad\qquad 2\sinh x \qquad\qquad -2\sinh x$$

2 Which one of the expressions below is **not** equal to zero?

Circle your answer.

[1 mark]

$$\lim_{x \to \infty}(x^2 \mathrm{e}^{-x}) \qquad \lim_{x \to 0}(x^5 \ln x) \qquad \lim_{x \to \infty}\left(\dfrac{\mathrm{e}^x}{x^5}\right) \qquad \lim_{x \to 0}(x^3 \mathrm{e}^x)$$

3 The determinant $A = \begin{vmatrix} 1 & 1 & 1 \\ 2 & 0 & 2 \\ 3 & 2 & 1 \end{vmatrix}$

Which one of the determinants below has a value which is **not** equal to the value of A?

Tick (✓) **one** box.

[1 mark]

$\begin{vmatrix} 3 & 1 & 3 \\ 2 & 0 & 2 \\ 3 & 2 & 1 \end{vmatrix}$ ☐

$\begin{vmatrix} 1 & 2 & 3 \\ 1 & 0 & 2 \\ 1 & 2 & 1 \end{vmatrix}$ ☐

$\begin{vmatrix} 2 & 2 & 2 \\ 1 & 0 & 1 \\ 3 & 2 & 1 \end{vmatrix}$ ☐

$\begin{vmatrix} 1 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 0 & 2 \end{vmatrix}$ ☐

4 It is given that $f(x) = \cosh^{-1}(x - 3)$

Which of the sets listed below is the greatest possible domain of the function f?

Circle your answer.

[1 mark]

$\{x : x \geq 4\}$ $\qquad$ $\{x : x \geq 3\}$ $\qquad$ $\{x : x \geq 1\}$ $\qquad$ $\{x : x \geq 0\}$

5   Josh and Zoe are solving the following mathematics problem:

> The curve C_1 has equation
>
> $$\frac{x^2}{16} - \frac{y^2}{9} = 1$$
>
> The matrix $\mathbf{M} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ maps C_1 onto C_2
>
> Find the equations of the asymptotes of C_2

Josh says that to solve this problem you **must** first carry out the transformation on C_1 to find C_2, and then find the asymptotes of C_2

Zoe says that you will get the same answer if you first find the asymptotes of C_1, and then carry out the transformation on these asymptotes to obtain the asymptotes of C_2

Show that Zoe is correct.

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

6 (a) Express $-5 - 5i$ in the form $re^{i\theta}$, where $-\pi < \theta \le \pi$

[2 marks]

_____

_____

_____

_____

_____

_____

6 (b) The point on an Argand diagram that represents $-5 - 5i$ is one of the vertices of an equilateral triangle whose centre is at the origin.

Find the complex numbers represented by the other two vertices of the triangle.

Give your answers in the form $re^{i\theta}$, where $-\pi < \theta \le \pi$

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

Turn over ▶

7   Show that

$$\sum_{r=11}^{n+1} r^3 = \frac{1}{4}(n^2 + an + b)(n^2 + an + c)$$

where a, b and c are integers to be found.

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

8   **A** is a non-singular 2×2 matrix and $\mathbf{A}^{\mathrm{T}}$ is the transpose of **A**

8 (a)   Using the result

$$(\mathbf{AB})^{\mathrm{T}} = \mathbf{B}^{\mathrm{T}}\mathbf{A}^{\mathrm{T}}$$

show that

$$(\mathbf{A}^{-1})^{\mathrm{T}} = (\mathbf{A}^{\mathrm{T}})^{-1}$$

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

8 (b)   It is given that $\mathbf{A} = \begin{bmatrix} 4 & 5 \\ -1 & k \end{bmatrix}$, where k is a real constant.

8 (b) (i)   Find $(\mathbf{A}^{-1})^{\mathrm{T}}$, giving your answer in terms of k

[2 marks]

_____

_____

_____

_____

_____

8 (b) (ii)   State the restriction on the possible values of k

[1 mark]

_____

_____

_____

Turn over ▶

9 The complex number z is such that

$$z = \frac{1+i}{1-ki}$$

where k is a real number.

9 (a) Find the real part of z and the imaginary part of z, giving your answers in terms of k

[2 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

9 (b) In the case where $k = \sqrt{3}$, use part **(a)** to show that

$$\cos \frac{7\pi}{12} = \frac{\sqrt{2} - \sqrt{6}}{4}$$

[5 marks]

_____

_____

_____

_____

_____

_____

Turn over ▶

10 The region R on an Argand diagram satisfies both $|z + 2i| \leq 3$ and $-\dfrac{\pi}{6} \leq \arg(z) \leq \dfrac{\pi}{2}$

10 (a) Sketch R on the Argand diagram below.

[3 marks]

10 (b) Find the maximum value of $|z|$ in the region R, giving your answer in exact form.

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

Turn over for the next question

Turn over ▶

11 The line l_1 passes through the points $A(6, 2, 7)$ and $B(4, -3, 7)$

11 (a) Find a Cartesian equation of l_1

[2 marks]

_____

_____

_____

_____

_____

_____

11 (b) The line l_2 has vector equation $\mathbf{r} = \begin{bmatrix} 8 \\ 9 \\ c \end{bmatrix} + \mu \begin{bmatrix} 1 \\ 1 \\ 2 \end{bmatrix}$ where c is a constant.

11 (b) (i) Explain how you know that the lines l_1 and l_2 are not perpendicular.

[2 marks]

_____

_____

_____

_____

_____

_____

11 (b) (ii) The lines l_1 and l_2 both lie in the same plane.

Find the value of c

[5 marks]

Turn over for the next question

14

12 The function f is defined by

$$f(n) = 3^{3n+1} + 2^{3n+4} \qquad (n \in \mathbb{Z}^+)$$

Prove by induction that $f(n)$ is divisible by 19 for $n \geq 1$

[6 marks]

15

Turn over for the next question

13 The quadratic equation $z^2 - 5z + 8 = 0$ has roots α and β

13 (a) Write down the value of $\alpha + \beta$ and the value of $\alpha\beta$

[2 marks]

_____

_____

_____

_____

_____

_____

13 (b) Without finding the value of α or the value of β, show that $\alpha^4 + \beta^4 = -47$

[4 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

13 (c) Find a quadratic equation, with integer coefficients, which has roots $\alpha^3 + \beta$ and $\beta^3 + \alpha$

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

Turn over for the next question

14 The function f is defined by

$$f(x) = \frac{1}{4x^2 + 16x + 19} \qquad (x \in \mathbb{R})$$

14 (a) Show, without using calculus, that the graph of $y = f(x)$ has a stationary point at $\left(-2, \dfrac{1}{3}\right)$

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

14 (b) Show that $\displaystyle\int_{-2}^{-\frac{1}{2}} f(x)\, dx = \frac{\pi\sqrt{3}}{18}$

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

14 (c)   Find the value of $\displaystyle\int_{-2}^{\infty} f(x)\,dx$

Fully justify your answer.

[2 marks]

15 (a) Given that $z = \cos\theta + i\sin\theta$, use de Moivre's theorem to show that

$$z^n - z^{-n} = 2i\sin n\theta$$

[2 marks]

_____

_____

_____

_____

_____

_____

15 (b) The series S is defined as

$$S = \sin\theta + \sin 3\theta + ... + \sin(2n-1)\theta$$

Use part **(a)** to express S in the form

$$S = \frac{1}{2i}(G_1) - \frac{1}{2i}(G_2)$$

where each of G_1 and G_2 is a geometric series.

[3 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

15 (c)   Hence, show that

$$S = \frac{\sin^2(n\theta)}{\sin\theta}$$

[5 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

Turn over ▶

22

16   A bungee jumper of mass m kg is attached to an elastic rope.
The other end of the rope is attached to a fixed point.

The bungee jumper falls vertically from the fixed point.

At time t seconds after the rope first becomes taut, the extension of the rope is x metres and the speed of the bungee jumper is v m s^{-1}

16 (a)   A model for the motion while the rope remains taut assumes that the forces acting on the bungee jumper are

- the weight of the bungee jumper

- a tension in the rope of magnitude kx newtons

- an air resistance force of magnitude Rv newtons

where k and R are constants such that $4km > R^2$

16 (a) (i)   Show that this model gives the result

$$x = e^{-\frac{Rt}{2m}}\left(A\cos\left(\frac{\sqrt{4km - R^2}}{2m}\right)t + B\sin\left(\frac{\sqrt{4km - R^2}}{2m}\right)t\right) + \frac{mg}{k}$$

where A and B are constants, and g m s^{-2} is the acceleration due to gravity.

You do not need to find the value of A or the value of B

[6 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

16 (a) (ii) It is also given that:

$$k = 16$$

$$R = 20$$

$$m = 62.5$$

$$g = 9.8 \, \text{m s}^{-2}$$

and that the speed of the bungee jumper when the rope becomes taut is $14 \, \text{m s}^{-1}$

Show that, to the nearest integer, $A = -38$ and $B = 16$

[6 marks]

Turn over ▶

Do not write outside the box

16 (b)   A second, simpler model assumes that the air resistance is zero.

The values of k, m and g remain the same.

Find an expression for x in terms of t according to this simpler model, giving the values of all constants to two significant figures.

[4 marks]

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

_____

END OF QUESTIONS

Do not write outside the box

There are no questions printed on this page

DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED

27

Question number	Additional page, if required. Write the question numbers in the left-hand margin.

Do not write outside the box

Question number	**Additional page, if required.** **Write the question numbers in the left-hand margin.**