

OCR


Oxford Cambridge and RSA

Monday 10 June 2024 – Morning**A Level Chemistry B (Salters)****H433/01 Fundamentals of Chemistry****Time allowed: 2 hours 15 minutes****You must have:**

- the Data Sheet for Chemistry B

You can use:

- a scientific or graphical calculator
- an HB pencil

Please write clearly in black ink. **Do not write in the barcodes.**

Centre number

| | | | | |
|--|--|--|--|--|
| | | | | |
|--|--|--|--|--|

Candidate number

| | | | |
|--|--|--|--|
| | | | |
|--|--|--|--|

First name(s) _____

Last name _____

INSTRUCTIONS

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- Answer **all** the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

INFORMATION

- The total mark for this paper is **110**.
- The marks for each question are shown in brackets [].
- Quality of extended response will be assessed in questions marked with an asterisk (*).
- This document has **36** pages.

ADVICE

- Read each question carefully before you start your answer.

2

Section A

You should spend a **maximum** of **40 minutes** on this section.

Write your answer to each question in the box provided.

- 1 An isotope of an element has a mass number of 25.

What is correct for an atom of this isotope?

- A It has 12 neutrons.
- B It has 25 electrons.
- C It has 25 protons.
- D The sum of its protons and neutrons is 25.

Your answer

[1]

- 2 What is correct about s orbitals?

- A Only elements in Group 1 or Group 2 have them.
- B They always contain 2 electrons.
- C They are spherical in shape.
- D They can contain up to 6 electrons.

Your answer

[1]

- 3 Which species have an electron configuration ending in s^1 ?

- A atoms of some d-block elements
- B atoms of Group 13
- C ions of Group 2
- D ions of Group 16

Your answer

[1]

3

4 Which row has the radiations arranged in order of **increasing** wavelength?

- A infrared < ultraviolet < visible
- B infrared < visible < ultraviolet
- C ultraviolet < infrared < visible
- D ultraviolet < visible < infrared

Your answer

[1]

5 What is correct about nuclear fusion reactions?

- A They are the way in which all elements are formed.
- B They join together atoms to make heavier atoms.
- C They only occur under conditions of high temperature and pressure.
- D They turn neutrons into protons.

Your answer

[1]

6 Which species is present in an acidic solution of glycine, $\text{H}_2\text{NCH}_2\text{COOH}$?

- A $\text{H}_3\text{N}^+\text{CH}_2\text{COOH}$
- B $\text{H}_3\text{N}^+\text{CH}_2\text{COO}^-$
- C $\text{H}_2\text{NCH}_2\text{COO}^-$
- D $\text{H}_2\text{NCH}_2\text{COOH}_2^+$

Your answer

[1]

4

7 In the mass spectrum of ethanoic acid ($M_r = 60$), what is the cause of the very small peak at $m/z = 61$?

- A the presence of an ethanol impurity in the ethanoic acid
- B the presence of small amounts of $\text{CH}_3\text{COOH}_2^+$
- C the presence of ^{13}C in the ethanoic acid
- D the presence of ^1H in the ethanoic acid

Your answer

[1]

8 Which compound has optical isomers?

- A $\text{CH}_3\text{CH}_2\text{COOH}$
- B $\text{CH}_3\text{CH}_2\text{CONH}_2$
- C $\text{H}_2\text{NCH}_2\text{COOH}$
- D $\text{H}_2\text{NCH}(\text{CH}_3)\text{COOH}$

Your answer

[1]

9 Which compound will react with sodium carbonate?

- A CH_3COCH_3
- B $\text{CH}_3\text{COOCH}_3$
- C $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$
- D $\text{CH}_3\text{CH}_2\text{COOH}$

Your answer

[1]


10 Which option occurred first in the discovery of atomic structure?

- A the Geiger-Marsden experiment
- B the idea of an atom with electrons embedded in positive charge
- C the idea that electrons are arranged in shells
- D the idea that the nucleus consists of protons and neutrons

Your answer

[1]

11 The mass spectrum of a sample of an element is shown:



What is the relative atomic mass of the sample?

- A 69.2
- B 69.5
- C 69.8
- D 70.0

Your answer

[1]

6

12 Which of the following is correct for a C=C bond?

- A All the electrons are between the C atoms.
- B There are two π -bonds.
- C There is one σ -bond.
- D The bond can freely rotate.

Your answer

[1]

13 The Table shows the concentrations of some noble gases in the atmosphere.

| Gas | Concentration by volume |
|-----|-------------------------|
| He | 5 ppm |
| Ne | 2 parts in 10^5 |
| Kr | 0.0001% |
| Xe | 1 part in 10^7 |

Which gas has the **highest** concentration?

- A He
- B Ne
- C Kr
- D Xe

Your answer

[1]

14 Which reaction has the **smallest** atom economy for the formation of the organic product?

- A $\text{C}_2\text{H}_4 + \text{HCl} \rightarrow \text{C}_2\text{H}_5\text{Cl}$
B $\text{C}_2\text{H}_4 + \text{HBr} \rightarrow \text{C}_2\text{H}_5\text{Br}$
C $\text{C}_2\text{H}_6 + \text{Cl}_2 \rightarrow \text{C}_2\text{H}_5\text{Cl} + \text{HCl}$
D $\text{C}_2\text{H}_6 + \text{Br}_2 \rightarrow \text{C}_2\text{H}_5\text{Br} + \text{HBr}$

Your answer

[1]

15 Which of 1-iodobutane and 1-chlorobutane has the higher boiling point and why?

- A 1-Chlorobutane because its molecules have the stronger permanent dipole–dipole bonds.
B 1-Chlorobutane because its molecules have the stronger instantaneous dipole–induced dipole bonds.
C 1-Iodobutane because its molecules have the stronger permanent dipole–dipole bonds.
D 1-Iodobutane because its molecules have the stronger instantaneous dipole–induced dipole bonds.

Your answer

[1]

16 Which reaction has an increase in entropy?

- A $\text{H}_2\text{O}(\text{g}) \rightarrow \text{H}_2\text{O}(\text{l})$
B $\text{N}_2\text{O}_4(\text{g}) \rightarrow 2\text{NO}_2(\text{g})$
C $2\text{SO}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{SO}_3(\text{g})$
D $\text{C}(\text{CH}_3)_4(\text{g}) \rightarrow \text{CH}_3(\text{CH}_2)_3\text{CH}_3(\text{l})$

Your answer

[1]

17 The compounds below have negative lattice enthalpies.

Which option shows the compounds in order of more negative lattice enthalpies from left to right?

- A NaF, KF, RbF
- B Na₂O, K₂O, Rb₂O
- C Na₂O, MgO, Al₂O₃
- D NaF, NaCl, NaBr

Your answer

[1]

18 A colorimeter is used to determine the concentration of copper(II) ions in solution.

What is correct?

- A Absorbance increases with the concentration of copper(II) ions.
- B In the colorimeter the filter is positioned between the cuvette and the detector.
- C The filter should be blue because copper(II) ions are blue.
- D The zero reading on the colorimeter is taken using an empty cuvette.

Your answer

[1]

19 What is correct about propanal, CH₃CH₂CHO, and propanone, CH₃COCH₃?

- A Only propanal can be oxidised by acidified dichromate.
- B Only propanal forms a 'silver mirror' with Fehling's solution.
- C Only propanone forms a cyanohydrin with hydrogen cyanide.
- D Propanal has a higher boiling point than propanone because propanal can form hydrogen bonds.

Your answer

[1]

- 20 A student is measuring the relative rates of reaction of 1-chlorobutane, 1-bromobutane and 1-iodobutane with aqueous silver nitrate.

Which statement is correct?

- A 1-Chlorobutane is slowest and gives a white precipitate.
- B 1-Bromobutane is slowest and gives a yellow precipitate.
- C 1-Iodobutane is fastest and gives a white precipitate.
- D 1-Iodobutane is slowest and gives a yellow precipitate.

Your answer

[1]

- 21 Wood is used to make a sample of bioethanol. The bioethanol is used to fuel a car engine.

What is correct?

- A Carbon dioxide is the only pollutant produced by the car engine.
- B The amount of carbon dioxide given out when the fuel is fully burned is equal to the amount of carbon dioxide absorbed when the wood is grown.
- C The fuel is made from the wood by cracking.
- D The wood is never grown where other crops could be planted.

Your answer

[1]

- 22 Which of the following represents poly(propene)?

- A $-\text{CH}_2-\text{CH}(\text{CH}_3)-$
- B $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$
- C $-\text{CH}=\text{C}(\text{CH}_3)-\text{CH}_2-$
- D $-\text{C}(\text{CH}_3)-\text{C}(\text{CH}_3)-$

Your answer

[1]

23 10 cm³ of an alkane react exactly with 35 cm³ oxygen in complete combustion at RTP.

What is the alkane?

- A CH₄
- B C₂H₆
- C C₃H₈
- D C₄H₁₀

Your answer

[1]

24 50 cm³ of 0.30 mol dm⁻³ Fe(II) in acid solution exactly reduces 25 cm³ of 0.10 mol dm⁻³ NaClO₃ solution. Fe(III) is formed.

What is the oxidation state of chlorine in the product?

- A +3
- B +2
- C 0
- D -1

Your answer

[1]

25 Methane reacts with chlorine by a radical mechanism.

There is an initiation reaction, two propagation reactions and various termination reactions.

The overall equation for the main reaction is $\text{CH}_4 + \text{Cl}_2 \rightarrow \text{CH}_3\text{Cl} + \text{HCl}$

Which statement is correct for this reaction?

- A CH_3Cl is formed in the first propagation reaction.
- B HCl is formed in the second propagation reaction.
- C One termination reaction involves the formation of ethane.
- D Two radicals react in the first propagation step.

Your answer

[1]

26 The solubility of lead(II) chloride is $0.0388 \text{ mol dm}^{-3}$ at 293 K.


What is the correct numerical value of the solubility product at this temperature?

- A 5.84×10^{-5}
- B 2.34×10^{-4}
- C 1.51×10^{-3}
- D 3.01×10^{-3}

Your answer

[1]

- 29 The graph shows how the rate of an enzyme catalysed reaction varies with substrate concentration.




At low substrate concentration, which of the following is/are correct?

- 1 Not all enzyme active sites have a substrate molecule attached.
 - 2 The reaction is first order with respect to substrate.
 - 3 The rate determining step involves both enzyme and substrate.
- A 1, 2 and 3
B Only 1 and 2
C Only 2 and 3
D Only 1

Your answer

[1]

30 A carbon atom has the bonding shown.



Which of the following is/are correct for this carbon atom?

- 1 The shape is trigonal around the carbon atom.
 - 2 The atoms attached to the bonds are all in the same plane.
 - 3 The angle between the single bonds is 109° .
- A 1, 2 and 3
B Only 1 and 2
C Only 2 and 3
D Only 1

Your answer

[1]

15
BLANK PAGE

DO NOT WRITE ON THIS PAGE
Turn over for Section B

Section B

31 Fireworks contain metal compounds that produce intense colours when they are burned.

(a) Give the colour from a firework containing barium carbonate.

..... [1]

(b) The atomic **emission** spectrum of barium appears as a series of coloured lines at specific frequencies on a black background.

Describe the appearance of the atomic **absorption** spectrum of barium.

How is the absorption spectrum related to the emission spectrum?

.....
.....
.....
..... [2]

(c) One line in the barium spectrum has a wavelength of 5.54×10^{-5} cm.

Calculate the frequency of this line (in Hz).

frequency = Hz [2]

(d) Strontium carbonate is another compound used in fireworks.

(i) Write the equation for the thermal decomposition of SrCO_3 .

[1]

(ii) There is a trend in the stability of the Group 2 carbonates.

A student says:

- Barium carbonate decomposes at a higher temperature than strontium carbonate.
- This is because the carbonate ion has a stronger bond to the more reactive barium ion.

Comment on the student's statements giving the correct chemistry where necessary.

.....

.....

.....

.....

.....

.....

.....

.....

.....


..... [4]

(e) A group of students identify an insoluble Group 2 carbonate, MCO_3 , by calculating its M_r .

The students' procedure and results are:

- They weigh out 3.02 g of MCO_3 .
- They add the MCO_3 to 50.0 cm³ of 2.00 mol dm⁻³ HCl (an excess).
- When the reaction has finished they make the solution up to 250 cm³ in a volumetric flask.
- They titrate 25.0 cm³ of the diluted solution with 0.108 mol dm⁻³ NaOH.
- Their mean titre is 26.15 cm³.

The reaction between HCl and NaOH is:



The reaction between MCO_3 and HCl is:



(i) Describe the procedure they should use for weighing out the 3.02 g of MCO_3 .

.....

.....

.....

..... [2]

(ii) Calculate the M_r of MCO_3 .

$M_r = \dots\dots\dots$ [5]

19

(iii) The 25.0 cm^3 of the solution used for the titration is measured using a volumetric pipette.

The uncertainty in reading the volumetric pipette is 0.06 cm^3 .

Calculate the percentage uncertainty in using this pipette.


Give your answer to an **appropriate** number of significant figures.

percentage uncertainty = % [1]

21
BLANK PAGE

DO NOT WRITE ON THIS PAGE
Question 32(b) begins on page 22

(b) Another reaction that occurs in a catalytic converter is shown in **Equation 32.2**.



(i) The rate equation for this reaction is:

$$\text{Rate} = k[\text{NO}_2]^2$$

A student says that this reaction takes place in a single step.

Explain why this cannot be correct and give a possible mechanism for the reaction, indicating the rate-determining step.

.....

.....

.....

.....


.....

.....

..... [3]

- (ii) A chemist studies the reaction in **Equation 32.2** and measures the rate constant, k , at different temperatures.

The chemist plots $\ln k$ against $1/T$ on a graph as shown.



Calculate the activation enthalpy (in kJ mol^{-1}) for the reaction in **Equation 32.2**.

activation enthalpy = kJ mol^{-1} [4]

33 Ammonium nitrate, NH_4NO_3 , is a fertiliser.

One of the steps in the manufacture of NH_4NO_3 is the oxidation of NO by the reaction in **Equation 33.1**.



(a) Give the appearance of $\text{NO}_2(\text{g})$.

..... [1]

(b) Give the test and its result that shows that a solid contains nitrate ions.

.....
.....
.....
..... [2]

(c) The reaction in **Equation 33.1** is an example of a chemical equilibrium.

(i) A student says that once the position of equilibrium has been reached, the reaction has stopped so the concentration of $\text{NO}_2(\text{g})$ remains constant.

Evaluate what the student says and give correct chemistry where necessary.

.....
.....
.....
..... [2]

(ii) For the reaction in **Equation 33.1** at a certain temperature,

$$K_c = 6.9 \times 10^2 \text{ dm}^3 \text{ mol}^{-1}$$

In an industrial plant, operating at that temperature:

Equilibrium concentration, $[\text{NO}(\text{g})] = 0.250 \text{ mol dm}^{-3}$

Equilibrium concentration, $[\text{O}_2(\text{g})] = 0.180 \text{ mol dm}^{-3}$

Write the expression for K_c and calculate the equilibrium concentration of $\text{NO}_2(\text{g})$.

equilibrium concentration, $[\text{NO}_2(\text{g})] = \dots\dots\dots \text{ mol dm}^{-3}$ [3]

(iii) Explain how increasing temperature affects the value of K_c for the reaction in **Equation 33.1**.

.....

.....

.....

..... [2]

Equation 33.1 is repeated: $2\text{NO}(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g}) \quad \Delta H = -114 \text{ kJ mol}^{-1}$ **Equation 33.1**


(d) The reaction in **Equation 33.1** uses a heterogeneous catalyst.

(i) Explain the meaning of the term **heterogeneous** here.

.....
..... [1]

(ii) Draw an enthalpy profile diagram for the reaction in **Equation 33.1** and label the catalysed and uncatalysed reactions.

Show the reactants and products and the activation enthalpies. [2]



(iii) The catalyst used in the reaction in **Equation 33.1** is a solid mixture of platinum and rhodium.

Explain why this solid is able to behave as a catalyst.

.....
.....
.....
..... [2]

(e) The standard entropies of the substances involved in **Equation 33.1** are shown in the table.

| Substance | $S^\circ/\text{J mol}^{-1}\text{K}^{-1}$ |
|-------------------------|--|
| $\text{O}_2(\text{g})$ | 205.0 |
| $\text{NO}(\text{g})$ | 210.7 |
| $\text{NO}_2(\text{g})$ | 240.0 |

Calculate the total entropy change, $\Delta_{\text{tot}}S$, (in $\text{J mol}^{-1}\text{K}^{-1}$) for the reaction in **Equation 33.1** at 50°C .

$$\Delta_{\text{tot}}S = \dots\dots\dots \text{J mol}^{-1}\text{K}^{-1} \text{ [4]}$$

34 The plastic welding agent butanone, $\text{CH}_3\text{CH}_2\text{COCH}_3$, is sold as “model cement” and used for joining the plastic parts of scale model kits.

(a) Butanone can be formed by the oxidation of the alcohol $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$.

Classify this alcohol as primary, secondary or tertiary.

Give the structural reason for the classification you have chosen.

Classification:

Reason:

..... [2]

(b) There are other alcohols that are structural isomers of $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$.

Draw the **skeletal** formulae for the **two branched-chain** isomers.

| | |
|--|--|
| | |
|--|--|

[2]

(c) The alcohol $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$ can be made by reacting $\text{CH}_3\text{CH}_2\text{CHBrCH}_3$ with aqueous sodium hydroxide.

Name the mechanism of this reaction.

..... [1]

(d) Alcohols can be dehydrated by heating with concentrated sulfuric acid.

When dehydrated, $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$ produces a mixture of three isomeric products.


Draw the structures of these **three** isomeric products.

| | | |
|--|--|--|
| | | |
|--|--|--|


[3]

(e) The mechanism for the dehydration of a cyclic alcohol using concentrated sulfuric acid takes place in two steps.

Step 1:



Step 2:






Add curly arrows to **Step 2** to show the electron movements in the mechanism.

[2]


(f)* A chemist sets out to identify **compound A** and its oxidation product, **compound B**.

The chemist has the mass spectrum, the IR spectrum and the ^1H NMR spectrum of **compound A**, and the IR spectrum of **compound B**.



You may do working on this page but it will not be marked.

35 Acid orange is a soluble azo dye that can be made from benzene by the following route:



(a)

(i) Name **compound C**.

..... [1]

(ii) **Reaction 1** is an electrophilic substitution using concentrated sulfuric acid.

Give the formula of the electrophile.

..... [1]

(b) Use the Data Sheet to give reagents and conditions for **reaction 3**.

..... [1]

(c) **Reaction 4** forms a diazonium salt.

Give the reagents and conditions for this reaction.

Reagents:

.....

Conditions:

..... [3]


(d) **Reaction 5** is a coupling reaction, using a phenol and sodium hydroxide.

Give the formula of the phenol.

[1]

(e)

(i) Draw a **ring** around the **chromophore** of acid orange.



[1]

(ii) One part of its structure makes acid orange more soluble than many other azo dyes.

Give the formula of this part.

..... [1]

Extra answer space if required.

.....

.....

.....

.....

.....

.....

.....

.....

END OF QUESTION PAPER

EXTRA ANSWER SPACE

If you need extra space use this lined page. You must write the question numbers clearly in the margin.

A large area of the page is reserved for writing answers. It is bounded on the left by a solid vertical line and on the top, bottom, and right by horizontal dotted lines. This creates a grid of approximately 20 rows for writing.

OCR

Oxford Cambridge and RSA

Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact The OCR Copyright Team, The Triangle Building, Shaftesbury Road, Cambridge CB2 8EA.

OCR is part of Cambridge University Press & Assessment, which is itself a department of the University of Cambridge.